

ACCEPTED
OCT 28 2002

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2724-275

[] indicates optional statements

PROPOXUR FLEA COLLAR FOR CATS RF-101

Long Lasting	Kills Fleas Fast
Fast Acting Long Lasting Protection	Fast Acting for Fleas
Works From Head to Tail	Quick Killing for Fleas
Rain or Shine	Quick Acting for Fleas
[Available through Veterinarians]	Fast Action for Fleas
[Step 2] Part of the [Brand name] [#] Step Pet Protection [Guarantee] Program	
Temporary wetting will not reduce effectiveness	[Snip notch at replacement month]
[Unique Dating Feature]	
5 month protection	
Kills Fleas and Ticks including Lyme Disease-Carrying Ticks	
Kills fleas and ticks up to 5 months [21 weeks][150 days]	
For cats of all sizes	
[Unique Breakaway Feature – graphic]	
Works from Head to Tail to Kill Fleas and Lyme Disease-Carrying Ticks up to 5 months	
[(Descriptor) red/(Descriptor) blue/(Descriptor) green/(Descriptor) clear/ (Descriptor) white/(Descriptor) purple/(Descriptor) pink/(Descriptor) yellow]	
[Unique one-piece buckle design][Integrated one-piece buckle]	
[Integrated plastic buckle] [Breakaway Collar][Expandable Collar][Stretch Band]	
Kills [deer] ticks (including <i>Ixodes scapularis</i>), which may transmit Lyme disease [<i>borreliosis</i>]	
Effective against [vectors, ticks, <i>Ixodes scapularis</i>] which may transmit Lyme disease [<i>borreliosis</i>]	
Controls [Kills] Lyme disease [<i>borreliosis</i>]-carrying ticks	
100% satisfaction guaranteed/guaranteed or your money back	

ACTIVE INGREDIENT:

Propoxur (CAS #114-26-1) 10.0%

OTHER INGREDIENTS:..... 90.0%

Total: 100.0%

DO NOT ALLOW CHILDREN TO PLAY WITH THIS COLLAR

CAUTION

See Precautions on back

EPA Reg. No. 2724-275

EPA Est. No.

NET CONTENTS

**READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON CATS
[Apply only one collar at a time]**

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. Buckle collar around cat's neck. [Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Since the plastic buckle is molded into the collar, the possibility of neck irritation and buckle failure is reduced.] Fit collar snug enough to prevent placement of the cat's leg or lower jaw through the collar, yet loose enough to be rotated about the neck.
2. Leave 1-2 extra inches on the collar for adjustment, cut off excess length and dispose of in trash.
3. Check periodically and adjust fit if necessary, especially when kittens are rapidly growing.
4. May be used in addition to regular collar. Replace collar when effectiveness diminishes. [To release the collar, squeeze buckle between thumb and finger and pull sharply.]

[for "breakaway" versions]

[UNIQUE FEATURE: This collar is designed to stretch under the weight of an average size adult cat. It helps the cat to free itself ("breakaway") if the collar gets snagged on a fixed object. If the collar is fitted improperly, so that it is loose enough for a cat to work the collar over its leg or jaw, this release feature may not stretch adequately to allow the extra room necessary for the cat to free itself.]

STORAGE and DISPOSAL: **Storage:** Store in original unopened container away from children. Do not reuse container or reuse collar. **Disposal:** Dispose of in trash.

WHAT TO EXPECT:

For Fleas:

1. This [brand name] collar is designed to kill adult fleas.
2. This collar starts killing fleas as soon as it is placed around the cat's neck.
3. Protection against fleas will be achieved within a few days, providing continuous protection for 5 months.
4. Replace collar every 5 months to help control [prevent] flea infestations on your pet and in your home.

For Ticks:

1. Ticks are tougher to kill than fleas and are killed more slowly.
2. Full protection against adult and immature ticks will be achieved within a few days, providing continuous protection for 5 months.
3. Adult ticks will be killed over the entire pet. Dead ticks will fall off or may be easily removed.
4. Replace collar every 5 months for continuous protection from ticks.

PRECAUTIONARY STATEMENTS


HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION: Do not get dust or collar in mouth, harmful if swallowed. Do not get dust in eyes, will cause temporary pupillary constriction.* The dust released by this collar is a cholinesterase inhibitor.

Humans: Do not open package until ready to use. Do not allow children to play with this collar. Dust will form on this collar during storage. Wash hands thoroughly with soap and water after handling collar.

Animals: Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing cats. Do not use on kittens under 12 weeks of age. Do not use this product on cats simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. However, flea and tick collars may be immediately replaced. When collar is first worn, observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amounts of water.

FIRST AID	
Call a poison control center or doctor immediately for treatment advice.	
If swallowed	<ul style="list-style-type: none"> • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-[phone number] for emergency medical treatment information.	
NOTE TO PHYSICIAN/VETERINARIAN: Contains an N-methyl carbamate that inhibits cholinesterase. *Atropine and *homatropine are antidotal only if symptoms of cholinesterase inhibition are present.	

[Note: First Aid statement paragraph format may be used if space is limited.]



Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

Wellmark International
Schaumburg, IL 60173

In case of emergency or for product use information, call [phone number] or visit our Web site:
[URL]

For information, or in emergency call [phone number]

[If not satisfied, send used collar and indication of purchase price paid to (company name/address). See top panel of box for details of unconditional guarantee.]

Made in USA