

ACCEPTED  
All...  
2-4-72

# ZOECON RF-156 COLLAR FOR DOGS

(Controls or Kills) fleas up to 7 months  
(Controls or Kills) ticks up to 7 months  
(Aids in control or aids in kill) of fleas and ticks for an additional 3 months  
Aids in the prevention of Sarcoptic Mange for up to 8 weeks.

ACTIVE INGREDIENT:  
N-(mercaptomethyl) phthalimide S-(0,0-dimethyl phosphorodithioate) . . . . . 15.0%  
INERT INGREDIENTS . . . . . 85.0%

Net Wt. 1oz (28.3g)  
Keep Out of Reach of Children  
**CAUTION**  
See other precautions on back.

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS - CAUTION** Dust will form on this collar during storage. Do not get dust on collar in mouth, or allow dust in eyes. Do not open protective pouch until ready to use. Do not allow children to handle collar.

**STATEMENT OF PRACTICAL TREATMENT:** If dust is on skin - Wash promptly with soap and water. Rinse thoroughly. If dust is in eyes - Rinse eyes with plenty of water. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** The dust released by this collar is a cholinesterase inhibitor. If signs of cholinesterase inhibition appear, Atropine is antidotal.

**DIRECTIONS FOR USE:** It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Buckle collar around dog's neck. Collar should be worn loosely to prevent irritation. Attach collar so two or three fingers may be inserted easily between collar and dog's neck. Cut off any excess length and dispose of it. When collar is first worn, observe neck area every few days for irritation. Remove collar at first sign of irritation or adverse reaction. Collar is intended for use only as an insecticide generator and is not to be taken internally by man or animals. Use in addition to regular collar. Do not use on sick or convalescing dogs. Do not use other pesticides on dogs while collar is worn. Product not to be used on puppies under 12 weeks of age. Remove collar when bathing.

The collar starts killing fleas as soon as it is placed around dog's neck. Fleas on the dog will be killed and new ones, which may temporarily appear on the dog, will also be killed while collar is worn. This product has been shown to provide effective flea (control or kill) up to 7 months plus aiding in (control or kill) for an additional 3 months. Replace collar when effectiveness diminishes.

Ticks are tough and are killed slowly. When the collar is first placed on dog, adult ticks will be killed over entire body in a few days and will fall off, or may then be easily removed. Ticks appear on dogs in three stages. In the first two stages, ticks are smaller than a match head and are difficult to see. The collar will also (control or kill) these immature stages. This product has been shown to provide effective tick (control or kill) up to 7 months plus aiding in (control or kill) for an additional 3 months. If ticks are a problem, this collar should be worn continuously. Kills ticks that may carry and transmit Rocky Mountain spotted fever and tularemia. Replace when effectiveness diminishes. The collar will also aid in the prevention of Sarcoptic Mange for up to 8 weeks.

**DISPOSAL:** Do not reuse empty container, pouch or used collar. Dispose in trash collection.

ZOECON INDUSTRIES  
12200 Denton Drive Dallas, Texas 75236  
EPA Reg. No. 2724-279 EPA Est. 2724-TX-1

RB7-A