

2764-282

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: IRRITANT. Do not burn, puncture, or  
until ready to use. Do not allow children to handle this collar. Wash hands  
thoroughly with soap and water after handling the collar. Do not use in areas  
containing dust. Do not use other insecticides on dogs while the collar is worn.  
Do not use on dogs under 10 weeks of age. This product contains a  
cholinesterase inhibitor. Some breeds of individuals may be sensitive. Remove  
collar if symptoms of cholinesterase inhibition may include salivation,  
vomiting, and/or diarrhea occur. Atropa belladonna only if symptoms of  
cholinesterase inhibition are present.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a  
manner inconsistent with its labeling.

Buckle collar around dog's neck. Collar should be worn loosely to prevent  
irritation. Attach collar so two or three fingers may be inserted easily  
between collar and dog's neck. Cut off any excess length in excess of 1/2".  
When collar is first worn, observe neck area every few days for irritation.  
Remove collar at the first sign of irritation or adverse reaction.

This collar starts killing fleas and ticks as soon as it is placed on the dog's  
neck. Fleas on the dog will be killed within 48 hours after collar  
attachment, and new ones which may temporarily appear on the dog will also be  
killed while collar is worn.

Ticks are tough and are killed slowly. When the collar is first placed on dog,  
adult ticks will be killed over entire body in a few days and will fall off or  
may then be easily removed. Ticks appear on dogs in three stages. In the first  
two stages ticks are smaller than a match head and are difficult to see. The  
collar will also control these mature stages in addition to killing  
collar.

This collar has been shown to provide effective flea control for up to 11  
months and tick control for up to 7 months plus activity in ticks control for  
killing for an additional 2 months. Re-evaluate effectiveness from time to time.

Kills ticks that may carry and transmit Rocky Mountain spotted fever and  
tularemia.

This collar will also aid in the prevention of Sarcoptic Mange for up to 5  
months.

DISPOSAL: Do not reuse container or used collars; wrap and put in trash  
collection.

Zoecon Industries  
1021 Canton Drive, Dallas, Texas 75204

EPA reg. no.  
E 17962-1000  
Made in USA  
1021

ZOECON

# COLLAR RF-159 FOR DOGS

Fits necks to 19"  
Fits necks to 19"  
Fits necks to 25"  
Fits necks to 25"

Long Lasting  
Long Lasting Protection

(Controls or kills) Fleas Up To 11 Months

(Controls or kills) Ticks Up To 7 Months

Aids In Tick Control For An Additional 2 Months

From Head To Tail

Aids In The Prevention of Sarcoptic Mange  
For Up To 5 Months

ACTIVE INGREDIENT:

Chlorpyrifos: 0,0-diethyl 0-(3,5,6-  
trichloro-pyridyl) phosphorothioate..... 8.0%  
Inert Ingredients..... 92.0%

Net Weight: 1 oz. (28.4 g)  
1.3 oz. (36.9 g)

Keep Out of Reach of Children

**CAUTION**

See other precautions on back