

# ZOECON COLLAR RF-159 FOR DOGS

Long Lasting  
Long Lasting Protection

With necks to 19"  
Fits necks to 19"  
With necks to 25"  
Fits necks to 25"

(Controls or Kills) Fleas Up To 11 Months

(Controls of Kills) Ticks Up To 7 Months

(Controls or Kills) Fleas and Ticks Up to 7 Months

Aids In Tick Control For An Additional 2 Months

From Head To Tail

Aids In The Prevention of Sarcoptic Mange  
For Up To 5 Months

**ACTIVE INGREDIENT:**

Chlorpyrifos: 0,0-Diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate..... 8.0%

**INERT INGREDIENTS:**..... 92.0%

**NET WEIGHT:** 1 oz. (28.4 g)  
1.3 oz. (36.9 g)

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Other Precautions On Back



RECEIVED  
2724-282  
JUL 27 1984

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION:** Do not open protective pouch until ready to use. Do not allow children to handle this collar. Wash hands thoroughly with soap and water after handling collar. Do not use on sick and convalescing dogs. Do not use other pesticides on dogs while collar is worn. Do not use on dogs under 12 weeks of age. This product contains a cholinesterase inhibitor. Some breeds or individuals may be sensitive. Remove collar if symptoms of cholinesterase inhibition (which may include salivation, vomiting, and/or diarrhea) occur. Atropine is antidotal only if symptoms of cholinesterase inhibition are present.

**DIRECTIONS FOR USE -** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Buckle collar around dog's neck. Collar should be worn loosely to prevent irritation. Attach collar so two or three fingers may be inserted easily between collar and dog's neck. Cut off any excess length and dispose of it. When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction.

This collar starts killing fleas and ticks as soon as it is placed around dog's neck. Fleas on the dog will be killed (within 4-8 hours after collar attachment) and new ones which may temporarily appear on the dog will also be killed while collar is worn.

Ticks are tough and are killed slowly. When the collar is first placed on dog, adult ticks will be killed over entire body in a few days and will fall off or may then be easily removed. Ticks appear on dogs in three stages. In the first two stages ticks are smaller than a match head and are difficult to see. The collar will also control these immature stages. Use in addition to regular collar.

This collar has been shown to provide effective flea (control or kill) up to 11 months and tick (control or kill) up to 7 months plus aiding in tick (control or kill) for an additional 2 months. Replace when effectiveness diminishes.

Kills ticks that may carry and transmit Rocky Mountain spotted fever and tularemia.

This collar will also aid in the prevention of Sarcoptic Mange for up to 5 months.

**STORAGE & DISPOSAL:**

Store in original unopened container; away from children. Do not reuse container or used collar; wrap and put in trash.

Zoecon Industries  
A Division of Zoecon Corporation  
12200 Denton Drive, Dallas, Texas 75234

EPA Reg. No. 2724-282

EPA Est.  
c 1982 Zoecon  
Made in USA  
684-B-2807A-0089A

BEST DOCUMENT AVAILABLE

# ZOECON COLLAR RF-159 FOR DOGS

Long Lasting  
Long Lasting Protection

With necks to 19"  
Fits necks to 19"  
With necks to 25"  
Fits necks to 25"

(Controls or Kills) Fleas Up To 11 Months

(Controls or Kills) Ticks Up To 7 Months

(Controls or Kills) Fleas and Ticks Up to 7 Months

Aids In Tick Control For An Additional 2 Months

From Head To Tail

Aids In The Prevention of Sarcoptic Mange  
For Up To 5 Months

**ACTIVE INGREDIENT:**

Chlorpyrifos: 0,0-Diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate..... 8.0%


**INERT INGREDIENTS:**..... 92.0%

**NET WEIGHT:** 1 oz. (28.4 g)  
1.3 oz. (36.9 g)

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Other Precautions On Back



BEST DOCUMENT AVAILABLE

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION:** Do not open protective pouch until ready to use. Do not allow children to handle this collar. Wash hands thoroughly with soap and water after handling collar. Do not use on sick and convalescing dogs. Do not use other pesticides on dogs while collar is worn. Do not use on dogs under 12 weeks of age. This product contains a cholinesterase inhibitor. Some breeds or individuals may be sensitive. Remove collar if symptoms of cholinesterase inhibition (which may include salivation, vomiting, and/or diarrhea) occur. Atropine is antidotal only if symptoms of cholinesterase inhibition are present.

**DIRECTIONS FOR USE -** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Buckle collar around dog's neck. Collar should be worn loosely to prevent irritation. Attach collar so two or three fingers may be inserted easily between collar and dog's neck. Cut off any excess length and dispose of it. When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction.

This collar starts killing fleas and ticks as soon as it is placed around dog's neck. Fleas on the dog will be killed (within 4-8 hours after collar attachment) and new ones which may temporarily appear on the dog will also be killed while collar is worn.

Ticks are tough and are killed slowly. When the collar is first placed on dog, adult ticks will be killed over entire body in a few days and will fall off or may then be easily removed. Ticks appear on dogs in three stages. In the first two stages ticks are smaller than a match head and are difficult to see. The collar will also control these immature stages. Use in addition to regular collar.

This collar has been shown to provide effective flea (control or kill) up to 11 months and tick (control or kill) up to 7 months plus aiding in tick (control or kill) for an additional 2 months. Replace when effectiveness diminishes.

Kills ticks that may carry and transmit Rocky Mountain spotted fever and typhus.

This collar will also aid in the prevention of Sarcoptic Mange for up to 5 months.

**STORAGE & DISPOSAL:** Store in original unopened container; away from children. Do not reuse container or used collar; wrap and put in trash.

Zoecon Industries  
A Division of Zoecon Corporation  
12200 Denton Drive, Dallas, Texas 75234

EPA Reg. No. 2724-282

EPA Est.  
c 1982 Zoecon  
Made in USA  
684-B-2807A:0089A