

[] Indicates optional statements

ZOECON COLLAR RF-159 FOR DOGS

With [Fits] necks to 19" Long Lasting
With [Fits] necks to 25" Long Lasting Protection

(Controls or Kills) Fleas Up To 11 Months

(Controls or Kills) Ticks Up To 7 Months

(Controls or Kills) Fleas & Ticks Up to 7 Months

Aids in Tick Control For An Additional 2 Months

From Head To Tail

Aids In The Prevention of Sarcoptic Mange
For Up To 5 Months

[One-piece buckle design]

[Integrated plastic buckle]

[Unique buckle design]

[New integrated one-piece buckle]

[One-piece buckle]

ACTIVE INGREDIENT:

Chlorpyrifos: 0,0-Diethyl 0-(3,5,6-
trichloro-2-pyridyl) phosphorothioate..... 8.0%
INERT INGREDIENTS..... 92.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See other precautions on back.

NET CONTENTS:

Reg # 2724-202
1/1/12

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION: Do not open protective pouch until ready to use. Do not allow children to handle this collar. Wash hands thoroughly with soap and water after handling collar. Do not use on sick and convalescing dogs. Do not use other pesticides on dogs while collar is worn. Do not use on dogs under 12 weeks of age. This product contains a cholinesterase inhibitor. Some breeds or individuals may be sensitive. Remove collar if symptoms of cholinesterase inhibition (which may include salivation, vomiting, and/or diarrhea) occur. Atropine is antidotal only if symptoms of cholinesterase inhibition are present.

DIRECTIONS FOR USE - It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Buckle collar around dog's neck. [Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Since the plastic buckle is molded into the collar, the probability of neck irritation and buckle failure is reduced.] Collar should be worn loosely to prevent irritation. Attach collar so two or three fingers may be inserted easily between collar and dog's neck. Cut off any excess length and dispose of it. When collar is first worn observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. [To release the collar, squeeze buckle between thumb and finger and pull sharply.]

This collar starts killing fleas and ticks as soon as it is placed around dog's neck. Fleas on the dog will be killed (within 4-8 hours after collar attachment) and new ones which may temporarily appear on the dog will also be killed while collar is worn.

Ticks are tough and are killed slowly. When the collar is first placed on dog, adult ticks will be killed over entire body in a few days and will fall off or may then be easily removed. Ticks appear on dogs in three stages. In the first two stages ticks are smaller than a match head and are difficult to see. The collar will also control these immature stages. Use in addition to regular collar.

This collar has been shown to provide effective flea (control or kill) up to 11 months and tick (control or kill) up to 7 months plus aiding in tick (control or kill) for an additional 2 months. Replace when effectiveness diminishes.

Kills ticks that may carry and transmit Rocky Mountain spotted fever and tularemia.

This collar will also aid in the prevention of Sarcoptic Mange for up to 5 months.

STORAGE & DISPOSAL:

Store in original unopened container, away from children. Do not reuse container or used collar. Wrap and put in trash.

ZOECON INDUSTRIES
A Division of Zoecon Corporation
12200 Denton Drive, Dallas, Texas 75234

EPA Reg. No. 2724-282
285-B:6943A:0144A

ACCEPTED
JAN 07 1985
Under Insecticide Act
Fungicide as a pesticide
as registered at
EPA Reg. No. 2724-282

EPA Est.
Made in USA
c 1982 Zoecon