

2724-282

1/27/98

1/3

[] indicates optional statements

SANDOZ DURSBAN COLLAR FOR DOGS (RF-9411)

- For dogs with [Fits] necks to [number of inches]
- Long Lasting Protection
- [Step 2][Pet Protection Program]
- [Part of the (Brand Name)(#) step guarantee program]
- [Controls or Kills][adult] fleas up to 11 months
- [Controls or Kills] ticks up to 7 months
- [Controls or Kills] fleas & ticks up to 7 months
- Aids in Tick control for an Additional 2 months
- From Head to Tail
- Aids in the Prevention of Sarcoptic Mange for up to 5 Months
- [One-piece buckle design]
- [Integrated plastic buckle]
- [Unique buckle design]
- [Integrated one-piece buckle]
- With [contains] Dursban®

ACTIVE INGREDIENT:

Chlorpyrifos (CAS #2921-88-2)..... 8.2%

OTHER INGREDIENTS:..... 91.8%

Total: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See other precautions on back

NET CONTENTS:

ACCEPTED

JAN 27 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 2724-282

READ ENTIRE LABEL BEFORE EACH USE
USE ONLY ON DOGS

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

1. Buckle collar around dog's neck. [Thread flat end of the collar through buckle hole and tighten as necessary, so that at least two ridges are covered by the buckle. Since the plastic buckle is molded into the collar, the possibility of neck irritation and buckle failure is reduced.] Collar should fit snug enough to prevent placement of the dog's leg or lower jaw through the collar, yet loose enough to be rotated about the neck. [Generally, if used on large dogs, two or three fingers should fit between the collar and dog's neck.]
2. Leave 2-3 extra inches on the collar for adjustment, cut off excess length and dispose of in trash.
3. Check periodically and adjust fit if necessary, especially when puppies are rapidly growing.
4. May be used in addition to regular collar. Replace collar when effectiveness diminishes. [To release the collar, squeeze buckle between thumb and finger and pull sharply.]

STORAGE and DISPOSAL:

Storage: Store in original unopened container away from children. **Disposal:** Do not reuse container or used collar. Securely wrap in several layers of newspaper and discard in trash.

WHAT TO EXPECT:

For Fleas:

1. This [brand name] collar is designed to kill adult fleas.
2. This collar starts killing fleas as soon as it is placed around dog's neck.
3. Protection against fleas will be achieved within a few days, providing continuous protection for up to 11 months.
4. Replace collar every 11 months to help control [prevent] flea infestations on your pet and in your home.

For Ticks:

1. Ticks are tougher to kill than fleas and are killed more slowly.
2. Full protection against adult and immature ticks will be achieved within a few days, providing continuous protection for 7 months plus aiding in tick [control or kill] for an additional 2 months.
3. Adult ticks will be killed over the entire pet. Dead ticks will fall off or may be easily removed.

Kills ticks that may carry and transmit Rocky Mountain spotted fever and tularemia. This collar will also aid in the prevention of Sarcoptic Mange for up to 5 months

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION: This product contains a cholinesterase inhibitor. Some breeds or individuals may be sensitive. Remove collar if symptoms of cholinesterase inhibition (which may include salivation, labored breathing, incoordination, and pin-point pupils) occur.

Humans: Do not open protective pouch until ready to use. Do not allow children to handle this collar. Wash hands thoroughly with soap and water after handling collar.

Animals: Consult a veterinarian before using this product on debilitated, aged, medicated, pregnant, or nursing dogs. Do not use this product on dogs simultaneously or within 30 days before or after treatment with or exposure to cholinesterase inhibiting drugs, pesticides, or chemicals. However, flea and tick collars may be immediately replaced. Do not use on dogs under 12 weeks of age. When collar is first worn, observe neck area every few days for irritation. Remove collar at the first sign of irritation or adverse reaction. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, remove collar and bathe your pet with mild soap and rinse with large amount of water.

FIRST AID: If Swallowed: Call a physician/veterinarian or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting. **If in Eyes:** Flush eyes with plenty of water. Call a physician if irritation persists. **If on Skin:** Wash with plenty of soap and water. Get medical attention if irritaiton persists.

NOTE TO PHYSICIAN/VETERINARIAN: Atropine is antidotal only if symptoms of cholinesterase inhibition are present.

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

For more information or in case of an emergency call [phone number]

Wellmark International
Bensenville, IL 60106

Dursban® is a registered trademark of DowElanco

EPA Reg. No. 2724-282
Made in USA

EPA Est. No.