1

ZOECON BLOCK IGR CATTLE SUPPLEMENT

PREVENTS ADULT HORN FLY EMERGENCE FROM MANURE OF TREATED CATTLE

A Nutritional supplement Containing Methoprene* Insect Growth Regulator for Continuous Feeding to Cattle through the Horn Fly Season.

ACTIVE INGREDIENT:

Methoprene [Isopropyl (E,E)-11-methoxy-3,7,11-trimethyl-2,4-dodecadi	enoate] 0.02%
INERT INGREDIENTS:	99.98%
TOTAL	100.00%


GUARANTEED ANALYS	IS
Calcium (Cā), not less than	
Calcium (Ca), not more than	
Phosphorus (P), not less than	
Salt (NaCl), not less than	
Salt (NaCl), not more than	
lodine (I), not less than	
Vitamin A, not less than	
USP units per pound	

FEED INGREDIENTS

Vitamin A Acetate, D-activated Animal Sterol (Source of Vitamin D₃-Stability improved), Dicalcium Phosphate, Defluorinated Phosphate, Calcium Carbonate, Low Fluorine Ground Rock Phosphate, Soybean Meal, Salt, Iron Sulfate, Iron Oxide, Potassium Iodide, Cobalt Carbonate, Silicon Dioxide, Copper Carbonate, Cane Molasses, Manganous Oxide, Ethylene Diamine Dihydriodide, Zinc Oxide, Magnesium Oxide, Paraffin Wax, Hemicellulose Extract

KEEP OUT OF REACH OF CHILDREN CAUTION

NET WEIGHT:



^{*}As the Insect Growth Regulator ALTOSID®

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Using it in a manner other than that described by the label could result in harm to aquatic invertebrates. Do not contaminate water when disposing of rinsate or equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

FEEDING AND MANAGEMENT INSTRUCTIONS

When used as directed ZOECON BLOCK IGR CATTLE SUPPLEMENT will provide sufficient methoprene insect growth regulator to prevent the emergence of adult horn flies from manure of treated cattle. Existing adult horn flies will not be affected. Start feeding before horn flies appear and continue use until cold weather marks the end of the horn fly season. Use the following management techniques to maintain continuous intake of Zoecon Block IGR Cattle Supplement at a level of 1/4 - 1/2 lb per 100 lb of animal body weight per month.

- Assure cattle are not starved for nutrients in Zoecon Block IGR Cattle Supplement by providing them prior to feeding Zoecon Block IGR Cattle Supplement.
- 2. Offer cattle 50 lb of Zoecon Block IGR Cattle Supplement for every fifteen head of cattle.
- 3. Remove all other self limiting nutritional supplements.
- 4. Locate Zoecon Block IGR Cattle Supplement where cattle congregate (watering, loafing, shade areas).
- 5. Replenish Zoecon Block IGR Cattle Supplement on a regular basis as needed.
- 6. If intake of Zoecon Block IGR Cattle Supplement is <u>below</u> 1/4 lb per 100 lb animal body weight per month increase the number of feeding locations and/or change location(s) to areas more frequented by cattle.
- 7. If intake of Zoecon Block IGR Cattle Supplement is <u>above</u> 1/2 lb per 100 lb animal body weight per month reduce the number of feeding locations and/or change location(s) to areas less frequented by cattle.
- 8. Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if immediate elimination of the adult fly population is desired.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal.

Storage: Store Zoecon Block IGR Cattle Supplement in a cool place.

Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities by burning. If burned, stay out of smoke.

EPA Reg. No.: 2724-373

Made in USA

EPA Est. No.:

ALTOSID® is a registered trademark of Wellmark International.

Wellmark International Bensenville IL 60106