



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

September 29, 2016

Nancy Huebl
Senior Registration Specialist
Wellmark International
1501 East Woodfield Road, Suite 200W
Schaumburg, Illinois 60173

Subject: Pesticide Registration Improvement Act (PRIA) Labeling Amendment – Label amendment plus supplemental labeling to add aerial applications and update tank mixing directions
Product Name: Altosid Liquid Larvicide
EPA Registration Number: 2724-392
Application Date: May 19, 2016
OPP Decision Number: 518020

Dear Ms. Huebl:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

The supplemental labeling contains some new and/or revised uses and/or directions that may be additional to the uses and/or directions found on the labeling on or attached to the container, but the supplemental labeling does not by itself constitute the complete set of use directions. The complete set of use directions is set forth on the container labeling as combined with the supplemental labeling.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 § CFR 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.


If you have any questions, please contact Menyon Adams of my branch by phone at (703) 347-8496 or via email at adams.menyon@epa.gov.

Sincerely,

A handwritten signature in blue ink that reads "Andrew C. Bryceland". The signature is fluid and cursive, with the first name being the most prominent.

Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure



ALTOSID® LIQUID LARVICIDE

An Insect Growth Regulator for Mosquito Control
PREVENTS EMERGENCE OF ADULT FLOODWATER MOSQUITOES

For control of mosquito larvae using ULV application

Microencapsulated formula

Prevents adult mosquito emergence (including those which may transmit West Nile virus, Zika, chikungunya and dengue)

ACTIVE INGREDIENT:

(S)-Methoprene (CAS #65733-16-6) 5%

OTHER INGREDIENTS: 95%

TOTAL: 100%

Formulation contains 0.43 lb/gal (51.3 g/liter) active ingredient

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

BECAUSE OF THE UNIQUE MODE OF ACTION OF A.L.L.™, SUCCESSFUL USE
REQUIRES FAMILIARITY WITH SPECIAL TECHNIQUES FOR APPLICATION TIMING AND
TREATMENT EVALUATION. SEE **GUIDE TO PRODUCT APPLICATION** OR CONSULT
LOCAL MOSQUITO ABATEMENT AGENCY.

EPA Reg. No. 2724-392
EPA Est No.

NET CONTENTS:

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS - CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

FIRST AID	
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.	

[Note: First Aid statement paragraph format may be used if space is limited.]

ENVIRONMENTAL HAZARDS: Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

CHEMIGATION: Refer to supplemental labeling entitled "**Guide to Product Application**" for use directions for chemigation. Do not apply this product through any irrigation system unless the supplemental labeling on chemigation is followed.

MIXING AND HANDLING INSTRUCTIONS:

1. **SHAKE WELL BEFORE USING.** A.L.L.TM may separate on standing and must be thoroughly agitated prior to dilution.
2. Do not mix with oil; use clean equipment.
3. Partially fill spray tank with water; then add the labeled amount of A.L.L.TM, agitate and complete filling. Mild agitation during application is desirable.
4. Use spray solution within 48 hours; always agitate before spraying.

APPLICATION INSTRUCTIONS - Introduction: A.L.L.TM must be applied to 2nd, 3rd, or 4th larval instars of target mosquitoes to prevent adult emergence. Treated larvae continue normal development to the pupal stage where they die. This insect growth regulator **has no effect when applied to pupae or adult mosquitoes.** A.L.L.TM has sufficient field life to be effective at appropriate rates when applied to larval stages under varying field conditions. For further information, see **Guide to Product Application.**

METHODS OF APPLICATION - Aerial: A.L.L.TM may be applied by aircraft to terrestrial locations where mosquito breeding occurs. Use the prescribed amount of A.L.L.TM listed below in sufficient water to give complete coverage. One-half to 5 gallons of spray solution per acre is usually satisfactory. Do not apply when weather conditions favor drift from areas treated.

Ground: Determine the average spray volume used per acre by individual operators and/or specific equipment. Mix A.L.L.TM in the appropriate volume of water to give the rate per acre as indicated below.

For Control of Mosquito Larvae Using ULV Application:

For ground and aerial application to terrestrial sites, apply at the rate of 3-4 fluid ounces of product per acre to water-holding containers and other small bodies of water that breed mosquitoes. Use equipment capable of applying a fine mist or ULV. Follow equipment manufacturer's recommendations when making applications. Direct spray applications to sites where mosquitoes breed. These sites include tires and tire piles, potted plants, tree holes, garbage bins, cans, birdbaths, rain barrels, and other water-holding containers and small bodies of water.

For aerial application to terrestrial sites, apply by fixed wing or rotary aircraft. Apply at the rate of 3 – 4 fluid ounces of product to acre diluted with water at a minimum of a 1:1 mix ratio with water. Apply using ULV equipped and capable aircraft. Unlike ULV sprays targeting flying mosquitoes, it is important that spray droplets deposit in targeted areas. Target terrestrial areas where mosquitoes breed. These sites include tires, open containers, garbage bins, birdbaths, and gutters, which hold small amounts of water. Spray equipment must be adjusted so that the volume median diameter (VMD) produced ranges from 60 microns ($D_{V0.5} < 60\mu$) to 100 microns ($D_{V0.5} < 100\mu$), and that 90% of the spray is contained in droplets smaller than 200 microns ($D_{V0.9} < 200\mu$). Directions from the equipment manufacturer or vendor, pesticide registrant, or test facility using a wind tunnel and laser-based measurement instrument must be used to adjust equipment to produce acceptable droplet size spectra. Application equipment must be calibrated annually to confirm that nozzle flow rate(s) are accurate. Do not apply at altitudes below 100 feet unless using unmanned aircraft designed for low application heights. Apply when wind speed on the ground is ≥ 1 mph and ≤ 10 mph. Apply when wind factors promoting drift are low. For best results, use Global Positioning System (GPS) equipped aircraft.

APPLICATION RATE: Apply 3 to 4 fl oz of A.L.L.™ per acre (219 to 293 ml/hectare) in water as directed.

For Control of Mosquito Larvae Using ULV Application:

Apply at 3-4 fluid ounces per acre. Reapply as breeding sites become reinfested or when monitoring indicates an increase in adult populations.

APPLICATION SITES - Pastures: A.L.L.™ may be applied after each flooding without removal of grazing livestock.

Rice: A.L.L.™ must be applied to 2nd, 3rd, and/or 4th instar larvae of mosquitoes found in rice, usually within 4 days after flooding. A.L.L.™ treatment may be repeated with each flooding.

Intermittently Flooded Noncrop Areas: A.L.L.™ may be applied as directed above when flooding may result in floodwater mosquito hatch.

Typical sites include freshwater swamps and marshes, salt marshes, woodland pools and meadows, dredging spoil sites, drainage areas, waste treatment and settling ponds, ditches and other natural and manmade depressions.

Crop Areas: A.L.L.™ may be applied to irrigated croplands after flooding to control mosquito emergence. Examples of such sites are: vineyards, rice fields (including wild rice), date palm orchards, fruit and nut orchards and berry fields and bogs. Irrigated pastures may be treated after each flooding **without** the removal of grazing livestock.

In dense vegetation or canopy areas, apply an A.L.L.™ sand or BIODAC® mixture using standard granular dispersal equipment. For detailed preparation instructions, refer to **Guide to Product Application**.

TANK MIXING INSTRUCTIONS

The user, at his discretion, can [tank] mix A.L.L.™ with registered adulticides or other larvicides such as *Bacillus thuringiensis* variety *israeliensis* (*B.t.i.*) or *Bacillus sphaericus* (*B.s.*) unless the tank-mix product label prohibits such mixing. Use the resulting tank mixture in accordance with the more restrictive label limitations and precautions.

STORAGE AND DISPOSAL: Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in cool place. Store product away from other pesticides, food, and feed. In case of leakage or spill, soak up with sand or another absorbent material. **Pesticide**

Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling:** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows. . Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

To the extent consistent with applicable law, Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

Wellmark International
1501 East Woodfield Road 200W
Schaumburg IL 60173

ALTOSID® Liquid Larvicide and A.L.L.™ are trademarks of Wellmark International.

For information, call 1-800-248-7763 or visit our website: www.altosid.com .

Made in USA

**ALTOSID® LIQUID LARVICIDE
MOSQUITO GROWTH REGULATOR**

PREVENTS EMERGENCE OF ADULT FLOODWATER MOSQUITOES

GUIDE TO PRODUCT APPLICATION

FOR THE FIRST TIME USER

ALTOSID® Liquid Larvicide (A.L.L.™) is the result of extensive research into the intricacies of natural biochemical and physiological development of insects. New chemical technology and biological findings were combined to develop a unique mosquito larvicide.

A.L.L.™, an insect growth regulator (IGR), acts by inducing morphological changes which interfere with normal development. These effects, not immediately apparent, result in the failure of adult mosquitoes to emerge from pupae. A.L.L.™ is **not** a conventional pesticide. It does not produce the nondiscriminatory rapid, directly toxic effects that are associated with traditional larvicides. A.L.L.™ differs from other larvicides you may have used only in the manner and time course of its action **after** application.

A.L.L.™ is applied to second, third, or fourth instar larvae using standard larviciding equipment in a manner similar to other larvicides. After application to second, third, or fourth instar larvae at the appropriate rates, absolutely no effects on larvae will be observed. They will continue developing normally and will pupate. Pupae will appear unaffected, but will eventually die. **Adults will not emerge.** Infrequently, a few adults may be seen at the water surface but they will have abnormalities preventing flight and will not survive. Because the effect of A.L.L.™ is neither larval death nor widespread mortality immediately following pupation, the **number of adults which emerge is the only criterion for accurately assessing control.** Checks by dip counts during larval and pupal stages will give no measure of effectiveness.

Refer to the following diagram and checklist, in addition to label instructions for guidance in timing of application and performance evaluation. They will assist you in obtaining the best possible results with this unique product.

Wellmark International
1501 East Woodfield Road 200W
Schaumburg, IL 60173

For information, call 1-800-248-7763

The information presented herein, while not guaranteed, is to the best of our knowledge true and accurate. No warranty or guarantee, expressed or implied, is made regarding the performance or stability of any product, since the manner of use and conditions of storage and handling are beyond our control.

CHECKLIST

Things to remember when using A.L.L.™


DO the following:

1. **DO** treat second, third, and/or fourth instar larvae, not pupae or adults. (First instar larvae are so small they are not readily detectable.)
2. **DO** wait until treated larvae have pupated. Then **collect pupae** and transfer to laboratory to observe for emergence of adults.
3. **DO** observe pupae for several days, since death of IGR treated mosquitoes occurs when pupae would normally emerge as adults. (Careful observation is necessary since dead pupae decompose rapidly and thus are not easily seen.)
4. **DO** monitor emerging adults at the treatment site. This absolutely requires that emergence traps be placed in treatment areas to capture adult mosquitoes as they emerge.

DO NOT do the following:

1. **DO NOT** take dip counts of larvae after treatment for the purpose of performance evaluation. **Normal looking larvae** will be present.
2. **DO NOT** take dip counts of pupae after treatment for the purpose of performance evaluation. **Normal looking pupae** will be seen but these will not develop into normal adults.
3. **DO NOT** think A.L.L.™ has failed if some adult mosquitoes are flying in treated areas; they probably have flown in from nearby untreated areas. Numbers 2 and 4 of the "**DO**" checklist are the only methods of accurately assessing effectiveness.
4. **DO NOT** spray again, either with A.L.L.™ or a conventional insecticide, because larvae or pupae are present after application. This is normal. The effectiveness of A.L.L. can only be measured by lack of adult emergence.

LIFE CYCLE OF MOSQUITO WHEN TO APPLY A.L.L.™



Preparation of ALTOSAND® Granular Formulation

An "On-Site" Method of Preparing a Granular Formulation of ALTOSID® Liquid Larvicide

INTRODUCTION

A method of application of ALTOSID® Liquid Larvicide (A.L.L.™), using sand as a carrier, has been developed for use in floodwater mosquito breeding areas with dense vegetation or canopy. The characteristics of ALTOSAND® provide excellent foliage penetration, ensuring that the active ingredient reaches the water where it is released from the sand.

ALTOSAND® will prevent the emergence of species of the floodwater mosquito complex when applied to second, third, or fourth larval instars at a rate of ten (10) to thirteen (13) pounds per acre.

PREPARATION INSTRUCTIONS

The following materials are required to prepare a 100 lb batch of ALTOSAND®:

96 lb washed, dry sand (20 to 45 mesh)
2 lb A.L.L. (15 fl oz./lb)
2 lb HiSil 233 (silicon dioxide)
Small Funnel
Cement Mixer

1. Measure the time required for a level funnel full of sand to empty.
2. Into a rotating-type mixer, place 96 lb of dry (20 to 45 mesh) sand. While the mixer is rotating, slowly pour 2 lb (30 fl oz) of A.L.L.™ onto the sand. (If better wetting is required, A.L.L.™ may be diluted in up to an equal volume of water.)
3. Mix until the sand is uniformly coated with A.L.L.™ (usually 5 to 10 minutes).
4. Stop the mixer and add 2 lb of HiSil 233. Cover the mixer to reduce dust. Start the mixer and run for approximately 5 minutes. (The quantity of HiSil 233 necessary to achieve a dry, free-flowing mixture will vary depending on the particle size distribution and moisture of the sand.)
5. Compare the flow rate of the ALTOSAND® mixture with that of untreated sand in Step No. 1. Add more HiSil if it flows significantly slower and reduce the amount of HiSil in subsequent batches if the mixture flows at the same or a faster rate and is excessively dusty.

APPLICATION RATE AND METHODS

Apply at a rate of 10 to 13 lb of the final mixture per acre, using standard granular dispersal equipment.

Preparation of ALTODAC™ Granular Formulation

An "On-Site" Method of Preparing a Granular Formulation of ALTOSID® Liquid Larvicide

INTRODUCTION

A method of ALTOSID® Liquid Larvicide (A.L.L.™) application, using BIODAC® as carrier, has been developed for use in mosquito breeding areas in floodwater and intermittently flooded noncrop areas including freshwater and saltwater marshes. The characteristics of A.L.L.™ using BIODAC® carrier provide excellent coverage, ensuring that the active ingredient reaches the water and is released from the BIODAC®.

A.L.L.™ will prevent the emergence of adult mosquitos when applied to second, third, or fourth larval instars at a rate of ten (10) to thirteen (13) pounds per acre.

PREPARATION INSTRUCTIONS

The following materials are required to prepare a 100 lb batch of ALTODAC™ using BIODAC® carrier:

96 lb BIODAC® 12/20
30 oz ALTOSID® Liquid Larvicide Mosquito Growth Regulator
32 oz water

1. Weigh the required amounts of A.L.L.™ into a tared container suitable for mixing.
2. Weigh the water into the vessel containing the A.L.L.™. Stir the contents in the vessel until a uniform mixture is achieved.
3. Add the appropriate amount of BIODAC® 12/20 to a blending device, e.g., a cement mixer with lifters, a munson blender, or any other device that will allow the granules to tumble through a spray.
4. Add the water/A.L.L.™ mixture to a spray unit or any pressurized device capable of delivering a cone-shaped, fine particle size spray to the contents in the blender.
5. Spray the mixture of A.L.L.™ and water onto the BIODAC® while the blender is tumbling the granules. Once the mixture has been applied to the BIODAC®, continue to blend until the granules appear to be dry (usually five to ten minutes).
6. Remove the granules and screen over a 12 mesh screen to remove agglomerates.

APPLICATION RATE AND METHODS

Apply at a rate of 10 to 13 lb of the final mixture per acre using granular dispersal equipment.

Chemigation: Apply this product only through flood (basin), furrow, or border irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Flood (Basin), Furrow and Border Chemigation: Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

A pesticide supply tank is recommended for the application of A.L.L.™ by chemigation.

ALTOSAND®, ALTODAC™, ALTOSID® and A.L.L.™ are trademarks of Wellmark International. BIODAC® is a registered trademark of Edward Lowe Industries, Inc.

EPA Reg. No. 2724-392

ALTOSID® LIQUID LARVICIDE

Supplemental Labeling

FOR CONTROL OF MOSQUITO LARVAE USING AERIAL ULV APPLICATION AND UPDATED TANK MIXING DIRECTIONS FOR USE

EPA Reg. No. 2724-392

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read the label affixed to the container for **Altosid® Liquid Larvicide** before using.

Important

Use of **Altosid® Liquid Larvicide** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Altosid® Liquid Larvicide**.

For Control of Mosquito Larvae Using ULV Application:

For aerial application to terrestrial sites, apply at the rate of 3 - 4 fluid ounces of product per acre to water-holding containers and other small bodies of water that breed mosquitoes. Use equipment capable of applying a fine mist or ULV. Follow equipment manufacturer's recommendations when making applications. Direct spray applications to sites where mosquitoes breed. These sites include tires and tire piles, potted plants, tree holes, garbage bins, cans, birdbaths, rain barrels, and other water-holding containers and small bodies of water. Reapply as breeding sites become reinfested or when monitoring indicates an increase in adult populations.

TANK MIXING INSTRUCTIONS

The user, at his discretion, can [tank] mix A.L.L.™ with registered adulticides or other larvicides such as *Bacillus thuringiensis* variety *israeliensis* (*B.t.i.*) or *Bacillus sphaericus* (*B.s.*) unless the tank-mix product label prohibits such mixing. Use the resulting tank mixture in accordance with the more restrictive label limitations and precautions.

For information or in case of emergency, contact 1-800-248-7763.

This labeling must be in the possession of the user at the time of application.

Altosid is a registered trademark of Wellmark International.

Registrant:

Wellmark International

1501 East Woodfield Road 200W

Schaumburg, Illinois 60173

www.altosid.com

A C C E P T E D

09/29/2016

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 2724-392

This supplemental label expires on September 29, 2019 and must not be used or distributed after this date.