

2724-404

[] Indicate optional statement

ZOECON RF-322 OVICIDAL PUMP SPRAY

EFFECTIVE

Kills & Repels: Ticks, fleas, lice, flies, mosquitoes and gnats.

Kills fleas before they mature into adults.

Kills flea eggs.

Prevents [Stops] flea eggs from becoming full grown for 30 days [4 weeks] [with PRECOR^R[VIGRENTM] Insect Growth Regulator]

One treatment works for 30 days [against flea eggs] [to keep flea eggs from hatching]

Effectively breaks the flea life cycle

Keeps working to keep flea eggs from hatching for 30 days [4 weeks] with PRECOR^R [VIGRENTM] Insect Growth Regulator

No need to pump, spray in any position, even upside down


Controls [Kills] [and Repels] Lyme disease-carrying ticks [including Ixodes dammini]

Reduces bot infestation by killing botfly eggs

Prevents botfly eggs from hatching

ACTIVE INGREDIENTS:

(S)-Methoprene [Isopropyl (2E, 4E, 7S)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate].....	0.25%
Pyrethrins.....	0.18%
Piperonyl butoxide, technical*.....	0.36%
N-octyl bicycloheptene dicarboximide.....	0.60%
INERT INGREDIENTS.....	98.61%
TOTAL.....	100.00%



*Equivalent to 0.29% (butylcarbityl) (6-propylpiperonyl) ether and 0.07% related compounds

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back Panel For Additional Precautionary Statements

EPA Reg. No. 2724-404
EPA Est No.

NET CONTENTS:

Zoecon RF-322 combines the insecticidal and repellency action of pyrethrins with Precor [VIGREN] IGR, an insect growth regulator. RF-322 controls ticks, lice, mosquitoes, gnats, fleas, house flies, stable flies, horn flies, face flies, flea larvae and horse bots. Botfly eggs and flea eggs sprayed with RF-322 will not hatch into adults.

To control [kill] adult fleas and prevent hatching of flea eggs, apply RF-322. The use of an effective product for controlling adult and larvae flea populations in the pets environment, like [name of Zoecon premise product(s)], will reduce the possibility of re-infestation of the animal.

DIRECTIONS FOR USE - It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

FOR CATS: A light, even coverage of the hair coat will provide effective flea control. Apply at a rate of 1 or 2 squeezes of the trigger sprayer per pound of body weight. Cover animal's eyes with hand and with a firm, fast stroke, spray head, ears and chest. With cloth, rub into face around mouth, nose and eyes. Then spray the neck, middle and hind quarters, finishing with legs last. For best penetration, spray against natural lay of the hair. On a heavily coated animal, ruffle the coat while spraying against the natural lay of the hair. Make sure spray thoroughly wets ticks. Do not allow animal to become chilled. Repeat as necessary.

FOR DOGS: Cover animal's eyes with hand and with a firm, fast stroke, spray head, ears and chest until damp (1 or 2 squeezes of the trigger sprayer per pound of body weight). With cloth, rub into face around mouth, nose and eyes. Then spray the neck, middle and hind quarters, finishing with legs last. For best penetration, spray against natural lay of the hair. On a heavily coated animal, ruffle the coat and spray against the lay of the hair. Make sure spray thoroughly wets ticks. Repeat as necessary.

FOR PUPPIES AND KITTENS: Puppies and kittens may be treated by wrapping the animal in a towel which has been lightly sprayed. Leave eyes, nose and mouth exposed. Keep the animal wrapped for 5 to 10 minutes. Adult fleas may then be removed from the coat by picking or combing. Larger-size puppies and kittens may also be treated the same as dogs and cats at a rate of 1 or 2 squeezes of the trigger sprayer per pound of body weight. Repeat only as necessary. Do not allow animal to become chilled. Do not treat newborn puppies and kittens.

FOR HORSES AND PONIES: Remove excess dirt and dust by brushing coat. Apply spray directly to horse's coat giving particular attention to legs, shoulders, flanks, neck and topline. To treat face, dampen a cloth with spray and rub on face and head, being careful to avoid eyes. Repeat as necessary. Do not treat horses destined for food.

FOR PREVENTION OF HATCH OF HORSE BOTFLY EGGS: Spray any infested areas directly at the first sign of visible eggs, paying particular attention to legs and shoulders. Spray to the point of dampness but not to the point of run-off of the spray. For a high degree of botfly egg control, spray animals every 3-5 days during the botfly season.

STORAGE & DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Do not store near heat or open flame. Store away from children. Do not reuse empty container. Wrap and put in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION: Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. May cause dermal sensitization.

Some cats may exhibit salivation. If concerned, call 1-800-[appropriate phone number] or consult your veterinarian.

STATEMENT OF PRACTICAL TREATMENT: If swallowed, call physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of the throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. If on skin, remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists. If in eyes, flush with plenty of water. Get medical attention if irritation persists.

PHYSICAL HAZARDS: Flammable. Keep away from heat and open flame.

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

SANDOZ AGRO INC.
1300 East Touhy
Des Plaines, IL 60018

VET-KEM [ZODIAC] , OVITROL [FLEATROL] [PETCOR] and PRECOR [VIGREN] are trademarks of Sandoz Ltd.

Made in USA

© 1995 Sandoz Agro, Inc.

For Information, call 1-800 [phone number]