

() Indicates optional statement

PM 90

2724-462

Page 1 of 1

ZOECON 9307 PET SPRAY

Kills Fleas

Egg Stop Formula

Kills Fleas [Dead] Plus Their Hatching Eggs [for up to 28 Days][on Your Dog]

Kills Flea Eggs [Before They Hatch]

[Effectively] Breaks the Flea Life Cycle

Keeps [Working to Keep] Flea Eggs from Hatching for 28 days [4 Weeks][with VIGRENtm Insect Growth Regulator]

Kills [and Repels] Fleas [and Ticks][on Dogs]

Kills Adult and Pre-Adult Fleas (Eggs)

Prevents Hatching Fleas From Becoming Full Grown [Fleas]

No Need to Pump, Sprays in Any Position, Even Upside Down

One treatment works for 28 Days to Keep Flea Eggs from Hatching

ACTIVE INGREDIENTS:

(S)-Methoprene [isopropyl (2E, 4E, 7S)-11-methoxy-3,7,11-trimethyl-2,4-dodecadienoate].....	0.10%
Pyrethrins.....	0.18%
Piperonyl butoxide, technical*.....	0.36%
N-octyl bicycloheptene dicarboximide.....	0.60%
INERT INGREDIENTS.....	98.76%

Equivalent to 0.29% (butylcarbityl) (6-propylpiperonyl) ether and 0.07% related compounds

KEEP OUT OF REACH OF CHILDREN

CAUTION ACCEPTED

See Back Panel For Additional
Precautionary Statements

FEB 27 1995

Under the Federal Insecticide, Fungicide, and
Rodenticide Act, as amended, for the
pesticide registered under
EPA Reg. No. 2724-462

NET CONTENTS:

DIRECTIONS FOR USE - It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USING.

FOR DOGS: Recommended dosage rate is [1 to 2 squeezes of the trigger sprayer][1 to 2 seconds of spray] per pound of body weight. Cover animal's eyes with hand and [with a firm, fast stroke,] spray head, ears and chest until damp. With cloth, rub into face around mouth, nose and eyes. Then spray the neck, middle and hind quarters, finishing with legs last. For best penetration, spray against natural lay of the hair. On a heavily coated animal, ruffle the coat and spray against the lay of the hair. Make sure spray thoroughly wets ticks. Repeat as necessary.

FOR PUPPIES: Puppies may be treated by wrapping the animal in a towel which has been lightly sprayed. Leave eyes, nose and mouth exposed. Keep the animal wrapped for 5 to 10 minutes. Adult fleas may then be removed from the coat by picking or combing. Larger-size puppies may also be treated the same as dogs with [1 to 2 squeezes of the trigger sprayer] [1 to 2 seconds of spray] per pound of body weight. Repeat only as necessary. Do not allow animal to become chilled. Do not treat newborn puppies.

STORAGE & DISPOSAL: Do not store near heat or open flame. Store away from children. Do not reuse empty container. Wrap and put in trash.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS-CAUTION: Harmful if bed through skin. Avoid contact with skin, eyes, or clothing. May skin sensitization in rare cases with repeated use.

STATEMENT OF PRACTICAL TREATMENT: If on skin, remove contaminated clothing and wash affected areas with soap and water. Get medical attention if irritation persists. If in eyes, flush with plenty of water. Get medical attention if irritation persists.

PHYSICAL HAZARDS: Flammable. Keep away from heat and open flame.

Zoecon Corporation
A Sandoz Company
12200 Denton Drive, Dallas, Texas 75234

EPA Reg. No. 2724-
Made in USA
01293-A

EPA Est.
© 1987 Zoecon

HTG: hman: 6