

2724-478

2/21/2009

10f19



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

FEB 21 2009

Ms. Nancy Huebl
Sr. Registratrion Specialist
Wellmark International
1501 East Woodfield Road
Suite 200W
Schaumburg, ILL 60173

RE: Notification to add Direction of Use for a similar application method on Yardex sublabel
for perimeter treatments

EPA Registration Number: 2724-478

Date of Submission: November 18, 2008

Dear Ms. Huebl:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated November 18, 2008, for the product Wellmark International/Mavrik Aquaflo^w Insecticide. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.


If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to be "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs

20f19

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
--	---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number Wellmark International/2724-478	2. EPA Product Manager Richard Gebken	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Wellmark International/Mavrik Aquaflow Insecticide	PM# 13	
5. Name and Address of Applicant / Registrant (Include ZIP Code) Wellmark International 1501 East Woodfield Road, Suite 200 West Schaumburg, Illinois 60173 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(C)(3) (b)(i), my product is similar or identical in composition and labeling to: <div style="text-align: center;"> NOTIFICATION FEB 21 2009 </div> EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.


Explanation: Use additional page(s) if necessary. (For section I and section II.)

Label notification to add DFU of similar application method on Yardex sublabel for perimeter treatments. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____	
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Unit Packaging wgt. No. per container		
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 1 pt., 1 qt., 1 gal.		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input checked="" type="checkbox"/> On Label Accompanying Product	
6. Manner in Which Label is Affixed to Product			<input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Nancy Huebl	Title Senior Registration Specialist	Telephone No. (Include Area Code) (847) 330-5376
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Senior Registration Specialist	
4. Typed Name Nancy Huebl	5. Date 18-Nov-08	

3 of 19

Wellmark International
1501 East Woodfield Road, Suite 200 West
Schaumburg, Illinois 60173
847 330 5300
www.wellmarkinternational.com



November 19, 2008

Document Processing Desk (NOTIF)
Office of Pesticide Programs 7504P
U.S. Environmental Protection Agency
Room S-4900 One Potomac Yard
2777 South Crystal Drive
Arlington, Virginia 22202-4501

ATTENTION: Richard Gebken, PM 13

SUBJECT: LABEL NOTIFICATION FOR
MAVRIK AQUAFLOW® INSECTICIDE/MITICIDE
EPA REG. NO. 2724-478
ADD APPLICATION METHOD ON YARDEX® SUBLABEL

Dear Mr. Gebken:

Enclosed is completed EPA Form 8570-1 and 2 copies of revised labeling for the above product. A highlighted version is enclosed for your convenience in viewing the changes (red is new text). Per PRN 98-10, §II.M.3, the change involves the addition of directions for use for a similar application method for perimeter treatments on the YARDEX® sublabel. This revision involves no change in dosage, concentration, timing or frequency of application. I have enclosed a self-addressed postcard for notification confirmation.

If you have any questions please, call me directly at 847-330-5376. Thank you for your assistance in this matter.

Yours sincerely,
WELLMARK INTERNATIONAL

Nancy Huebl
Senior Registration Specialist

enclosure

cc: J. McFadden/C. Elmi/H. Friedli

cc: J. McFadden/C. Elmi/H. Friedli

40f19

MAVRIK AQUAFLOW® INSECTICIDE/MITICIDE

For commercial use only on indoor plants in greenhouses, interior plantscapes, and flower and foliage cuttings and on outdoor plants in ornamental plantings, plantscapes (including trees, shrubs), and containerized nursery stock. Also for Perimeter Treatments around buildings and outside surfaces

ACTIVE INGREDIENT:

Tau-fluvalinate (CAS #102851-06-9)..... 22.3%

OTHER INGREDIENTS: 77.7%

TOTAL 100.0%

MAVRIK AQUAFLOW is a flowable formulation containing 2 pounds tau-fluvalinate per gallon (240 g/L).

EPA Registration No. 2724-478

EPA Est. No. 2724-TX-1


KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional Precautionary Statements and First Aid

SHAKE WELL BEFORE USING

Net Contents



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. **Avoid Breathing Spray Mist.** Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For 24 Hour Medical or Chemical Emergency Assistance, call Wellmark International [Phone Number]	

Sensitive individuals may temporarily experience an itching, burning, or tingling sensation, with or without a rash, following exposure. These symptoms will usually subside without requiring medical treatment. Avoid hand or sleeve-to-face contact. Prior to exposure or after washing, an application of corn oil to exposed skin may reduce these symptoms. Certain persons may be sensitive to MAVRIK AQUAFLOW®'s fine spray particles.


If workers enter the treated area and/or contact treated surfaces:

- Pesticide residues that get on their skin may cause itching or irritation that can be severe, and
- They should avoid skin contact with treated foliage and/or surfaces, and
- They should wash the affected skin immediately if irritation begins to occur.

Personal Protective Equipment (PPE):

WPS USES:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.



All mixers, loaders, and applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) -- in general, agricultural-plant uses are covered -- must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, butyl rubber (≥ 14 mils), nitrile rubber (≥ 14 mils), neoprene rubber (≥ 14 mils), polyvinyl chloride (PVC) (≥ 14 mils), or viton (≥ 14 mils)
- Shoes plus socks
- For exposures outdoors, a NIOSH approved respirator with any R, P, or HE filter.
- For exposures in enclosed areas, a NIOSH approved respirator with an organic vapor (OV) cartridge or canister with any R, P, or HE prefilter.

NON-WPS USES:

Applicators and other handlers who handle this pesticide for any use **NOT** covered by the Worker Protection Standard (40 CFR Part 170) -- in general, agricultural-plant uses are covered by the WPS -- must wear:

- Protective clothing
- Rubber gloves while handling; workers in frequent contact with treated foliage should wear light barrier gloves to reduce excess exposure.
- Indoor exposures: Use a properly fitted NIOSH approved respirator with any R, P, or HE filter for use with pesticides to avoid potential irritation to nose and throat from fine spray particles. NOTE: Persons other than applicator should not be present at time of application.
- Outdoor exposures: For sensitive persons, use a NIOSH approved respirator with any R, P, or HE filter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Engineering Controls Statements:

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment breakdown.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. This product is toxic to honey bees if bees are exposed to direct application. However, dried residues of this product are non-toxic to honey bees. Treat during non-foraging periods to minimize adverse effects. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Use with care when applying in areas adjacent to any body of water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

PHYSICAL/CHEMICAL HAZARDS

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store away from direct sunlight or heat. Pesticides must be stored in a secured area away from other products, food, or feed. In case of spill or leak, soak up with soil or other absorbent material. Do not use caustic agents.

Pesticide Disposal: Wastes resulting from the use of MAVRIK AQUAFLOW may be disposed of on site or at any approved waste disposal facility.

Container Disposal: Triple rinse container (or equivalent). Puncture and dispose of in a sanitary landfill, by incineration or if allowed by state and local authorities by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides.

It contains requirements for training, decontamination, notification, and emergency assistance.

It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, butyl rubber (≥14 mils), nitrile rubber (≥14 mils), neoprene rubber (≥14 mils), polyvinyl chloride (PVC)(≥14 mils), or viton (≥14 mils)
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170).

The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected people, children, and pets out of treated area until spray has dried. Some sensitive pets eating or coming in contact with freshly treated plants may demonstrate vomiting and respiratory distress.

Do not apply this product through any type of irrigation system. Not for broadcast use in nurseries. Not for use on sod farms, on grass grown for seed, or on golf course turf. Do not apply this product by aerial application.

GENERAL DIRECTIONS


MAVRIK AQUAFLOW controls insects and controls or suppresses mite pests when used at labeled rates. MAVRIK AQUAFLOW works primarily through contact action, good spray coverage is necessary. Treat when pests are immature or at a susceptible stage and populations are manageable, before plant damage occurs. Repeat applications as needed, according to the specific use directions given in this label. Residues of MAVRIK AQUAFLOW can continue to control certain insect pests for a week or longer. Growth dilution, climatic variables, and pest habits determine re-treatment needs.

Dilute MAVRIK AQUAFLOW in sufficient water to give thorough spray coverage [based on previous experience]. For best results under varying conditions, a range of use rates is recommended for most uses. Select the higher rates when pest pressure or damage potential is severe. The lower rates often give acceptable economic results. [Buffer spray water to pH 5-7, if necessary.]

For severe mite infestations, use the highest rate of MAVRIK AQUAFLOW and use an acaricide in an alternate treatment program. The use of a spreader-sticker, insecticidal soaps, or piperonyl butoxide may improve control of mites and some insects, such as, whiteflies, aphids, thrips, and mealybugs. MAVRIK AQUAFLOW may be used as a tank-mix. The more stringent product label will apply when tank-mixing two products. When MAVRIK AQUAFLOW is tank-mixed with wettable powders, add powders to the tank first using good agitation. Check for physical and biological compatibility prior to use.

Dried residues of MAVRIK AQUAFLOW are non-toxic and non-repellant to honey bees. Treat during non-foraging periods to minimize adverse effects.

Consult your local extension office, pest management consultant, or a Wellmark International specialist for guidance in selecting a rate and timing schedule or tank mix combination that has demonstrated good performance under specific conditions. Follow all use directions and precautions for products used in tank mix.



9/29/19

CROP/SITE	PEST	AMOUNT MAVRIK AQUAFLOW IN WATER TO MAKE		SPECIFIC DIRECTIONS
		100 GALLONS	5 GALLONS	
<p>For Outdoor Applications: Use lower rates and greater re-treatment intervals for light to moderate insect pressures. Use higher rates and shorter re-treatment intervals for green peach aphids, unexposed insects and heavy pressures of other insects. Mite suppression can be obtained using the lower rate; for more residual control of mites, use up to 10 fl oz/100 gal water. For maximum mite control, tank mix with an acaricide.</p>				
<p>Perimeter Treatments to include Outside Surfaces</p>	<p>Aphids, Thrips, Mites, Clover Mites, Two-Spotted Spider Mites, Whiteflies, Mealybugs, Leaf- Feeding Caterpillars (i.e., Tent, Orlando, Gypsy Moth, Bagworm, Canker Worm, etc.), Ticks, Brown Dog Ticks, American Dog Ticks, Deer Ticks, Fleas, Ants, Fire Ants, Crickets, Armyworms, Cutworms, Springtails (Collembola)*, Adelgids, Mosquitoes, Chiggers*, Earwigs, Lygus Plant Bugs Beetles (Flea, Cucumber, Elm Leaf) Leafhoppers Weevils* (Strawberry Root, Obscure Root, Blackvine Root) Psyllids, Millipedes* American Cockroaches Smoky Brown Cockroaches, Oriental Cockroaches, Boxelder Bugs, Spiders, Silverfish, Firebrats, Chinch Bugs, Asian Lady Beetles*, Sowbugs, Glassy-Wing Sharpshooters</p> <p>* Except in California</p>	<p>4-10 fl oz</p>	<p>0.2-0.5 fl oz (1.3-3 teaspoons)</p>	<p>Apply MAVRIK AQUAFLOW to perimeters and outside surfaces of residential, non-commercial, and commercial buildings such as industrial offices, institutional buildings, private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, storage sheds, patios</p> <p>Apply MAVRIK AQUAFLOW using a low pressure, coarse fan spray, applied in even bands to surfaces and perimeters where pests are a problem. Apply the spray mixture at a volume equivalent to 5 gallons per 1,000 square feet. Depending on the severity of pest infestations, a re-treatment may be necessary at 7-14 days. Apply no more than 4 times per month or no more than 20 times per year. Typical re-treatment interval is 28 days.</p> <p>Perimeter applications are made to a band of soil and/or vegetation 6 to 10 feet wide around and adjacent to the structure. Higher volumes of water may be needed if mulch or leaf litter is present or foliage is dense. These sites may include refuse dumps, vegetation areas, soil, trunks of woody ornamentals, and fence lines adjacent to or around the structure, and other areas near a structure where pests congregate or have been seen.</p> <p>Surface applications can be made to all outside surfaces of structures listed above and includes porches, window and door frames, eaves, and foundations. Apply as a crack and crevice treatment using a pinstream nozzle to pest harborage sites. Refrain from applying to the point of runoff.</p>

CROP/SITE	PEST	AMOUNT MAVRIK AQUAFLOW IN WATER TO MAKE		SPECIFIC DIRECTIONS
		100 GALLONS	5 GALLONS	
Containerized Nursery Stock- Outdoors	Aphids, Thrips, Mites, Clover Mites, Whiteflies, Leaf-Feeding Caterpillars (i.e., Tent, Orlando, Gypsy Moth, Bagworm, Canker Worm, etc.) Lygus Plant Bugs Beetles (Flea, Cucumber, Elm Leaf) Leafhoppers, Earwigs, Ants, Crickets, Armyworms, Cutworms, Sowbugs, Chiggers*, Millipedes*, Springtails (Collembola)* Adelgids, Mosquitoes, Glassy -Wing Sharpshooters * Except in California	4-10 fl oz	0.2-0.5 fl oz (1.3-3 teaspoons)	For application to small, immature ornamental plants in containers up to 5 gallons in size: Mix 4-10 fl oz of MAVRIK AQUAFLOW in 100 gallons of water. Apply an equivalent of 100 gallons of spray solution per 20,000 square feet of growing area using a low pressure, coarse fan spray application. Spray interval is typically 14-28 days but not more than 4 applications per month. Do not exceed 16 applications per year. Use sufficient water to thoroughly cover the surfaces to be treated. Use lower rates for light to moderate insect pressure. Apply spray to foliage, around base of plant, and to the main trunk or stems.
Woody and Herbaceous Ornamentals- Outdoors	Aphids, Thrips, Mites, Clover Mites, Whiteflies, Leaf-Feeding Caterpillars (i.e., Tent, Orlando, Gypsy Moth, Bagworm, Canker Worm, etc.) Lygus Plant Bugs Beetles (Flea, Cucumber, Elm Leaf) Leafhoppers, Earwigs, Ants, Crickets, Armyworms, Cutworms, Sowbugs, Chiggers*, Millipedes*, Springtails (Collembola)* Adelgids, Mosquitoes, Glassy -Wing Sharpshooters, Strawberry Root Weevils*, Obscure Root Weevils*, Blackvine Root Weevils*	4-10 fl oz	0.2-0.5 fl oz (1.3-3 teaspoons)	Make applications to large, mature ornamental specimen plants or plants in containers larger than 5 gallons in size. Mix 4-10 fl oz of MAVRIK AQUAFLOW in 100 gallons of water (equivalent to 0.2- 0.5 fl oz or 1.3-3 teaspoons per 5 gallons of water). Apply 100 gallons of spray solution per equivalent of 20,000 square feet of growing area using a low pressure, coarse fan spray applications. Spray interval is typically 14-28 days but not more than 4 applications per month. Do not exceed 24 applications per year. Use sufficient water to thoroughly cover the surfaces to be treated. Use lower rates for light to moderate insect pressure. Spray on ground around base of tree or shrub and/or up main trunk or stem and foliage.

110919

CROP/SITE	PEST	AMOUNT MAVRIK AQUAFLOW IN WATER TO MAKE		SPECIFIC DIRECTIONS
		100 GALLONS	5 GALLONS	
Eugenia and Pepper Tree	Psyllids * Except in California	5-10 fl oz	0.25-0.5 fl oz (1.5-3.0 teaspoons)	Apply 5-10 fl oz of MAVRIK AQUAFLOW per 100 gallons of water to protect new foliage, depending on pest pressure. Use the higher rate of MAVRIK AQUAFLOW for heavier infestations of psyllids. Apply 2 applications at 14 day intervals under heavy reinfestation pressure or if new foliage is present. If pruning is needed, prune the entire hedge, then delay application until adults and new shoots are present.
Ant Mound Treatment	Fire Ants and other mounding ant pests	10 fl oz	0.5 fl oz (3 teaspoons)	<u>Individual ant mound treatment:</u> Apply as drench using 1 gallon of spray solution per mound to control mounding ant colonies, including the fire ant. Thoroughly mix 2/3 teaspoons (0.1 fl oz.) of MAVRIK AQUAFLOW with 1 gallon of water. Drench the mound using 1 gallon of dilution per mound. Apply the dilution starting on the perimeter of the mound and ending at the center of the mound until the entire mound is wet but not to the point of runoff.

Indoor applications:


Greenhouse Applications	Aphids, Thrips, Mites, Whiteflies, Earwigs, Beetles (Flea, Cucumber, Elm Leaf), Leafhoppers, Springtails (Collembola)*, Adelgids, Leaf-Feeding Caterpillars (i.e. Tent, Orlando, Gypsy Moth, Bagworm, Canker Worm, etc.), Lygus Plant Bugs * Except in California	4-10 fl oz	0.2-0.5 fl oz (1.3-3 teaspoons)	<u>Broadcast, fogger, or bench applications.</u> Mix 4-10 fl oz of MAVRIK AQUAFLOW in 100 gallons of water (equivalent to 0.2-0.5 fl oz per 5 gallons of water). Use a higher rate of MAVRIK AQUAFLOW for more residual activity. Apply 100 gallons of spray solution which will typically cover 20,000 square feet of greenhouse space using a low pressure, coarse fan spray for bench or broadcast applications. Spray interval is typically 14-28 days but not more than 4 applications per month or 16 applications per crop. When using ElectroStatic Spraying Systems (ESS®), PulsFog®, cold foggers, or other ULV delivery systems, apply the labeled amount of MAVRIK AQUAFLOW over the square footage with the equipment manufacturer's requirement of water. Follow manufacturer's calibration procedures.
-------------------------	--	------------	---------------------------------	--

CROP/SITE	PEST	AMOUNT MAVRIK AQUAFLOW IN WATER TO MAKE		SPECIFIC DIRECTIONS
		100 GALLONS	5 GALLONS	
Indoor Ornamentals (Interior Plantscapes)	Aphids, Whiteflies, Thrips, Mites, Earwigs, Beetles (Flea, Cucumber, Elm Leaf), Leafhoppers, Springtails (Collembola)*, Adelgids, Leaf-Feeding Caterpillars (i.e. Tent, Orlando, Gypsy Moth, Bagworm, Canker Worm, etc.) Lygus Plant Bugs * Except in California	4-10 fl oz	0.2-0.5 fl oz (1.3-3 teaspoons)	Mix 0.1–0.25 fl oz of MAVRIK AQUAFLOW in 2.5 gallons of water (equivalent to 0.2–0.5 fl oz per 5 gallons of water). Apply 5 gallons of spray solution per 1,000 square feet using a low pressure, coarse fan spray for plantscape applications.
Flower and Foliage Cuttings (for Dipping)	Mealybugs, Aphids, Two-Spotted Spider Mites	5 fl oz	0.25 fl oz (1.5 teaspoons)	Immerse basket of cuttings into a diluted solution of MAVRIK AQUAFLOW until thoroughly wet (usually 1 minute or less).

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

ESS® is a registered trademark of ElectroStatic Spraying Systems.
PulsFog® is a registered trademark of Dramm International.
MAVRIK AQUAFLOW® is a registered trademark of Wellmark International.

Wellmark International
1501 East Woodfield Road 200W
Schaumburg, IL 60173
Made in USA



Box 19

[] optional text

ZOËCON®

**OUTDOOR FLEA/ANT* CONTROL
YARDEX® INSECTICIDE**

Alternate brand name: MAVRIK® Perimeter

For commercial use only on outdoor plants in ornamental plantings, plantscapes (including trees and shrubs), Also for (Perimeter Treatment) around buildings and outside surfaces

Small, easy to use 8 oz. Bottle

Contains tau-fluvalinate

Ideal for applications without the worry of watering-in

Perimeter treatment against a wide range of insects including mosquitoes and fire ants

ACTIVE INGREDIENT:

Tau-fluvalinate (CAS #102851-06-9)..... 22.3%

OTHER INGREDIENTS: 77.7%

TOTAL **100.0%**

YARDEX is a flowable formulation containing 2 pounds tau-fluvalinate per gallon (240 g/L)

EPA Registration No. 2724-478

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN

CAUTION

READ COMPLETE DIRECTIONS AND PRECAUTIONS INSIDE

SHAKE WELL BEFORE USING

Not use on plants or trees being grown for sale or other commercial use, for commercial seed production, for the production of timber or wood products, or for research purposes. For use only on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks and grounds.

* Except in California

Net Contents

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Harmful if swallowed, inhaled, or absorbed through skin. **Avoid Breathing Spray Mist.** Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. For 24 Hour Medical or Chemical Emergency Assistance, call Wellmark International [Phone Number]	

Sensitive individuals may temporarily experience an itching, burning, or tingling sensation, with or without a rash, following exposure. These symptoms will usually subside without requiring medical treatment. Avoid hand or sleeve-to-face contact. Prior to exposure or after washing, an application of corn oil to exposed skin may reduce these symptoms. Certain persons may be sensitive to YARDEX®'s fine spray particles.

If workers enter the treated area and/or contact treated surfaces:

- Pesticide residues that get on their skin may cause itching or irritation that can be severe, and
- They should avoid skin contact with treated foliage and/or surfaces, and
- They should wash the affected skin immediately if irritation begins to occur.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- Protective clothing
- Rubber gloves while handling; workers in frequent contact with treated foliage should wear light barrier gloves to reduce excess exposure.
- For sensitive persons, use a NIOSH approved respirator with any R, P, or HE filter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic organisms. This product is toxic to honey bees if bees are exposed to direct application. However, dried residues of this product are non-toxic to honey bees. Treat during non-foraging periods to minimize adverse effects. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Use with care when applying in areas adjacent to any body of water. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not apply when weather conditions favor drift from the target area. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

PHYSICAL/CHEMICAL HAZARDS

Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store away from direct sunlight or heat. Pesticides must be stored in a secured area away from other products, food, or feed. In case of spill or leak, soak up with soil or other absorbent material. Do not use caustic agents.

Pesticide Disposal: Wastes resulting from the use of YARDEX may be disposed of on site or at any approved waste disposal facility.

Container Disposal: Triple rinse container (or equivalent). Puncture and dispose of in a sanitary landfill, by incineration or if allowed by state and local authorities by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GENERAL PRECAUTIONS/RESTRICTIONS

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Keep unprotected people, children and pets out of treated area until spray has dried. Some sensitive pets eating or coming in contact with freshly treated plants may demonstrate vomiting and respiratory distress.

Do not apply this product by aerial application or through any type of irrigation system.



GENERAL DIRECTIONS


YARDEX controls insects and controls or suppresses mite pests when used at labeled rates. YARDEX works primarily through contact action, good spray coverage is necessary. Treat when pests are immature or at a susceptible stage and populations are manageable, before damage occurs. Repeat applications as needed, according to the specific use directions given in this label. Residues of YARDEX can continue to control certain insect pests for a week or longer. Growth dilution, climatic variables, and pest habits determine retreatment needs.

Dilute YARDEX in sufficient water to give thorough spray coverage [based on previous experience]. For best results under varying conditions, a range of use rates is recommended for most uses. Select the higher rates when pest pressure or damage potential is severe. The lower rates often give acceptable economic results. [Buffer spray water to pH 5-7, if necessary.]

For severe mite infestations, use the highest rate of YARDEX and use an acaricide in an alternate treatment program. [The use of a spreader-sticker, insecticidal soaps, or piperonyl butoxide may improve control of mites and some insects, such as, whiteflies, aphids, thrips, and mealybugs.] When YARDEX is mixed with wettable powders, add powders to the tank first using good agitation. Check for physical and biological compatibility prior to use.

Dried residues of YARDEX are non-toxic and non-repellant to honey bees. Treat during non-foraging periods to minimize adverse effects.

Consult your local extension office, pest management consultant, or a Wellmark International specialist for guidance in selecting a rate and timing schedule or mixing combination that has demonstrated good performance under specific conditions.



17619

CROP/SITE	PEST	AMOUNT YARDEX IN WATER TO MAKE		SPECIFIC DIRECTIONS
		5 GALLONS	1 GALLON	
<p>For Outdoor Applications: Use lower rates and greater re-treatment intervals for light to moderate insect pressures. Use higher rates and shorter re-treatment intervals for green peach aphids, unexposed insects and heavy pressures of other insects. Mite suppression can be obtained using the lower rate; for more residual control of mites, use up to 0.5 fl oz/5 gallons water [0.1 fl oz/1 gallon of water].</p>				
Perimeter Treatments to include Outside Surfaces	<p>Aphids, Thrips, Mites, Clover Mites, Two-Spotted Spider Mites, Whiteflies, Mealybugs, Leaf- Feeding Caterpillars (i.e., Tent, Orlando, Gypsy Moth, Bagworm, Canker Worm, etc.), Ticks, Brown Dog Ticks, American Dog Ticks, Deer Ticks, Fleas, Ants, Fire Ants, Crickets, Armyworms, Cutworms, Springtails (Collembola)*, Adelgids, Mosquitoes, Chiggers*, Earwigs, Lygus Plant Bugs Beetles (Flea, Cucumber, Elm Leaf) Leafhoppers Weevils* (Strawberry Root, Obscure Root, Blackvine Root) Psyllids, Millipedes* American Cockroaches Smoky Brown Cockroaches, Oriental Cockroaches, Boxelder Bugs, Spiders, Silverfish, Firebrats, Chinch Bugs, Asian Lady Beetles*, Sowbugs, Glassy-Wing Sharpshooters</p> <p>* Except in California</p>	0.2-0.5 fl oz (1 1/3-3 teaspoons) (6-15 ml)	.04-0.1 fl oz (1/3-2/3 teaspoons) (1.2-3 ml)	<p>Apply YARDEX to perimeters and outside surfaces of residential, non-commercial, and commercial buildings such as industrial offices, institutional buildings, private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, storage sheds, patios</p> <p>Apply YARDEX using a low pressure, coarse fan spray, applied in even bands to surfaces and perimeters where pests are a problem. Apply the spray mixture at a volume equivalent to 5 gallons per 1,000 square feet. Depending on the severity of pest infestations, a re-treatment may be necessary at 7-14 days. Apply no more than 4 times per month or no more than 20 times per year. Typical re-treatment interval is 28 days.</p> <p>When using cold foggers or other ULV delivery systems for perimeter applications, apply the labeled amount of YARDEX over the square footage with the equipment manufacture's requirement of water. Follow manufacture's calibration procedures.</p> <p>Perimeter applications are made to a band of soil and/or vegetation 6 to 10 feet wide around and adjacent to the structure. Higher volumes of water may be needed if mulch or leaf litter is present or foliage is dense. These sites may include refuse dumps, vegetation areas, soil, trunks of woody ornamentals, and fence lines adjacent to or around the structure, and other areas near a structure where pests congregate or have been seen.</p> <p>Surface applications can be made to all outside surfaces of structures listed above and includes porches, window and door frames, eaves, and foundations. Apply as a crack and crevice treatment using a pinstream nozzle to pest harborage sites. Refrain from applying to the point of runoff.</p>

CROP/SITE	PEST	AMOUNT YARDEX IN WATER TO MAKE		SPECIFIC DIRECTIONS
		5 GALLONS	1 GALLON	
Woody and Herbaceous Ornamentals-Outdoors	Aphids, Thrips, Mites, Clover Mites, Whiteflies, Leaf-Feeding Caterpillars (i.e., Tent, Orlando, Gypsy Moth, Bagworm, Canker Worm, etc.) Lygus Plant Bugs Beetles (Flea, Cucumber, Elm Leaf) Leafhoppers, Earwigs, Ants, Crickets, Armyworms, Cutworms, Sowbugs, Chiggers*, Millipedes*, Springtails (Collembola)* Adelgids, Mosquitoes, Glassy -Wing Sharpshooters, Strawberry Root Weevils*, Obscure Root Weevils*, Blackvine Root Weevils*	0.2-0.5 fl oz (1 1/3-3 teaspoons)	.04-0.1 fl oz (1.2-3 ml) (1/3-2/3 teaspoons)	Make applications to large, mature ornamental specimen plants or plants in containers larger than 5 gallons in size. Mix 0.2-0.5 fl oz or 1 1/3-3 teaspoons of YARDEX in 5 gallons of water. Apply 5 gallons of spray solution per equivalent of 1,000 square feet of growing area using a low pressure, coarse fan spray applications. Spray interval is typically 14-28 days but not more than 4 applications per month. Do not exceed 24 applications per year. Use sufficient water to thoroughly cover the surfaces to be treated. Use lower rates for light to moderate insect pressure. Spray on ground around base of tree or shrub and/or up main trunk or stem and foliage.
Eugenia and Pepper Tree	Psyllids * Except in California	0.25-0.5 fl oz (1 1/2-3. teaspoons)	0.05-0.1 fl oz (1/3-2/3 teaspoons)	Apply 0.25-0.5 fl oz (1 1/2-3 teaspoons of YARDEX per 5 gallons of water to protect new foliage, depending on pest pressure. Use the higher rate of YARDEX for heavier infestations of psyllids. Apply 2 applications at 14 day intervals under heavy reinfestation pressure or if new foliage is present. If pruning is needed, prune the entire hedge, then delay application until adults and new shoots are present.
Ant Mound Treatment	Fire Ants and other mounding ant pests	0.5 fl oz (15 ml) (3 teaspoons)	0.1 fl oz (3 ml) (2/3 teaspoons)	<u>Individual ant mound treatment:</u> Apply as drench using 1 gallon of spray solution per mound to control mounding ant colonies, including the fire ant. Thoroughly mix 2/3 teaspoons (0.1 fl oz/3 ml) of YARDEX with 1 gallon of water. Drench the mound using 1 gallon of dilution per mound. Apply the dilution starting on the perimeter of the mound and ending at the center of the mound until the entire mound is wet but not to the point of runoff.

For more information or in case of emergency, call [phone number] or visit our Web site [URL]

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

MAVRIK®, YARDEX® Insecticide and ZOËCON® are registered trademarks of Wellmark International.

Wellmark International
1501 East Woodfield Road 200W
Schaumburg, IL 60173

Made in USA

NOTIFICATION
FEB 21 2009

