

2724-483

7/16/2002

1/6

[ ] indicates optional statements

### SANDOZ 9404 SPRAY

[Treats # square feet]  
 [Part of the (Brand Name) [Step #][Guarantee] Pet Protection Program]  
 Contains the Synthetic Pyrethroid **PERMANONE™**  
 Contains [With] **PRECOR®** [**VIGREN™**] Insect Growth Regulator plus permethrin, an adulticide  
 Stops Hatching [flea] Eggs (Larvae) From Developing Into Adult Fleas For 210 Days [30 weeks][7 months]  
 Controls Residential, Industrial and Garden Insects  
 Kills [Controls] Both Adult and Preadult (Flea Eggs & Larvae) Fleas & Ticks for [7 months][210 days][30 weeks]  
 Contains an Insect Growth Regulator – Kills Flea Eggs  
 Kills adult ticks  
 For Carpet/Upholstery  
 Satisfaction 100% guaranteed  
 Kills Preadult (Larvae) Fleas Before They Grow Up to Bite You  
 Leaves No Lingering Odor (Bad Smell), No Stains, No Sticky Mess  
 Effectively Breaks The Flea Life Cycle  
 For Use On Dogs and On Premises to Control Fleas and Ticks  
 Controls Fleas and Ticks on Dogs for 14 Days  
 A Water Based Aerosol  
 Kills Lice and Their Eggs  
 For Use On Bedding, Furniture and Other Inanimate Objects Infested with Lice  
 Easy to Apply  
 Prevents Reinfestation and Flea Build-Up For 210 Days [30 weeks][7 months]  
 Not for Use on Humans

**ACTIVE INGREDIENTS:**

(S)-Methoprene (CAS #65733-16-6)	0.09%
Permethrin (CAS #52645-53-1)	0.50%

**OTHER INGREDIENTS:**

99.41%

**TOTAL:**


...100.00%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See back panel for additional precautionary statements

Net Contents:



**READ ENTIRE LABEL BEFORE USE  
USE ONLY ON DOGS**

2/6

**DIRECTIONS FOR USE:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not use in food areas of food handling establishments, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods are served such as dining rooms, but excluding areas where foods may be prepared or held. In the home, all food processing surfaces and utensils should be covered during treatment or thoroughly washed before use. Exposed food should be covered or removed. Nonfood areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling). Avoid contamination of feed and foodstuffs.

**SHAKE WELL BEFORE USING**

**This product is not for use on humans.**

A single can (16 oz.) treats 2000 square feet. Remove protective cap, hold container *upside down* and spray from a distance of 2 - 3 feet. Remove pets and birds and cover and unplug aquariums before spraying. Vacate room after treatment and ventilate before reoccupying. Do not allow children or pets to contact treated areas until surfaces are dry.

**Houseflies, Mosquitoes, Gnats, Wasps, Hornets, Bees, Black Flies, Skipper Flies, Blow Flies and Small Flying Moths:** Close all doors and windows and spray room at the rate of 2 to 3 seconds per 1000 cubic feet. Direct spray to all parts of the room, especially windows and other light sources which attract these insects. Keep nozzle in constant motion.

**Cockroaches, Waterbugs, Palmetto Bugs, Silverfish, Firebrats, Crickets, Scorpions, Spiders, Millipedes, Centipedes, Sowbugs and Pillbugs:** Spray into hiding places such as cracks, crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible.

**Ants:** Spray trails, nests and points of entry. Spray on ants where possible.

**Indian Meal Moths, Angoumois Grain Moths:** Close doors and windows, direct spray into all parts of room especially around stored food product containers, pallets and darkened areas. Spray 7 to 10 seconds for each 1000 cubic feet of space. Keep area closed for 15 minutes following treatment.

**Exposed stages of Confused Flour Beetles, Saw-Toothed Grain Beetles, Red Flour Beetles, Rice Weevils, Granary Weevils, Grain Mites:** Spray into cracks and crevices, processing and handling equipment, around cartons or containers or other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Infested stored food products in the home should be disposed of.

Carpet Beetles: Spray directly on beetles, floors, baseboards, shelves and carpets. Bedbugs: Spray mattresses, particularly around tufts and seams. Take beds apart and spray into all joints. Treat baseboards, moldings and floors. Clothes Moths and Carpet Beetles on wool and other proteinaceous fibers: Remove infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing, infested articles may be treated by spraying lightly, keeping nozzle at least three feet from fabric to avoid staining.

In Animal Quarters and Milk Rooms: For rapid control of House flies, Mosquitoes, Gnats, Wasps, Hornets, Bees and Small-flying Moths, close windows and doors and spray at the rate of 2 seconds per 1000 cubic feet of space. For best results, keep area closed for 10 minutes after spraying. Do not remain in treated area and ventilate after treatment is completed.

To Treat Livestock: To protect beef cattle, dairy cattle, goats, sheep, hogs and horses from attacks of Stable Flies, Horse Flies, Deer Flies, Face Flies, House Flies, Horn Flies, Mosquitoes, and Gnats: Spray about 3 seconds on each side being careful to spray back, withers and forelegs thoroughly. To protect from Face Flies, spray the face and head, but do not spray into eyes. Repeat treatment when flies are troublesome. For Swine: Do not ship animals for slaughter within 5 days of treatment.

To control Blood Sucking Lice: Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required. To control Poultry Lice, spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist.

For control of Bedbugs and Mites in poultry houses, spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

#### **Fleas (Adults and Larvae) and Ticks (Adults):**

##### **FOR USE IN THE HOME**

For best results, pay special attention to areas your pets frequent.

Avoid thoroughly wetting the area that needs treating.

To kill adult fleas, flea larvae and adult ticks in your home, hold container upside down 2-3 feet from surface, point actuator towards the surface and push button.

#### **For Surface / Carpet Spray:**

Vacuum thoroughly before spraying.

A test application should be made to upholstery or drapery fabrics in an inconspicuous place before use. Apply uniformly with a sweeping motion to rugs, carpets, drapes and all surface of upholstered furniture. Treat surfaces uniformly from one end to the other. Only complete coverage of carpets, upholstery, drapes and other fabrics will provide thorough flea/tick elimination.

#### **For Use in Pet Areas:**

Pay special attention to areas where pets frequent, such as in and around sleeping and resting areas. Treat pet bedding and other pet resting areas as these are primary hiding areas for fleas/ticks. For best results, pets should be treated before they return to the treated areas.

**FOR USE ON DOGS**

To Control Fleas, Ticks and Lice on dogs: Starting at the tail, stroke against the lay of the hair, spraying the parted fur in short bursts, directly behind the hand to insure penetrating the coat. Dampen the fur up to the head. Do not treat face, eyes or genitals. Repeat application weekly if necessary to control infestations. **NOTE:** Do not use on puppies under 12 weeks old.

**To kill Lice and Louse eggs:** Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Hold container upright with nozzle away from you. Depress valve and spray from a distance of 8 to 10 inches. Spray each square foot for three seconds. Spray only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all sprayed articles to dry thoroughly before use. If lice infestations should occur on humans, use RID Lice Killing Shampoo. If lice infestations should occur on humans, use an EPA-registered product labeled for use on humans. If lice infestations should occur on humans, consult either your physician or pharmacist for a product for use on humans.

**For the Control of Garden Insects:** For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray, as well as Whiteflies, Aphids, exposed Thrips, Spiders, Red Mites and Leafminers. May be used on plants, as well as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas.

May also be used on other plants such as African violets, azaleas, begonias, camellias, carnations, chrysanthemums, delphiniums, dogwood, English Ivy, euonymus, fuschia, geraniums, crassula, grape vine, Kentia palm, laurel, marigolds, rhododendrons, rubber plants, snapdragons, stocks, wandering Jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spiders and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Army Worms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Spray in one second bursts. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90F. Contact spray insects directly whenever possible.

**STORAGE and DISPOSAL:** **Storage:** Store in a cool, dry area. Do not transport or store below 32°F. **Disposal:** Do Not Puncture or Incinerate! **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMAL - CAUTION:** Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling and before smoking or eating. Consult a veterinarian before using this product on debilitated, aged, pregnant, animals on medication, or nursing animals. Sensitivities may occur after using ANY pesticide product for pets. If signs of sensitivity occur, bathe your pet with mild soap and rinse with large amounts of water. If signs continue, consult a veterinarian immediately.

<b>FIRST AID</b>	
Call a poison control center or doctor immediately for treatment advice	
If swallowed	<ul style="list-style-type: none"> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or a doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-[phone number] for emergency medical treatment information.	

[Note: First Aid statement paragraph format may be used if space is limited.]

**PHYSICAL OR CHEMICAL HAZARDS:** Contents under pressure. Do not use or store near heat or open flame. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

*Contains no CFC or other ozone-depleting substances. Federal regulations prohibit CFC propellants in aerosols.*

Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and handling are contrary to label instructions.

For information or in an emergency call 1-800 (###-####) or visit our Web site: [web site name]

PRECOR<sup>®</sup> Insect Growth Regulator and vIGRen<sup>®</sup> Insect Growth Regulator are registered trademarks of Wellmark International.

PERMANONE<sup>™</sup> is a trademark of AgrEvo Environmental Health.

Wellmark International  
Schaumburg IL 60173

EPA Reg. No. 2724-483  
Made in USA

EPA Est. No.