

ANNUAL SOW THISTLE
ARROWHEAD
BEGGAR TICKS
BINDWEED
BUCKHORN
BULL THISTLE
BUTTERPRINT
CATNIP
CHECKORY
CLOVER
DANDELION
DOCKS

FANWEED
FIGWORT
FOUR O'CLOCK
FRENCH WEED
GROUND IVY
HEDGE BINDWEED
HENBIT
HORSETAIL
JIMSONWEED
KNOTWEEDS
KOCHIA
LAMBSQUARTERS
LAWN PENNYWORT

LOCO WEEDS
MALLOW
MARESTAIL
MORNING GLORY
MUSTARD
NETTLES
PIG WEED
PLAINTAINS
PRICKLY LETTUCE
PUNCTURE VINE
PURSLANE
RAGWEEDS
RUSSIAN THISTLE

SHEPHERD'S PURSE
SMART WEED
SOW THISTLE
STINKWEED
SUMACS
SUNFLOWER
WILD MUSTARD
WILD PARSNIP
WILD RADISH
WILD ROPE
WILD SWEET POTATO

#1
PESTS

Some weeds above listed as susceptible may not be completely killed but growth is so retarded by proper application that they are no longer a factor affecting growth and yield of the crop.

When conditions are such as to promote vigorous growth. Consult local agricultural authorities for recommendations for specific uses.

WARNINGS

All plant life is more susceptible to 2,4-D esters than other types of 2,4-D forms and therefore requires less. Since esters are volatile, care should be exercised to avoid vapors drifting to sensitive desirable plants. Coarse sprays are less likely to drift.

Do not allow 2,4-D or its vapors to come into contact with flowers, fruits, grapes or cotton.

Do not store 2,4-D near fertilizers, seeds, insecticides or fungicides.

Do not use 2,4-D spray equipment for any other purpose until it has been thoroughly cleaned and neutralized with ammonia.

Avoid spray drift to desirable plants.

Do not use on Bent grass, St. Augustine grass and clovers as damage may occur.

CAUTIONS

Avoid contact with eyes, skin and clothing. Do not take internally. May cause irritation. Do not contaminate irrigation ditches or water used for domestic purposes. Harmful if swallowed.

ACCEPTED
6/9/70
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTS REGISTERED
EPA UNDER NO. 24377

Comment