

HANDLING INSTRUCTIONS

DANGER. Potentially fatal if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. **DO NOT GET ON SKIN, IN EYES OR ON CLOTHING. DO NOT BREATHE SPRAY OR MIST.** wear protective gloves, clothing and goggles while spraying this product. **DO NOT TAKE INTO MOUTH. KEEP AWAY FROM FOOD AND FOOD PRODUCTS.** Do not spray fruit or truck crops later than at least one month prior to harvest. If spilled on skin, wash immediately with soap and warm water. Wash hands, arms and face after handling and before eating or smoking. Clothing that has become contaminated should be removed immediately and thoroughly washed with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Parathion protection. When used constantly during the day, the filters must be changed several times.

Decontaminate all containers with lye or alkali; wash, then puncture containers and place on private dump.

FIRST AID TREATMENT

Persons who show any contraction of eye pupils, or have headaches, nausea or other signs of illness while or soon after handling Parathion must be taken to a doctor at once. The physician should be informed that repeated treatment with atropine, to the limit of the patient's tolerance, is the most effective antidote or give 2 P. A. M. intravenously at the rate of 1 gm. for adults and .25 gm. for children.

CALL A PHYSICIAN AT ONCE

SKIN: Wash immediately with plenty of soap and water. Remove all contaminated clothing and wash before re-use.

EYES: Flush immediately with large amounts of water.

INTERNAL: If swallowed, induce vomiting by rubbing a down throat, or administer a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Place victim in a drowsy and keep quiet.

CALL A PHYSICIAN AT ONCE


NOTICE OF WARRANTY

This product is warranted to be free from mechanical defects in the container and to be of the quality and quantity stated on the label. The manufacturer is not responsible for the results of the use of this product when it is applied in violation of the directions on the label. The user assumes all liability for the results of the use of this product when it is applied in violation of the directions on the label.

SURE KILL

8-lb. Parathion-Miscible

8 LBS. ETHYL PARATHION PER GALLON
EMULSIFIABLE CONCENTRATE



ACTIVE INGREDIENTS	84%
Parathion (O, O-Dimethyl O-paranitrophenyl thiophosphate)	
XYLENE	11%
INERT INGREDIENTS	5%
TOTAL	100%

POISON DANGER -- Poisonous if

swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. **KEEP OUT OF THE REACH OF CHILDREN.** See side panels for additional required warnings and antidote statement.

Pueblo Chemical Company

611 SANTA FE DRIVE

PUEBLO, COLORADO

DIRECTIONS

Exercise extreme care in handling this product. Do not inhale or get on skin.

DIRECTIONS FOR USE

SURE KILL PARATHION must be diluted with water or oil before using.

Use in ground or airplane, custom and power sprayers. Add the required amount of Parathion to only a small quantity of water and agitate well before adding the remainder of the water required.

Spray all foliage thoroughly with the diluted spray solution. Repeat as often as necessary for complete control.

Application should be made before heads of wheat, barley and oats begin to form.

AIRPLANE SPRAYING. (Greenbug on wheat, oats, barley and sorghum). Use from 1/4 to 1/2 lb. Parathion dissolved in 2 gallons of water and apply at the rate of 2 gallons per acre.

CORN: Spider Mites and Aphids use 1/2 lb. Parathion per acre. Airplane use minimum 1 gallon spray, ground rig use minimum 4 to 5 gallon spray per acre. Do not apply less than 12 days before harvest or cutting for forage.

WHEAT, BARLEY, OATS AND SORGHUM Mites. Use 1/2 to 1 lb. Parathion per acre. Airplane use minimum 1 gallon spray, ground rig use minimum 4 to 5 gallon spray per acre. Do not apply less than 15 days before harvest on wheat, barley and oats. Do not apply less than 12 days before harvest or cutting for forage on sorghums.

BROCCOLI, BRUSSEL SPROUTS, CABBAGE, CAULIFLOWER, Cabbage, Looper and cabbage worm, aphids. Use 1/2 lb. Parathion per acre with the necessary amount of water to cover the crop. Do not apply less than 7 days before harvest.

ONION: Fleas use 1/2 lb. Parathion with the necessary amount of water per acre. Do not apply less than 15 days before harvest.

DO NOT APPLY TO VEGETABLE CROPS WITHIN 10 DAYS OF HARVEST UNLESS NOTED OTHERWISE.

DANGER: Do not use this product if the operator or a bystander is wearing clothing or shoes that are contaminated with dirt. Vacated clothing should be washed until the dirt has been completely removed. Do not use if the operator or a bystander has been drinking alcohol.