

ETHION MISCIBLE

Apply the listed amount per acre when insects first appear and repeat as required to maintain effective control. Use in sufficient water for thorough coverage, unless otherwise noted for listed crop. Coverage of upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use pattern, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in () following the crop.

When applying this material by aircraft, mix the recommended amount with sufficient water to provide a minimum of 1 gallon finished spray per acre. Where more than 2 quarts are recommended mix with sufficient water to provide a minimum of finished spray equal to twice the amount of Ethion Miscible used. Do not use in undiluted form.

BEANS (SUCCULENT AND DRY) (2): Bean Leaf Beetle, Mexican Bean Beetle, Two-spotted Spider Mite - 0.5 to 1 pt. per acre. Potato Leafhopper - 1 pt. per acre. Do not feed treated foliage to livestock.

BEANS (SUCCULENT AND DRY) (4): Lygus Bugs - 1 qt. per acre. Do not feed treated foliage to livestock.

CORN, FIELD (50): Pacific Spider Mite, Two-spotted Spider Mite - 1 qt. per acre. Do not make more than 1 application after ear formation. Do not feed treated fodder to livestock.

CUCUMBERS (0), SUMMER SQUASH (0): Serpentine Leaf Miner, Two-spotted Spider Mite - 1 pt. per acre.

EGGPLANT (0): Two-spotted Spider Mite - 0.5 to 1 pt. per acre.

MELONS (0): Melon Leaf Miner, Serpentine Leaf Miner, Two-spotted Spider Mite - 1 pt. per acre. Melon Leafhopper - 1.5 to 2 pts. per acre.

PEPPER (0), PIMENTO (0): Serpentine Leaf Miner - 1 pt. per acre. Two-spotted Spider Mite - 0.5 to 1 pt. per acre.

SORGHUM (30): Sorghum Midge, Sorghum Mite - 0.5 to 1.0 pt. per acre. For Midge, apply at first bloom and repeat 4 to 10 days later. Do not utilize for forage within 30 days of treatment. Do not apply more than 3 times per season. Some leaf burning may occur on phosphate-sensitive varieties.


TOMATOES: (2): Leaf Miner, Tropical Spider Mite, Two-spotted Spider Mite - 1 pt. per acre.

APPLES: Apple Aphid, European Red Mite, Rosy Apple Aphid, San Jose Scale - 0.6 pt. plus 2 gals. dormant oil per 100 gals. water. Apply during dormant or delayed dormant period. ("Superior" type oil of 60 or 70 second viscosity preferred for delayed dormant use).

ORNAMENTALS (GREENHOUSE): Millipedes - **SOIL TREATMENT.** Use 10 fl. ozs. in 75 gals. of water and apply to 2500 sq. ft. of bench space. Wear mask or respirator approved by U.S.D.A. for Ethion protection.

ORNAMENTALS (OUT-OF-DOORS): Two-spotted Spider Mite - 0.5 pt. per 100 gals. May cause some needle drop on Anderson Yew.

Rev. 2-18-71



ETHION MISCIBLE

ACTIVE INGREDIENTS:

*Ethion (O, O, O', O'-Tetraethyl
S, S'-Methylene bisphosphorodithioate 46.5%
Xylene 35 %
Petroleum Distillate 10 %

INERT INGREDIENTS 8.5%

TOTAL 100.0%

*U.S. Patent No. 2,873,228
Contains 4 lb Ethion per gallon

WARNING -- KEEP OUT OF THE REACH OF CHILDREN. Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. See side panels for additional required warnings and antidote statement.

Manufactured by

PUEBLO CHEMICAL & SUPPLY CO.

611 SANTA FE DRIVE PUEBLO, COLORADO

Contains _____ Gal.

ACCEPTED
APR 19 1971 | 2737-25

ORNAMENTAL TURF Weevils: Use 0.33 to 1 gal. of water or 1.67 to 3.33 gal. To treat one acre, use 1.67 to 3.33 gal. It is important that the spray reach the insects. It helps prevent. Not for use or storage in

Keep children and animals thoroughly watered into Workers entering treated clothing.

Avoid contact with skin. Wash after handling.

Do not contaminate or

Persons who show a nausea or other signs of must be taken to a doctor.

SKIN: Wash immediately contaminated clothing and

EYES: Flush immediately

INTERNAL: If swallowed (throat) or administer a repeat until vomit is clear.

CALL PH

Atropine is antidotal. 2-P in conjunction with Atropine

USA

Do not use if material is Do not use or store near

Do not store at temperature

This product is toxic to water. Birds feeding on turf

only as specified on this

This product is toxic to actively visiting the area.

Rinse equipment and or burying at least 18 inch

supplies. Do not reuse or wastes. Do not burn.

DO NOT ALLOW THIS A

ING CROPS OR NON-CR

AT A TIME OTHER TH

RECTIONS BECAUSE A

LEGAL RESIDUES OR

OCCUR.

WARRANTY OF

SELLER warrants that the

tion and is reasonably fit

used in accordance with

and BUYER assumes the

SELLER makes no other

other express or implied

ABILITY, and no agent or

writing with a specific re

SELLER's liability for an

price of the material as to