

EPA Reg. No. 2737-36

Net Content

**READ ENTIRE LABEL  
BEFORE USING THIS PRODUCT**  
GENERAL INFORMATION

AMINE 4 LB is a selective herbicide for control of many broadleaf weeds in certain crops, around buildings, along fences, roads and pastures. A partial list of weeds controlled by AMINE 4 LB is shown below.

|            |              |             |
|------------|--------------|-------------|
| Broadleaf  | Kochia       | Pasture     |
| Carpetweed | Lambquarters | Ragweed     |
| Cocklebur  | Mustards     | Synthetic   |
| Dandelion  | Morningglory | Toxic       |
| Dock       | Nutsedge     | Wild oat    |
| Goldenrod  | Peppercorn   | Wild radish |
| Knotweed   |              |             |

NOTE: Local conditions, crop varieties and application methods may vary and may affect use of this herbicide. Consult your local extension agent or state or federal regulatory agencies for recommendations and restrictions.

Aerial application may be used for control of weeds in certain crops where there would be no danger of drift to susceptible crops. Applications should only be made by application in accordance with the use of 2,4-D formulations. Regulations may vary with application of herbicides are in effect in many states. Consult local regulatory agencies concerning requirements before making applications.

TREATING SMALL AREAS: One tablespoon of AMINE 4 LB in 1 1/2 gallons of water is about equal to 1 quart of 100 gals.

When stored at temperatures below freezing, it may be necessary to warm contents to 40 F and mix thoroughly before using.

**DIRECTIONS FOR USE**

PREPARATION OF THE SPRAY: Fill sprayer with water to required amount of water. Then add the amount of AMINE 4 LB and mix thoroughly. The amount of water required for low volume applications may vary from 5 to 25 gallons per acre. For high volume applications, 100 gallons or more of water per acre is recommended. In any case, use the same amount of 2,4-D recommended per acre.

TIME OF APPLICATION: Best results are obtained when AMINE 4 LB is used on young weeds that are actively growing. Application of lower rates to susceptible annual weeds is usually sufficient. For perennial weeds and other resistant types, such as in very dry areas where it is difficult to use high rates, AMINE 4 LB may be used as a selective spray on crops in the early stages of growth of the crop and be considered. Some woody plants and weeds are listed for control. For additional applications may be necessary.

SMALL GRASS (Wheat, Barley, Rye): Apply AMINE 4 LB in 1 1/2 to 2 gallons of water for uniform coverage on small grasses. Apply to 1/4 to 1/2 inch of growth (4 to 8 inches tall) but before head emergence. Do not apply to crop until after the 2nd or 3rd leaf has emerged. Crop injury may result if applied after the 2nd or 3rd leaf has emerged. Do NOT use on grain under any conditions, such as alfalfa or clover, except where the label specifically indicates. DO NOT graze or feed from treated areas until 2 weeks after treatment. See chart for recommended rates.

For late season control of weeds in certain crops, apply AMINE 4 LB in 1 1/2 to 2 gallons of water to 1/4 to 1/2 inch of growth. Do not apply to crop until after the 2nd or 3rd leaf has emerged. Do NOT graze or feed from treated areas until 2 weeks after treatment. See chart for recommended rates.

CORN: POST-EMERGENCE: Apply AMINE 4 LB in 1 1/2 to 2 gallons of water to 1/4 to 1/2 inch of growth. Do not apply to crop until after the 2nd or 3rd leaf has emerged. Do NOT graze or feed from treated areas until 2 weeks after treatment. See chart for recommended rates.

DO NOT apply to corn until after the 2nd or 3rd leaf has emerged. Do NOT graze or feed from treated areas until 2 weeks after treatment. See chart for recommended rates.

**High**  
**ACCEPTED PLAINS**  
4 MAR 1980

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2737-36

**AMINE 4 LB.  
Weed Killer**

Contains 3.8 Pounds 2,4-D Acid Equivalent Per Gallon

|   |        |
|---|--------|
| ACTIVE INGREDIENT:                                    |        |
| Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* | 46.5%  |
| INERT INGREDIENTS:                                    | 53.5%  |
| TOTAL   | 100.0% |

\*Isomer Specific by AOAC Method No. 6 D01-5  
\*Equivalent to 2,4-Dichlorophenoxyacetic Acid 38.6%.

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN.  
See Side Panels For Additional Cautions.

MANUFACTURED BY  
Pueblo Chemical & Supply Co.  
631 Santa Fe Drive • Pueblo, Colorado

FORUM: Apply AMINE 4 LB when forum plants are 6 to 12 inches tall. Do not apply to forum plants after they have begun to flower. Do NOT apply to forum plants after they have begun to flower. Do NOT apply to forum plants after they have begun to flower.

SOYBEAN CANE: Apply as a pre-emergence application before cane emergence and throughout the growing season. Do NOT apply to soybean cane after it has emerged.

**Amount of AMINE 4 LB. per Acre**

| Crop (See Detailed Directions Above) | For average Conditions |
|--------------------------------------|------------------------|
| Wheat, Barley, Rye - Annual Weeds    | 1 to 1.5 pt            |
| Wheat, Barley, Rye - Perennial Weeds | 1 pt                   |
| Corn - Post-emergence                | 1 to 1.5 pt            |
| Sorghum - Post-emergence             | 1 pt                   |
| Soybean Cane                         | 1 to 2 quarts          |

PASTURES: To control many broadleaf weeds in pastures, apply AMINE 4 LB in 1 1/2 to 2 gallons of water per acre. Do NOT apply to pastures where there are young calves or other animals. Do NOT apply to pastures where there are young calves or other animals.

LAWNS, GOLF COURSES, AND SIMILAR TURF: Apply 2 to 3 pints of AMINE 4 LB per 1000 sq ft of water to provide uniform coverage. Do NOT apply to lawns, golf courses, and similar turf where there are young children or other animals.

GENERAL WEED CONTROL: Apply AMINE 4 LB in 1 1/2 to 2 gallons of water to 1/4 to 1/2 inch of growth. Do NOT apply to crops until after the 2nd or 3rd leaf has emerged.

**CAUTION**

Harmful if swallowed or absorbed through the skin. CAUSES IRRITATION OF EYES AND SKIN. DO NOT get in eyes. In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention. DO NOT allow contact with skin or clothing. Wash exposed skin gently with plenty of soap and water. DO NOT store near fertilizers, seeds, insecticides or fungicides. DO NOT contaminate irrigation ditches or water used for irrigation or domestic purposes. Use care to avoid spray drift to 2,4-D susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees, and ornamentals. Trace amounts of spray drift may cause severe injury to both growing and dormant plants. Coarse sprays are less likely to drift. Spray only on calm days using low pressure and lowered boom. Flush sprayer out on suitable non crop area after use. DO NOT use the same spray equipment for applying other materials to plants as injury will result. DO NOT reuse empty container. Destroy by perforating and crushing. Bury or discard in a safe place.

**WARRANTY AND LIMITATION OF DAMAGES**

AMINE 4 LB is warranted to perform as described on the label. The user assumes all responsibility for the proper use of this product. The manufacturer shall not be liable for damages, including consequential damages, arising out of the use of this product. The manufacturer shall not be liable for damages, including consequential damages, arising out of the use of this product.