

DIRECTIONS FOR USE

TIME OF APPLICATION: Best results are obtained when BUTYL 6 is used on weeds that are young and in a rapidly growing condition. Application of lower rates to susceptible annual weeds usually will be satisfactory, but for perennial weeds and other conditions where kill is difficult, use higher rates. When used as a selective spray on crops, the stage of growth must be considered. Some woody plants and weeds are hard to kill and repeat applications may be necessary.

The following plants are susceptible to BUTYL-6: Pigweed, Russian Thistle, Kochia, Lambsquarter, Willows, Buckbrush, Cocklebur, Devils Claw, Sunflower.

SMALL GRAIN CROPS (Wheat, Oats, Barley, Rye): Apply BUTYL-6 at the rate of 1/3 to 2/3 pt. per acre in sufficient water for uniform coverage on small grains when fully tilled or stooled (4 to 8 inches tall) but before head emerges from the "boot". Do not use on grain undersown with legumes such as alfalfa or clovers, except where some legume injury can be tolerated. Do not apply when grain is headed. Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D.

CORN: Pre emergence, use BUTYL-6 at the rate of 1/2 to 2/3 pts. in sufficient water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting, before corn emerges. DO NOT apply to light sandy soils. Post emergence, apply at the rate of 1/3 pt. per acre. When spraying corn above 10 inches in height, use nozzle extensions ("corn drops"), directing the spray down to the base of the corn plant. DO NOT apply from tasseling to dough stage.

SORGHUM: Use BUTYL-6 at the rate of 1/3 to 2/3 pint per acre when the sorghum plants are from 6 to 12 inches high and the secondary roots are well established. Use drop nozzles when sorghum is over 10 inches high. DO NOT apply from tasseling to dough stage.

PASTURES: To control many broadleaved weeds in pastures, meadows and rangelands, use 1 1/2 to 3 quarts per acre of BUTYL-6 in sufficient water to provide for uniform application. Treat pastures when weeds are growing actively. DO NOT apply to recently seeded pastures until grass is well established. Most legumes are usually injured or killed at the rates recommended. For control of wild garlic and onion in pastures, apply 1 1/2 to 2 quarts per acre in very early spring and repeat annually until these weeds are controlled. DO NOT graze treated areas by dairy animals within seven days after treatment. Bent grass and St. Augustine grass may be damaged by BUTYL-6.

WOODY PLANT CONTROL: To control 2,4-D susceptible woody plants such as Willow, Alder, Buckbrush, Sumac, and Elderberry in non-cropped land such as roadsides, fence rows or other waste areas, use BUTYL-6, 1 to 2 qts. in 100 gallons of water. Wet thoroughly all parts of the plants, including foliage and stems, to the point of run-off.

GENERAL WEED CONTROL: Along fence rows, drainage ditch banks, around farm buildings and other waste areas, use BUTYL-6 1 1/3 to 2 pints in 100 gallons of water per acre. Thoroughly wet all foliage to point of run off.

VAPORS from this product may injure susceptible plants in the immediate vicinity.

Rev. 10-1-72 RMBN

BUTYL-6

WEED KILLER

ACCEPTED

FEB 1980

2737-40

ACTIVE INGREDIENTS:

Butyl Ester of 2,4-Dichlorophenoxyacetic acid * 73.2%
(2,4-Dichlorophenoxyacetic acid equivalent 58.5%)

INERT INGREDIENT: 26.8%

TOTAL 100%
5.6

Each gallon contains 4 lbs. 2,4-D acid.
Soluble in Oil or Water.

* Isomer Specific by AOAC Method No. 6 D01-5.

LICENSED UNDER U.S. PATENTS 2,394,916; 2,412,510;
2,390,941; 2,396,513; 2,472,347; RE 23,115

CAUTION: KEEP OUT OF THE REACH OF CHILDREN. See Side Panels For Other Cautions.

BEST DOCUMENT AVAILABLE

NET CONTENTS GALS

MANUFACTURED FOR

Pueblo Chemical & Supply Co.

611 SANTA FE DRIVE • PUEBLO, COLORADO 81006

GENERAL INFORMATION

PREPARATION OF THE SPRAY: Fill the spray tank with half the required quantity of water, then add the recommended amount of BUTYL 6 with agitation and continue filling the spray tank with water. The amount of water required in low volume applications may vary from 5 to 25 gallons per acre. For high volume applications, 50 to 100 gallons of water or more may be necessary for good coverage. In any case, use the same amount of 2,4-D recommended per acre.

CAUTION

Avoid contact with skin, eyes or clothing. Harmful if swallowed. In case of eye contact, wash with plenty of water. Avoid spray drift to susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals. Spray drift may cause severe injury to both growing and dormant plants. Coarse sprays are less likely to drift.

DO NOT contaminate irrigation ditches or water used for domestic purposes.

DO NOT use the same spray equipment for applying other materials to plants, as injury will result.

DO NOT store near fertilizers, seeds, insecticides or fungicides.

DO NOT contaminate water by cleaning equipment or disposal of wastes.

DO NOT re-use container. Destroy by rinsing with water, crush and bury in a safe place. Do not burn.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and BUYER assumes the risk of any use contrary to such directions. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing with a specific reference to this warranty. In no event shall SELLER's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

... 0737-KS-1

EPA REG NO 27740