

DIRECTIONS FOR USE

TIME OF APPLICATION: Best results are obtained when LV-4 is used on weeds that are young and in the early stages of growth. Application of LV-4 should be made on weeds usually under the following conditions: (1) on weeds in either condition when they are 3 to 6 inches high; (2) on weeds 5 to 8 inches high; (3) on weeds in the early stages of growth or on a late stage of growth; (4) on weeds that are not well established. Some weeds may require more than one application of LV-4 to be controlled.

The following plants are susceptible to LV-4:
Soybean, Corn, Sorghum, Cotton, Rice, Peanut, Sunflower, Cowpea, Bean, Cow, Turnip.

SMALL GRAIN CROPS: Weeds in the early stages of growth may be controlled by LV-4 at the rate of 1 to 2 quarts per acre. Uniform coverage should be obtained by spraying to a height of 8 inches, but before the weeds have reached a height of 12 inches. Do not use LV-4 on grain crops until the grain is headed. Do not apply LV-4 to clovers, except where some legume is present in the crop. Do not apply when grain is headed. Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D.

CORN: Pre-emergence use LV-4 at the rate of 1 to 2 quarts per acre in sufficient water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting but before grain emerges. **DO NOT** apply to light sandy soils. Post-emergence apply at the rate of 1/2 to 1 pint per acre. When spraying corn above 10 inches in height, use nozzle extension. Do not apply LV-4 to corn until the corn is headed. **DO NOT** apply LV-4 to corn until the corn is headed.

SORGHUM: LV-4 at the rate of 1 to 2 quarts per acre when the sorghum plants are 6 to 12 inches high and the secondary roots are well established. Use drop nozzles when sorghum is over 10 inches high. **DO NOT** apply LV-4 to sorghum until the grain is headed.

PASTURES: LV-4 can be applied to pastures containing weeds and large weeds. It is effective against many of the most common weeds in pastures. It is also effective against weeds when they are growing in pastures. **DO NOT** apply LV-4 to pastures until the grain is headed. **DO NOT** apply LV-4 to pastures until the grain is headed. **DO NOT** apply LV-4 to pastures until the grain is headed.

WOODY PLANT CONTROL: LV-4 is effective against many woody plants such as locoweed. Apply LV-4 to locoweed in the early stages of growth. Apply LV-4 to locoweed in the early stages of growth. Apply LV-4 to locoweed in the early stages of growth.

GENERAL WEED CONTROL: LV-4 is effective against many weeds in ditch banks, roadsides, and other areas. Apply LV-4 at the rate of 1 to 2 quarts per acre. Apply LV-4 at the rate of 1 to 2 quarts per acre.

LV-4 WEED KILLER

(Low Volatile Emulsifiable)

ACTIVE INGREDIENTS:	
2-Ethylhexyl (Isobutyl) Ester of 2,4-Dichlorophenoxyacetic Acid	65.5%
INERT INGREDIENTS:	34.5%
TOTAL	100.0%

Equivalent to 2,4 Dichlorophenoxyacetic Acid 43.4%
Contains 3.8 lbs. 2,4-D Acid Equivalent Per Gallon
*Isomer Specific by AOAC Method No. 6. Do 1-5

CAUTION:
KEEP OUT OF REACH OF CHILDREN.
HARMFUL IF SWALLOWED.
See Side Panels For Other Cautions.

NET CONTENTS GALS.

MANUFACTURED BY

Pueblo Chemical & Supply Co.

1111 GARDNER DRIVE, PUEBLO, COLORADO 81001

GENERAL INFORMATION

PREPARATION OF THE SPRAY: LV-4 is a low volatile emulsifiable concentrate. It should be diluted with water to form a spray. The dilution rate will vary with the type of weed to be controlled and the type of nozzle used. For best results, use a spray nozzle that will give a fine mist. Do not use a spray nozzle that will give a coarse spray. Do not use a spray nozzle that will give a coarse spray.

CAUTION

Do not spray LV-4 on susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals. Do not spray LV-4 on susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals.

DO NOT allow spray to drift to susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals. Spray drift may cause severe injury to both growing and dormant plants. These sprays are less likely to drift. Although this product is much less volatile than butyl or decyl esters of 2,4-D, high temperatures may produce vapors which may cause irritation to the eyes and throat.

DO NOT use LV-4 for domestic purposes.

DO NOT use the same equipment for spraying LV-4 as you use for spraying other pesticides.

DO NOT spray LV-4 on susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals.

DO NOT spray LV-4 on susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals.

WARRANTY AND LIMITATION OF DAMAGES

The purchaser of LV-4 warrants that the product is free from defects in material and workmanship at the time of sale. The purchaser warrants that the product is free from defects in material and workmanship at the time of sale. The purchaser warrants that the product is free from defects in material and workmanship at the time of sale.