

DIRECTIONS FOR USE

TIME OF APPLICATION: Best results are obtained when LV-6 is used on weeds that are young and in a fairly growing condition. Application of lower rates to susceptible broad leaf weeds usually will be satisfactory. Control of woody plants and other conditions where results differ at various locations. When used as a selective spray on crops, the following conditions should be considered. Some woody plants and weeds may be controlled and repeat applications may be necessary. The following plants are susceptible to LV-6: Pigweed, Sunflower, Thistle, Kochia, Lambquarters, Waterhemp, Barnyard, Cocklebur, Devils Claw, Sunflower.

SMALL GRAIN CROPS: (Wheat, Oats, Barley, Rye, Corn) LV-6 at the rate of 1/2 to 1 pint per acre in the early stages of uniform coverage on small grains when the plants are 4 to 8 inches tall but before most tillers appear. Do not use on grain under stress with low soil moisture. Do not use on clovers, except where clover leys are required. Do not use on alfalfa. Do not apply when grain is heading. Do not use on grain treated with granular insecticides. Do not use on grain treated with granular insecticides within 2 weeks after the application of LV-6.

CORN: Pre-emergence, use LV-6 at the rate of 1 1/2 to 2 pints in sufficient water for uniform coverage. Best results are obtained when applied 3 to 5 days after planting but before corn emerges. **DO NOT apply to light sandy soils.** Post-emergence apply at the rate of 1/2 pint per acre. When spraying corn above 10 inches in height, use nozzle combinations (corn drops), directing the spray down to the base of the corn plant. **DO NOT apply from tasseling to dough stage.**

SORGHUM: Use LV-6 at the rate of 1/2 to 1 pint per acre when the sorghum plants are from 6 to 12 inches high and the secondary tillers are well developed. Use drop nozzles when spraying above 10 inches in height. **DO NOT apply from tasseling to dough stage.**

PASTURES: Use LV-6 at the rate of 1/2 to 1 pint per acre in meadows and pastures where the weeds are in the early stages of growth. Use drop nozzles when spraying above 10 inches in height. **DO NOT apply to pastures where the weeds are in the late stages of growth.** For control of weeds and other plants in pastures, use LV-6 at the rate of 1/2 to 2 quarts per acre in very early spring or late fall. With these weeds and conditions, **DO NOT** graze the pastures by dairy animals within 30 days after application. Best control is obtained by spraying the weeds before they are grazed by LV-6.

WOODY PLANT CONTROL: Use LV-6 at the rate of 1/2 to 1 pint per acre on woody plants such as Willow, Alder, Bushy hick, Boxelder, Elderberry in open, grassy areas. Use drop nozzles when spraying above 10 inches in height. Use LV-6 at the rate of 1/2 to 1 pint per acre on water. Wet thoroughly at parts of the plant, including the stems, to the point of runoff.

GENERAL WEED CONTROL: Use LV-6 at the rate of 1/2 to 1 pint per acre on ditch banks, around farm buildings, and in other areas where weeds are growing. Use drop nozzles when spraying above 10 inches in height. Wet thoroughly at parts of the plant, including the stems, to the point of runoff.

LV-6 WEED KILLER

(Low Volatile Emulsifiable Concentrate)
Contains 5.6 lbs. 2,4-D Acid Equivalent Per Gallon

ACTIVE INGREDIENT:
Isooctyl ester of 2,4-D
Dichlorophenoxyacetic Acid* 88.8%
INERT INGREDIENTS: 11.2%
TOTAL 100.0%
*Equivalent to 2,4-D Dichlorophenoxyacetic Acid 58.9%
Isomer Specific by AOAC Method No. 6.D01-5

CAUTION:
KEEP OUT OF REACH OF CHILDREN
HARMFUL IF SWALLOWED.
See Side Panels For Other Cautions.

NET CONTENTS GALS.

MANUFACTURED BY

Pueblo Chemical & Supply Co.

1000 E. 10th St., Pueblo, Colorado 81001

GENERAL INFORMATION

PREPARATION OF THE SPRAY: Fill the spray tank with half the recommended quantity of water. Then add the recommended amount of LV-6 with agitation and continue filling the spray tank with water. The amount of water required in low volume applications may vary from 5 to 25 gallons per acre. For high volume applications, 50 to 100 gallons of water or more may be required. For best coverage, in any case, use the same amount of 2,4-D as the recommended product.

CAUTION

Do not spray when wind is blowing. Control of weeds is best when plants are in contact with plenty of water. Do not spray on plants such as cotton, tomatoes, and ornamentals. Spray drift may be harmful to both growing and dormant plants. Coarse sprays are less likely to drift. Although this ester is much less volatile than butyl or isopropyl esters of 2,4-D, high temperatures may produce vapors after application which might injure sensitive plants growing nearby. Contamination of shallow standing waters at application rates recommended in this label may kill fish.

DO NOT contaminate irrigation ditches or waterways for domestic purposes.

DO NOT use the same spray equipment to apply to other materials to plants as in this label.

DO NOT use on plants listed in the Restricted Use section of this label.

DO NOT use on plants listed in the Restricted Use section of this label.

DO NOT use on plants listed in the Restricted Use section of this label.

NON WARRANTY

PUEBLO CHEMICAL & SUPPLY CO. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE. EXCEPT FOR THE LIMITED WARRANTY SET FORTH IN THIS LABEL, THE USER ASSUMES ALL LIABILITY FOR THE USE OF THIS PRODUCT.

U.S. PAT. NO. 2,447,442
U.S. PAT. NO. 2,757,853