

MANEB 80 WP

FUNGICIDE

ACCEPTED

DEC 8 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT CIDE ACT
FOR ECONOMIC POISON REGISTERED
EPA UNDER NO. 2749-22 SUBJECT
TO ATTACHED COMMENTS.

ACTIVE INGREDIENT:

Maneb (Manganese ethylenebisdithiocarbamate) 80%

INERT INGREDIENTS:

20%
TOTAL 100%

EPA REG. NO. 2749-22

NET WEIGHT _____

CAUTION: KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS.

May cause irritation of eyes, nose, throat and skin. May be harmful if inhaled or swallowed.

Do not breathe spray mist. Use with adequate ventilation. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water. For eyes - get medical attention.

This product is toxic to fish. Keep out of lakes, streams ponds and other bodies of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not let MANEB 80 WP become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness and create vapors which may be flammable. Keep MANEB 80WP away from fires and sparks. Keep container closed when not in use. Do not reuse empty container. Destroy empty container by perforating or crushing and burying in a safe place.

DIRECTIONS FOR USE:

SPRAYS - Use 1 1/2 lbs. of Aceto MANEB 80 WP per 100 gals. of spray, unless otherwise stated. If foliage is difficult to wet, add 4 fl. ozs. of a spreader-sticker to each 100 gals. of spray. During severe disease outbreaks or in high humidity conditions favor disease, use 2 lbs. per 100 gals. and shorten interval between applications.

Applications for specific crops:

CROPS
ALMONDS

SPRAY

1 1/2 to 2 lbs. per 100 gals. 150 gals. per acre

CABBAGES

150 gals. per acre

LETTUCE & ENDIVE

LIMA BEANS

200 gals. per acre

POTATOES

100 gals. per acre

SUGAR BEETS

1 1/2 to 2 lbs. per 75-125 gals. of water

SWEET CORN

3 lbs. per 100 gals. of water

A.C. ...
Dec - 8 - 1972
2749-22

ALMONDS (CALIFORNIA): Shot-hole (Fungus) and brown rot blossom and twig blight - Apply at the popcorn, full bloom and post-fall stages.

CABBAGE: Alternaria leaf spots and downy mildew - Apply 7 to 10 days after planting. If disease is present, apply earlier. Repeat every 3 days until plants are set in the field or until field-seeded plants are thinned. Do not apply within 7 days of picking.

CANTALOUPE AND SQUASH (TEXAS): Downy Mildew - Apply when disease threatens; repeat as needed every 7 to 10 days. Do not apply within 5 days of harvest.

CARROTS: Leaf spot diseases - Apply when plants are 6 weeks old and repeat every 7 to 10 days.

CELERY: Early blight (Cercospora) and late blight (Septoria) - Apply when plants are established in the field. Repeat every 7 days. Do not apply within 14 days of harvest.

If there are severe disease conditions, repeat every 3 to 5 days. Remove residues by stripping, trimming and washing.

CUCUMBERS: Downy Mildew, Anthracnose - Apply when disease threatens, repeat every 7 to 10 days. Do not apply within 5 days of harvest.

GRAPES: (In Great Lakes area): Black rot - Apply just before bloom, repeat after bloom and 7 to 10 days later.

LETTUCE AND ENDIVE: Downy Mildew - Apply when disease is present. Repeat every 7 to 10 days. Do not apply within 7 days of harvest.

Remove residues from leaf lettuce by stripping and washing. Remove residues from leaf lettuce and endive by thorough washing.

LIMA BEANS: Downy Mildew and Anthracnose - Apply when disease first appears. Repeat every 7 to 10 days. Do not apply within 7 days of harvest.

POTATOES: Late blight - Apply when disease first appears. Repeat every 7 to 10 days. Do not apply within 7 days of harvest.

SUGAR BEETS: Downy Mildew - Apply when disease first appears. Repeat every 7 to 10 days. Do not apply within 7 days of harvest.

Dec-3-1973

2744-35

SPINACIA: Downy mildew - Apply when plants are 2 to 6 inches high. Repeat every 4 to 7 days for 3 weeks if disease threat.

SPINACIA: Downy mildew and white rust - Apply during first 10 days of growth. Repeat every 4 to 7 days. Do not apply within 10 days of harvest. Remove residue by thorough washing.

SPINACIA: Downy mildew - Apply when plants are 2 to 6 inches high. Repeat every 4 to 7 days for 3 weeks if disease threat. Do not apply within 10 days of harvest. Remove residue by thorough washing.

SPINACIA: Downy mildew - Apply when plants are 2 to 6 inches high. Repeat every 4 to 7 days for 3 weeks if disease threat. Do not apply within 10 days of harvest. Remove residue by thorough washing.

SPINACIA: Downy mildew - Apply when plants are 2 to 6 inches high. Repeat every 4 to 7 days for 3 weeks if disease threat. Do not apply within 10 days of harvest. Remove residue by thorough washing. Do not use on tender young seedlings in greenhouse as injury may occur.

Camerman's Note
Poor Copy

MANUFACTURED FOR: ACETO CHEMICAL CO., INC.
Agricultural Chemicals Division
118-02 Northern Boulevard
Flushing, New York 11368