

1084  
2749-22

BEST AVAILABLE COPY

# MANEB 80-W FUNGICIDE

<b>ACTIVE INGREDIENTS:</b>	
Maneb (Manganese Ethylenebisdithiocarbamate) (Metallic Manganese Equivalent 16.5%)	80%
<b>INERT INGREDIENTS:</b>	20%
<b>TOTAL INGREDIENTS</b>	<b>100%</b>

ACCEPTED  
 NOV 29 1984  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 2749-22

KEEP OUT OF REACH OF CHILDREN  
CAUTION

### STATEMENT OF PRACTICAL TREATMENT

- If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.
- If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.
- If in Eyes: Flush with plenty of water for 15 minutes. Get medical attention.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONS

NET CONTENTS: 50 LBS.

Manufactured For: **Aceto Agricultural Chemicals Corp.**  
 126-02 Northern Boulevard  
 Flushing, New York 11368

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Wash thoroughly after handling. Avoid contact with skin, eyes or clothing. Avoid breathing dust ON SPRAY MIST.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not apply where runoff is like to occur and when weather conditions favor drift from areas treated. Do not contaminate water <sup>BY</sup> cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is a reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers.

Written or oral warnings must include the following information:

CAUTION

(insert area or field description) treated with *M...*  
on (insert date of application.

Do not enter treated areas without protective clothing until sprays have been dried.

Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

BEST AVAILABLE COPY

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

**STORAGE:** Store in a safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang can affect package strength.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide and Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**DIRECTIONS FOR**

**SPRAYS:** Use 1X lbs. of Aceto Meneb 50W per acre. If wet, add 4 fl. ozs. of a spreader-sticker to each 100 gallons of water. For control of late leaf diseases, use 2 lbs. per 100 gals. of spray solution.

**DUSTS:** Aceto Meneb 50W may be used at other dilutions. For control of late leaf diseases, mix with a neutral diluent. Use dust mixtures as soon as possible.

Applications for special crops:

- CROPS
- ALMONDS
- CABBAGES
- LETTUCE & ENDIVE
- LIMA BEANS
- POTATOES
- SUGAR BEETS

**BEST AVAILABLE COPY**

**SWEET CORN**

**ALMONDS (CALIFORNIA)** - All bloom and post-fall crop

**CABBAGE:** Alternating leaf; weevil damage; early blight. Repeat every 7 days; full plants 7 days of picking.

**CANTALOUPE AND SQUASH (TEXAS):** Do not apply to 10 days.

**CARROTS:** Leaf spot diseases - Apply when plants are young.

**CELERY:** early blight (Cercospora) and late blight. Repeat every 7 days.

If there are several disease conditions, repeat every 7 days.

**CUCUMBERS:** Downy mildew, anthracnose. Apply when plants are young. Repeat every 7 days.

**GRAPES:** (In Great Lakes Area): Black rot. Apply just before harvest. Repeat 7 or 10 days later.

**LETTUCE AND ENDIVE:** Downy mildew. Apply when plants are young. Repeat every 7 days. Do not apply within 7 days of picking. Remove residues from leaf lettuce and endive by thorough washing.

**LIMA BEANS:** Downy mildew and anthracnose. Apply when plants are young. Repeat every 7 days. Do not apply within 4 days before picking.

**ONIONS:** Downy mildew, purple blotch (Blight). Repeat every 7 days as needed.

**POTATOES:** Early and late blights. Apply when plants are young. Repeat every 7 days as blights threaten.

**SPINACH:** Downy mildew (Blue Mold) and white rust. Do not apply within 7 days of picking. Remove residues by thorough washing.

**SUGAR BEETS:** Cercospora leaf spot. Apply when plants are young. Repeat every 7 days. Do not apply within 30 days of picking. Do not treat treated crops.

**SWEET CORN:** Downy mildew, anthracnose.

