

9-26-72  
2744-36



# AMAZIN

## Zinc Enriched Maneb 80W Fungicide

# WITH 2% ZINC

### ACTIVE INGREDIENT:

Maneb (Manganese Ethylenebisdithiocarbamate) 80%  
(Metallic Manganese Equivalent) 10.5

### INERT INGREDIENTS

TOTAL INGREDIENTS 100%

Net weight  
50 LBS.

EPA REG. NO. 2749-35

## CAUTION:

KEEP OUT OF THE REACH OF CHILDREN.  
DO NOT APPLY TO OPEN WOUNDS OR IRRITATED SKIN.  
MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN.  
MAY BE HARMFUL IF INHALED OR SWALLOWED.

**THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS, PONDS AND OTHER BODIES OF WATER. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.**

**APPLICATIONS FOR SPECIFIC CROPS:**  
**APPLES** - Scab, Cedar apple rust, Black rot, Fly speck, Sooty blotch, Bitter rot, White rot - Use 1 to 2 lbs. per 100 gals. or up to 10 1/2 lbs per acre in concentrated sprays. Apply in pre-pink and repeat in prebloom, bloom and cover sprays. Follow state spray schedules.  
**APPLICABLE MANEBS** - Use 1 1/2 lbs. of Amazin Maneb 80W per 100 gals. of spray, unless otherwise stated. If foliage is difficult to wet, add 4 FL. ozs. of a spreader-sticker to each 100 gals. of spray. During severe disease outbreaks or if weather conditions favor disease use 2 lbs. per 100 gals. of spray and shorten interval between application.  
**DUSTS** - Amazin Maneb 80W may be used at concentrations of 6 to 8% (4.8 to 6.4% active) by diluting and thoroughly mixing with a neutral diluent. Use dust mixtures as soon as possible after preparation.

**CONTROLS DISEASES ON THESE CROPS:**

FIELD CROPS	FRUITS & NUTS	VEGETABLES	VEGETABLES (cont'd)
Sugar Beets	Apples, Grapes	Broccoli, Cucumbers	Onions, Sweet Corn
Tobacco	Apricots, Nectarines	Cabbage, Endive	Peppers, Squash
	Cranberries, Peaches	Cantaloupe, Lettuce	Potatoes, Tomatoes
		Carrots, Lima Beans	Pumpkins, Watermelon
		Celery	Spinach

parting. If disease is present, apply earlier. Repeat every 7 days until plants are out of the field or until field seeded parts are finished. Do not apply within 7 days of harvest.  
**CANTALOUPE AND SQUASH (TEXAS, DOW)** - Mildew - Apply when disease threatens. Repeat as needed every 7 to 10 days.  
**CARROTS** - Leaf spot diseases - apply when plants are 6 weeks old and repeat every 7 to 10 days.  
**CELERY** - Early blight (Cercospora) and late blight (Septoria) - Apply when plants are established in the field. Repeat every 7 days. If there are severe disease conditions, repeat every 3 to 5 days. Remove residues by stripping, trimming and washing. **DO NOT APPLY WITHIN 14 DAYS OF HARVEST.**

**CRANBERRIES (MASSACHUSETTS, NEW JERSEY AND WISCONSIN)** - Fruit rots - Use three to six pounds per acre. Apply at mid bloom and repeat per state spray schedules. Do not apply within 30 days of harvest.

**CUCUMBERS** - Downy Mildew, anthracnose - Apply when disease threatens. Repeat every 7 to 10 days. Do not apply more than 3 lbs. per acre per application. **DO NOT APPLY WITHIN 5 DAYS OF HARVEST.**

**GRAPES (IN GREAT LAKES AREA)** - Black rot - apply just before bloom, just after bloom and 7 to 10 days later.

**LETTUCE & ENDIVE** - Downy Mildew - As a spray use up to 150 gals. per acre. As a dust use up to 30 lbs. per acre. Apply when disease is present. Repeat every 7 to 10 days. Do not apply within 7 days of picking. Remove residues from lettuce by stripping and trimming. Remove residues from leaf lettuce and endive by thorough washing. **DO NOT APPLY WITHIN 10 DAYS OF HARVEST.**

**LIMA BEANS** - Downy Mildew and anthracnose - As a spray use up to 200 gals. per acre. As a dust use 40 lbs. per acre. Apply when disease first appears. Repeat every 4 to 7 days as needed. Do not apply within 4 days before picking.

**ONIONS** - Downy Mildew, purple blotch (Blight) and blight (Botrytis leaf blight) - apply when first spots appear. Repeat every 7 days as needed.

**PEACHES** (Pacific States) and **NECTARINES** (California) - Shot hole (fungus) and peach leaf curl (California only) apply dormant and popcorn sprays. For brown rot blossom blight apply at pink bud and 25% to 75% full bloom (Add a petal fall spray in Pacific Northwest). For fruit brown rot (California only) apply 2 lbs. per 100 gals. Four weeks before harvest if rot is found on green fruit apply earlier and repeat at 7 to 14 days intervals as needed. On peaches if applied within 14 days of harvest remove residue by brushing. Do not apply later than 2 days before harvest. On Nectarines do not apply later than 14 days before harvest.

**PEPPERS** - Anthracnose, Ripe rot, Cercospora leaf spot (Frog eye spot) - Use 2 lbs. per 100 gals. For Cercospora leaf spot apply when disease threatens. Repeat as needed at 7 to 10 days intervals. For Anthracnose and ripe rot apply when the first fruit spots start to form. Repeat as long as disease threatens at 7 to 10 days intervals. Do not apply more than 3.4 lbs. per acre per application.

**POTATOES** - Early and late blights - Use 1 to 2 lbs. per acre. Apply when plants are 2 to 6 inches high. Repeat every 5 to 10 days for as long as blight threatens.

**PUMPKINS** - Downy Mildew, Angular leaf spot - Use 3 lbs. per acre in a sufficient amount of water to wet plants thoroughly. Apply when disease threatens. Repeat at 7 to 10 days intervals as needed.

**ROSES** - Cercospora leaf spot (TEXAS) Rust (California), Black spot - Use 1 1/2 lbs. Spray when first leaves unfold. Repeat every 7 days as needed.

**SPINACH** - Downy Mildew (Blue Mold) and white rust - Apply during first sign of disease. Repeat every 4 to 7 days. Do not apply within 7 days of picking. Remove by thorough washing. **DO NOT APPLY WITHIN 14 DAYS OF HARVEST.**

**SUGAR BEETS** - Cercospora leaf spot - As a spray use 1 1/2 to 2 lbs. per acre in 75 to 125 gals. of water. As a dust use 30 to 40 lbs. per acre. Apply when disease threatens. Repeat every 7 to 10 days as needed. Do not apply within 30 days of picking.

**SWEET CORN** - Helminthosporium maydis leaf blight - Use 2 lbs. per acre. Apply when disease threatens. Repeat every 4 to 7 days as needed. Do not feed to livestock.

**TOMATOES** - Blue Mold, Anthracnose - Use 1/2 lbs. per 100 gals. or 1 1/2 to 5 gals. of diluted suspension per 100 sq. yards. Spray plant beds when plants are the size of a dime or when disease is reported in the area. Repeat every 3 days or more if worked with rain. Continue until transplanting is complete.

**TOMATOES** - Early blight, late blight, Septoria leaf spot, gray leaf spot (Stemphylium) and

# WITH 2% ZINC

**ACTIVE INGREDIENT:**

Maneb (Manganese Ethylenebisdithiocarbamate) 80%  
(Metallic Manganese Equivalent 15%)

**INERT INGREDIENTS**

**TOTAL INGREDIENTS**

Net weight  
**50 LBS.**

EPA REG. NO. 2749-35

**CAUTION:**

KEEP OUT OF THE REACH OF CHILDREN.  
DO NOT FEED TO ANIMALS.  
MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN.  
MAY BE HARMFUL IF INHALED OR SWALLOWED.

Avoid contact with skin. Wash thoroughly with plenty of water. For eyes, flush with water.

**THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS, PONDS AND OTHER BODIES OF WATER. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.**

Store in a cool, dry place. Avoid contact with water. This may lead to certain diseases. Keep container closed when not in use. Do not use in windy conditions. Do not apply to plants by pouring or drenching.

**DIRECTIONS FOR USE:**

**SPRAYS** - Use 1 1/2 lbs. of Amazin Maneb 80W per 100 gals. of spray, unless otherwise stated. If foliage is difficult to wet, add 4 FL. ozs. of a spreader-sticker to each 100 gals. of spray. During severe disease outbreaks or if weather conditions favor disease, use 2 lbs. per 100 gals. of spray and shorten interval between application.

**DUSTS** - Amazin Maneb 80W may be used at concentrations of 6 to 8% (4.8 to 6.4% active) by diluting and thoroughly mixing with a neutral diluent. Use dust mixtures as soon as possible after preparation.

**CONTROLS DISEASES ON THESE CROPS:**

FIELD CROPS	FRUITS & NUTS	VEGETABLES	VEGETABLES (cont'd)
Sugar Beets	Apples, Grapes	Broccoli, Cucumbers	Onions, Sweet Corn
Tobacco	Apricots, Nectarines	Cabbage, Endive	Peppers, Squash
	Cranberries, Peaches	Cantaloupe, Lettuce	Potatoes, Tomatoes
		Carrots, Lima Beans	Pumpkins, Watermelon
		Celery	Spinach

**APPLICATIONS FOR SPECIFIC CROPS:**

**APPLES** - Scab, Cedar apple rust, Black rot, Fly speck, Spotty blotch, Bitter rot, White rot - Use 1 to 2 lbs. per 100 gals. or up to 10 1/2 lbs. per acre in concentrated sprays. Apply in pre pink and repeat in prebloom, bloom and cover sprays. Follow state spray schedules.

**APRICOTS (CALIFORNIA)** - Brown rot and shot-hole (fungus) - Use 2 lbs. at red bud, early bloom and fullbloom. On Green rot or Jacket rot - use 2 lbs. at full bloom and petal fall.

**BROCCOLI** - Downy Mildew - Apply 4 lbs. per acre. As a spray use 100 to 150 gals. per acre. As an 8% dust use 50 lbs. per acre. Apply when disease threatens and continue as needed at 7 to 14 days intervals. Shorten intervals when disease conditions are severe. Repeat after rains. If applied less than 3 days before harvest, remove residues by trimming and washing.

**CABBAGE** - Alternaria leaf spots and Downy Mildew - As a spray use 1 1/2 lbs. per 100 gals. and apply up to 150 gals. per acre. As a dust use up to 40 lbs. per acre. Apply 7 to 10 days after

six pounds per acre. Apply at mid bloom and repeat per state spray schedules. Do not apply within 30 days of harvest.

**CUCUMBERS** - Downy Mildew, Anthracnose - Apply when disease threatens. Repeat every 7 to 10 days. Do not apply within 30 days of harvest. Do not apply within 5 days of harvest.

**GRAPES** - IN GREAT BASINS AND ALL BASINS - Apply just before bloom and repeat every 7 to 10 days after.

**LETTUCE & ENDIVE** - Downy Mildew - As a spray use up to 150 gals. per acre. As a dust use up to 50 lbs. per acre. Apply when disease is present. Repeat every 7 to 10 days. Do not apply within 7 days of picking. Remove residues from head lettuce by stripping and trimming. Remove residues from leaf lettuce and endive by thorough washing. Do not apply within 10 days of harvest.

**LIMA BEANS** - Downy Mildew and anthracnose - As a spray use up to 200 gals. per acre. As a dust use 40 lbs. per acre. Apply when disease first appears. Repeat every 4 to 7 days as needed. Do not apply within 4 days before picking.

**ONIONS** - Downy Mildew, purple blotch (Blight) and blast (Botrytis leaf blight) - apply when first spots appear. Repeat every 7 days as needed.

**PEACHES** (Pacific States) and **NECTARINES** (California) - Shot-hole (fungus) and peach leaf curl (California only) apply dormant and popcorn sprays. For brown rot blossom blight apply at pink bud and 25% to 75% full bloom (Add a petal fall spray in Pacific Northwest). For fruit brown rot (California only) apply 2 lbs. per 100 gals. Four weeks before harvest if rot is found on green fruit apply earlier and repeat at 7 to 14 days intervals as needed. On peaches if applied within 14 days of harvest remove residue by brushing. Do not apply later than 2 days before harvest. On Nectarines do not apply later than 14 days before harvest.

**PEPPERS** - Anthracnose, Ripe rot, Cercospora leaf spot (Frog-eye spot) - use 2 lbs. per 100 gals. For Cercospora leaf spot apply when disease threatens. Repeat every 7 to 10 days intervals. For Anthracnose and ripe rot apply when the first fruits begin to turn red. Repeat as long as disease threatens at 7 to 10 days intervals. Do not apply more than 3 1/2 lbs. per acre per application.

**POTATOES** - Early and late blights - Use 1 to 2 lbs. per acre. Apply when plants are 2 to 6 inches high. Repeat every 5 to 10 days for as long as blight threatens.

**PUMPKINS** - Downy Mildew, Angular leaf spot - use 3 lbs. per acre in a sufficient amount of water to wet plants thoroughly. Apply when disease threatens. Repeat at 7 to 10 days intervals as needed.

**ROSES** - Cercospora leaf spot (TEXAS), Rust (California), Black spot - Use 1 1/2 lbs. Spray when first leaves unfold; repeat every 7 days as needed.

**ST. NACH** - Downy Mildew (Blue Mold) and white rust - Apply during first sign of disease. Repeat every 4 to 7 days. Do not apply within 7 days of picking. Remove by thorough washing. Do not apply within 10 days of harvest.

**SUGAR BEETS** - Cercospora leaf spot - As a spray use 1 1/2 to 2 lbs. per acre in 75 to 125 gals. of water. As a dust use 30 to 40 lbs. per acre. Apply when disease threatens. Repeat every 7 to 10 days as needed. Do not apply within 30 days of picking.

**SWEET CORN** - Helminthosporium maydis leaf blight - Use 2 lbs. per acre. Apply when disease threatens. Repeat every 4 to 7 days as needed. Do not feed treated forage to livestock.

**TOBACCO** - Blue Mold, Anthracnose - use 1/2 lb. per 100 gals. Use 2 1/2 to 5 gals. of diluted suspension per 100 sq. yards. Spray plant beds when plants are the size of a dime or when disease is reported in the area. Repeat every 3 days or more often if worked with rain. Continue until transplanting is complete.

**TOMATOES** - Early blight, late blight, Septoria leaf spot, gray leaf spot (Stemphylium) and anthracnose - Apply when fruit of first clusters is well formed. Repeat every 7 to 10 days. When weather conditions favor disease, more frequent applications may be necessary. When anthracnose is a serious problem on canning tomatoes use 3 lbs. per acre. Do not apply more than 3 1/2 lbs. per acre per application. Do not use on tender young seedlings in greenhouses as injury may occur. Do not apply within 5 days of harvest.

**WATERMELONS** - Downy Mildew, Anthracnose, Cercospora leaf spot, gummy stem blight - use 1 1/2 to 2 lbs. per 100 gals. of water. Apply as a full coverage spray up to 250 gals. per acre. Apply when runners form or when disease threatens. Repeat as needed at 7 to 10 days intervals. Under severe conditions shorten intervals to 3 to 5 days. Do not apply within 5 days of harvest.

Manufactured for: ACETO CHEMICAL CO., INC., Agricultural Chemicals Division, 126-02 Northern Boulevard, Flushing, New York 11368

Camerman's Note  
Poor Copy

# Zinc Linched Maneb 80W Fungicide

# WITH 2% ZINC

### ACTIVE INGREDIENT:

Maneb (Manganese Ethylenebisdithiocarbamate) 80%  
(Metallic Manganese Equivalent) 5

### INERT INGREDIENTS

20  
TOTAL INGREDIENTS 100%

Net weight  
50 LBS.

EPA REG. NO. 2749-36

## CAUTION:

KEEP OUT OF THE REACH OF CHILDREN.  
DO NOT FEED TO ANIMALS AND DO NOT  
MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN.  
MAY BE HARMFUL IF INHALED OR SWALLOWED.

Keep the dust or spray mixture with adequate ventilation. Avoid contact with skin and eyes. Wash thoroughly with plenty of water. For eye contact, get medical attention.

**THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS, PONDS AND OTHER BODIES OF WATER. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.**

Store in a cool, dry place. Do not store in a place where children can reach. This may lead to certain diseases. Do not store in a place where children can reach. Do not store in a place where children can reach. Do not store in a place where children can reach. Do not store in a place where children can reach. Do not store in a place where children can reach.

### DIRECTIONS FOR USE

**SPRAYS** - Use 1 1/2 lbs. of Amazin Maneb 80W per 100 gals. of spray, unless otherwise stated. If foliage is difficult to wet, add 4 FL. ozs. of a spreader sticker to each 100 gals. of spray. During severe disease outbreaks or if weather conditions favor disease, use 2 lbs. per 100 gals. of spray and shorten interval between application.

**DUSTS** - Amazin Maneb 80W may be used at concentrations of 6 to 8 % (4.8 to 6.4 % active) by diluting and thoroughly mixing with a neutral diluent. Use dust mixtures as soon as possible after preparation.

### CONTROLS DISEASES ON THESE CROPS:

FIELD CROPS	FRUITS & NUTS	VEGETABLES	VEGETABLES (cont'd)
Sugar Beets	Apples Grapes	Broccoli Cucumbers	Onions Sweet Corn
Tobacco	Apricots Nectarines	Cabbage Endive	Peppers Squash
	Cranberries Peaches	Cantaloupe Lettuce	Potatoes Tomatoes
		Carrots Lima Beans	Pumpkins Watermelon
		Celery	Spinach

### APPLICATIONS FOR SPECIFIC CROPS:

**APPLES** - Scab, Cedar apple rust, Black rot, Fly speck, Spotty blotch, Bitter rot, White rot - Use 1 to 2 lbs. per 100 gals. or up to 10 1/2 lbs. per acre in concentrated sprays. Apply in pre-bloom and repeat in prebloom, bloom and cover sprays. Follow state spray schedules.

**APRICOTS (CALIFORNIA)** - Brown rot and shot-hole (fungus) - Use 2 lbs. at red bud, early bloom and fullbloom. On Green rot or Jacket rot - use 2 lbs. at full bloom and petal fall.

**BROCCOLI** - Downy Mildew - Apply 4 lbs. per acre. As a spray use 100 to 150 gals. per acre. As a dust use 50 lbs. per acre. Apply when disease threatens and continue as needed at 7 to 14 days intervals. Shorten intervals when disease conditions are severe. Repeat after rains. If applied less than 3 days before harvest, remove residues by trimming and washing.

**CABBAGE** - Alternaria leaf spots and Downy Mildew - As a spray use 1 1/2 lbs. per 100 gals. and apply up to 150 gals. per acre. As a dust use up to 40 lbs. per acre. Apply 7 to 10 days after

ONION BULBS - MASSACHUSETTS NEW JERSEY AND WISCONSIN - Use 10 to 15 pounds per acre. Apply at mid bloom and repeat per state spray schedules. Do not apply within 30 days of harvest.

**CUCUMBERS** - Downy Mildew and anthracnose - Apply when disease threatens, repeat every 7 to 10 days. Do not apply more than 3 1/2 lbs. per acre per application. **DO NOT APPLY WITHIN 5 DAYS OF HARVEST**

**GRAPE (IN GREAT AREAS)** - Downy Mildew - Apply at mid bloom and repeat every 7 to 10 days after.

**LETTUCE & ENDIVE** - Downy Mildew - As a spray use up to 100 gals. per acre. As a dust use up to 50 lbs. per acre. Apply when disease threatens. Repeat every 7 to 10 days. Do not apply within 7 days of picking. Remove residues from heads by trimming and washing. Remove residues from leaf lettuce and endive by thorough washing. **DO NOT APPLY WITHIN 10 DAYS OF HARVEST**

**LIMA BEANS** - Downy Mildew and anthracnose - As a spray use up to 200 gals. per acre. As a dust use 40 lbs. per acre. Apply when disease first appears. Repeat every 4 to 7 days as needed. Do not apply within 4 days before picking.

**ONIONS** - Downy Mildew, purple blotch (Blight) and blast (Botrytis leaf blight) - apply when first spots appear. Repeat every 7 days as needed.

**PEACHES** (Pacific States) and **NECTARINES** (California) - Shot hole (fungus) and peach leaf curl (California only) apply dormant and popcorn sprays. For brown rot blossom blight apply at pink bud and 25% to 75% full bloom (Add a petal fall spray in Pacific Northwest). For fruit brown rot (California only) apply 2 lbs. per 100 gals. Four weeks before harvest if rot is found on green fruit apply earlier and repeat at 7 to 14 days intervals as needed. On peaches if applied within 14 days of harvest remove residue by brushing. Do not apply later than 2 days before harvest. On Nectarines do not apply later than 14 days before harvest.

**PEPPERS** - Anthracnose, Ripe rot, Cercospora leaf spot (Frog eye spot) - use 2 lbs. per 100 gals. For Cercospora leaf spot apply when disease threatens. Repeat as needed at 7 to 10 days intervals. For Anthracnose and ripe rot apply when the first fruit begins to turn red. Repeat as long as disease threatens at 7 to 10 days intervals. Do not apply more than 3 1/2 lbs. per acre per application.

**POTATOES** - Early and late blights - Use 1 to 2 lbs. per acre. Apply when plants are 2 to 6 inches high, repeat every 5 to 10 days for as long as blight threatens.

**PUMPKINS** - Downy Mildew, Angular leaf spot - use 3 lbs. per acre in a sufficient amount of water to wet plants thoroughly. Apply when disease threatens; repeat at 7 to 10 days intervals as needed.

**ROSES** - Cercospora leaf spot (TEXAS), Rust (California) Black spot - Use 1 1/2 lbs. Spray when first leaves unfold, repeat every 7 days as needed.

**SPINACH** - Downy Mildew (Blue Mold) and white rust - Apply during first sign of disease. Repeat every 4 to 7 days. Do not apply within 7 days of picking. Remove by thorough washing. **DO NOT APPLY WITHIN 5 DAYS OF HARVEST**

**SUGAR BEETS** - Cercospora leaf spot - As a spray use 1 1/2 to 2 lbs. per acre in 75 to 125 gals. of water. As a dust use 30 to 40 lbs. per acre. Apply when disease threatens. Repeat every 7 to 10 days as needed. Do not apply within 30 days of picking.

**SWEET CORN** - Helminthosporium maydis leaf blight - Use 2 lbs. per acre. Apply when disease threatens. Repeat every 4 to 7 days as needed. Do not feed treated forage to livestock.

**TOBACCO** - Blue Mold, Anthracnose - use 1/2 lbs. per 100 gals. Use 2 1/2 to 5 gals. of diluted suspension per 100 sq. yards. Spray plant beds when plants are the size of a dime or when disease is reported in the area. Repeat every 3 days or more often if worked with rain. Continue until transplant is complete.

**TOMATOES** - Early blight, late blight, Septoria leaf spot, gray leaf spot (Stemphylium) and anthracnose - Apply when fruit of first clusters is well formed. Repeat every 7 to 10 days. When weather conditions favor disease, more frequent applications may be necessary. When anthracnose is a serious problem on canning tomatoes use 3 lbs. per acre. Do not apply more than 3 1/2 lbs. per acre per application. Do not use on tender young seedlings in greenhouses as injury may occur. **DO NOT APPLY WITHIN 5 DAYS OF HARVEST**

**WATERMELONS** - Downy Mildew, Anthracnose, Cercospora leaf spot, gummy stem blight - use 1 1/2 to 2 lbs. per 100 gals. of water. Apply as a full coverage spray up to 250 gals. per acre. Apply when runners form or when disease threatens. Repeat as needed at 7 to 10 days intervals. Under severe conditions shorten intervals to 3 to 5 days. **DO NOT APPLY WITHIN 5 DAYS OF HARVEST**

Manufactured for: ACETO CHEMICAL CO., INC., Agricultural Chemicals Division, 126-02 Northern Boulevard, Flushing, New York 11368

Cameraman's Note  
Poor Copy