

DDVP TECHNICAL INSECTICIDE

ACTIVE INGREDIENT:

2,2-Dichlorovinyl dimethylphosphate


INERT INGREDIENTS:

98%
2%
100%

TOTAL

EPA REG. NO. 2749-38
NET WEIGHT

DANGER



2749-38

DANGER: KEEP OUT OF REACH OF CHILDREN

ANTIDOTE AND FIRST AID TREATMENT

ANTIDOTE IS ATROPINE SULFATE 2 PAM IS ALSO ANTIDOTAL AND MAY BE USED IN CONJUNCTION WITH ATROPINE.

INTERNAL: IF THE MATERIAL HAS BEEN SWALLOWED, INDUCE VOMITING IMMEDIATELY. THIS MAY BE DONE BY INTRODUCING A FINGER INTO THE THROAT OR BY GIVING WARM SALT WATER (1 TABLESPOON OF SALT TO A GLASS OF WATER). NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

EXTERNAL: IF THE MATERIAL HAS BEEN SPILLED ON THE SKIN, IMMEDIATELY REMOVE PATIENT FROM THE VICINITY OF THE INSECTICIDE, REMOVE ALL CONTAMINATED CLOTHING AND WASH SKIN WITH SOAP AND RUNNING WATER. IF THE MATERIAL GETS INTO THE EYES, WASH IMMEDIATELY WITH RUNNING WATER FOR AT LEAST TEN MINUTES.

CALL A PHYSICIAN IMMEDIATELY.

POISONOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN. RAPIDLY ABSORBED THROUGH SKIN.

DO NOT SWALLOW OR GET IN EYES, ON SKIN OR ON CLOTHING.

WEAR CLEAN RUBBER GLOVES, CLEAN WATERPROOF PROTECTIVE CLOTHING AND GOGGLES. REPLACE GLOVES FREQUENTLY AND DESTROY USED GLOVES. WEAR A MASK OR RESPIRATOR OF A TYPE PASSED BY THE U. S. DEPARTMENT OF MINES FOR PROTECTION AGAINST THIS MATERIAL. WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING AND BEFORE EATING OR SMOKING. IF THE CHEMICAL GETS INTO THE EYES, IMMEDIATELY FLUSH WITH WATER FOR AT LEAST 15 MINUTES AND GET MEDICAL ATTENTION. IN CASE OF CONTACT REMOVE ALL CONTAMINATED CLOTHING AND WASH SKIN WITH SOAP AND WATER. WASH CLOTHING BEFORE REUSE. IN CASE OF SPILLAGE, WASH DOWN WITH LARGE AMOUNTS OF WATER.

IF WARNING SYMPTOMS APPEAR:

KEEP PATIENT PRONE AND QUIET.

START ARTIFICIAL RESPIRATION IMMEDIATELY IF PATIENT IS NOT BREATHING. TRANSPORT THE PATIENT IMMEDIATELY TO THE NEAREST PHYSICIAN.

NOTE TO PHYSICIAN

WARNING SYMPTOMS: Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps

TREATMENT. Atropine is the specific therapeutic antagonist of choice against the parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10-minute intervals in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. Clear chest by postural drainage. Oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

DO NOT STORE NEAR FEED OR FOODSTUFFS.

This product is toxic to fish, birds and other wildlife. Do not contaminate water by cleaning, or disposal of wastes.

Do not use empty container. Do not reuse container by perforating, burning, and reusing in any place.

DIRECTIONS FOR USE: For use by registered trade Formulators for manufacturing purposes only. Not for resale under this label.



Manufactured by
ACE TO CHEMICAL CO. INC.
AGRI-CHEMICAL CHEMICAL DIVISION
12602 NORTHERN BOULEVARD, FLEMING, N. Y. 11734