


U.S. PATENT OFFICE
WASHINGTON, D.C. 20540

[Faint, illegible text, likely bleed-through from the reverse side of the page]



ACCEPTED
FEB 1 1974
 UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT
 FOR PESTICIDES REGISTERED UNDER...



ACTIVE INGREDIENT		
SODIUM SALT OF DALAPON	7 Dils. a prop. 85	
INERT INGREDIENT	Equivalent 74	15
TOTAL		100

EPA REG. NO. 2749 52
NET WEIGHT 50lbs

WARNING

MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. MAY BE HARMFUL IF INHALED OR SWALLOWED. AVOID BREATHING VAPOUR OR SPRAY MIST. USE ONLY WITH ADEQUATE VENTILATION. AVOID CONTACT WITH EYES AND SKIN. WASH THOROUGHLY AFTER HANDLING. IN CASE OF CONTACT, FLUSH EYES WITH PLENTY OF WATER FOR AT LEAST 15 MINUTES AND GET MEDICAL ATTENTION. FOR SKIN, REMOVE CONTAMINATED CLOTHING AND WASH SKIN WITH PLENTY OF WATER. WASH CLOTHING BEFORE REUSE.

THIS PRODUCT IS TOXIC TO SHRIMP. KEEP OUT OF TIDAL MARSHES AND ESTUARIES. DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOUR DRIFT FROM AREAS TREATED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE. OBSERVE ALL CAUTIONS AND LIMITATIONS ON LABELLING OF ALL PRODUCTS USED IN MIXTURES. DALAPON IN THE SOIL MAY INHIBIT SEED GERMINATION AND PLANT GROWTH FOR 30 DAYS OR MORE DEPENDING ON DOSAGE APPLIED, SOIL TYPE, TEMPERATURE AND AMOUNT OF SOIL MOISTURE AND RAINFALL. ACCORDINGLY, ON LAND TO BE PLANTED TO SENSITIVE CROPS SUCH AS GRASS, SMALL GRAINS, CORN OR BEANS, DELAY PLANTING FOR AT LEAST 30 DAYS AFTER PLOWING AREAS SPRAYED WITH DALAPON. DECOMPOSITION OF DALAPON PROCEEDS MOST RAPIDLY IN WARM, MOIST SOIL. IF SOIL REMAINS COOL OR DRY, DECOMPOSITION MAY BE SLOWER AND MORE TIME WILL BE REQUIRED AFTER PLOWING BEFORE CERTAIN CROPS CAN BE PLANTED SAFELY TO AVOID THE POSSIBILITY OF RESIDUES IN MEAT OR MILK. DO NOT GRAZE MEAT OR DAIRY ANIMALS ON VEGETATION TREATED WITH DALAPON TO AVOID INJURY TO DESIRABLE PLANTS. DO NOT ALLOW SPRAY OR SPRAY DRIFT CONTAINING DALAPON TO CONTACT FOLIAGE OR ROOTS OF SUCH PLANTS INCLUDING LAWNS AND OTHER DESIRABLE TURF. FLUSH SPRAYING EQUIPMENT WITH PLENTY OF WATER IMMEDIATELY AFTER USE TO AVOID CONTAMINATION OF LATER SPRAYS OR POSSIBLE OTHER HERBICIDES. BE SURE TO READ THE DIRECTIONS AND PRECAUTION ON THE HERBICIDE CONTAINER.

DALAPON IS USED TO CONTROL ALL ANNUAL GRASSES SUCH AS CRABGRASS, FOXTAIL AND WATER GRASS AND ESTABLISHED PERENNIAL GRASSES SUCH AS BLUEGRASS, BERMUDA GRASS, JOHNSON GRASS, PHRAGMITES, PARAGRASS AND CATTAILS. DALAPON IS MOST EFFECTIVE AS A FOLIAGE SPRAY. ALTHOUGH IT MAY ALSO BE ABSORBED THROUGH PLANT ROOTS, DALAPON MAY BE USED IN COMBINATION WITH HERBICIDES SUCH AS MCP AMINE AND SIMLX. APPLY DALAPON WHEN GRASSES ARE GROWING VIGOROUSLY BUT BEFORE SEED STALKS DEVELOP. SPRAY THOROUGHLY AND UNIFORMLY. SEVERAL WEEKS MAY BE REQUIRED FOR MAXIMUM EFFECT. REPEAT APPLICATION AS NEEDED.

DIRECTIONS FOR USE

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.

Apply 1.5 to 2.0 lbs. of active ingredient per acre of grasses to be controlled. For control of annual grasses, apply when grasses are in the seedling stage. For control of established perennials, apply when grasses are in the flowering stage. For control of crabgrass, foxtail, and water grass, apply when grasses are in the seedling stage. For control of bluegrass, bermuda grass, johnson grass, phragmites, paragrass, and cattails, apply when grasses are in the flowering stage. For control of grasses in lawns and other desirable turf, apply when grasses are in the seedling stage. For control of grasses in pastures, apply when grasses are in the seedling stage. For control of grasses in fields, apply when grasses are in the seedling stage. For control of grasses in woods, apply when grasses are in the seedling stage. For control of grasses in brush, apply when grasses are in the seedling stage. For control of grasses in meadows, apply when grasses are in the seedling stage.