

ACCEPTED

OCT 16 1972

UNLAWFUL GENERAL INSECTICIDE
FUNGICIDE AND RODENT CIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 2749-70 SUBJECT
TO ATTACHED COMMENTS.

SPUD NIC

EMULSIFIABLE CONCENTRATE
POTATO SPROUT INHIBITOR

EPA REG. NO. 2749-70
NET CONTENTS 30 GALLONS

ONE GALLON OF SPUD NIC EMULSIFIABLE CONCENTRATE
WILL TREAT 2,880 X LBS. SACKS (288,000 LBS.) OF POTATOES.

ACTIVE INGREDIENT:
ISOPROPYL-M-CHLOROCARBANILATE* 36.0%
INERT INGREDIENTS: 64.0%
TOTAL 100.0%

*CONTAINS 3 LBS. OF ACTIVE INGREDIENT PER GALLON.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.
HARMFUL IF SWALLOWED. DO NOT TAKE INTERNALLY.
AVOID BREATHING OF VAPOR OR SPRAY MIST. USE WITH
ADEQUATE VENTILATION. AVOID CONTACT WITH EYES,
SKIN AND CLOTHING. IN CASE OF CONTACT, FLUSH WITH
PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION.
KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS.
AVOID CONTAMINATION OF FERTILIZERS, SEED, PLANTS
AND PESTICIDES.

KEEP CONTAINER CLOSED WHEN NOT IN USE. DO NOT
REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER
BY PERFORATING OR CRUSHING AND BURYING IN A SAFE
PLACE.

AVOID STORAGE OF SPUD NIC BELOW FREEZING TEMPERA-
TURES. PROLONGED STORAGE COULD RESULT IN CRYSTAL
FORMATION. IF THIS HAPPENS WARMING CONTENTS AND
AGITATION WILL RESTORE MATERIAL TO USABLE CON-
DITION. DO NOT HEAT ABOVE 250°F.


SPUD NIC EMULSIFIABLE CONCENTRATE IS USED FOR
TREATING SPROUTING POTATOES AFTER STORAGE AND
WASHING. SPUD NIC EMULSIFIABLE CONCENTRATE WILL
PREVENT WOUND PERIDERM FORMATION OF POTATOES,
AND THEREFORE, IT SHOULD ONLY BE USED AFTER
BRUISES AND CUTS HAVE HEALED. THIS NORMALLY TAKES
A MINIMUM OF 2 WEEKS.

DO NOT APPLY IN FIELD. DO NOT USE ON SEED POTATOES.
WASH EQUIPMENT THOROUGHLY WITH SOAP AND WATER
BEFORE USING THE EQUIPMENT FOR SEED POTATOES.

DIRECTIONS FOR USE:

TO INHIBIT SPROUTING OF POTATOES: THE INHIBITION OF
SPROUTING AT RECOMMENDED RATES IS USUALLY EF-
FECTIVE FOR UP TO 1 YEAR. SPUD NIC EMULSIFIABLE
CONCENTRATE IS APPLIED AS A 1% EMULSION ON THE
DRYING TABLE AFTER THE POTATOES HAVE BEEN WASHED.
DILUTE SPUD NIC EMULSIFIABLE CONCENTRATE TO A 1%
EMULSION BY ADDING 1 GALLON OF SPUD NIC EMULSIFI-
ABLE CONCENTRATE TO 35 GALLONS OF WATER. AGITATE

TO FORM A UNIFORM EMULSION. APPLY AT RATE OF 1
QUART OF 1% EMULSION PER 20 X 100 LBS. SACKS (2,000
LBS.) OF POTATOES. SPRAY NOZZLE SHOULD BE SET TO
SPRAY EVENLY ACROSS THE WASHER ROLLS MOVING THE
POTATOES. THE DRYER ROLLS WILL DISTRIBUTE THE
DILUTED SPUD NIC TO ASSURE COMPLETE COVERAGE OF
EACH POTATO.



Manufactured for:

ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICAL DIV.

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368

CHLORPROPHAM TECHNICAL

ACTIVE INGREDIENT:

Isopropyl N-(3-Chlorophenyl) Carbamate 98.0%

INERT INGREDIENTS: 2.0%

TOTAL 100.0%

NET WEIGHT _____
USDA REG. NO. 2749-117

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

KEEP OUT OF THE REACH OF DOMESTIC ANIMALS AND FOODSTUFFS.

KEEP OUT OF LAKES, STREAMS, PONDS AND OTHER BODIES OF WATER.

KEEP AWAY FROM HEAT AND OPEN FLAME.

May be harmful if inhaled or swallowed. Use with adequate ventilation. Do not inhale spray mist. Avoid contact with eyes, skin, nose and throat. In case of contact, flush with plenty of water. If swallowed, induce vomiting and get medical attention. Do not reuse empty container. Destroy empty container by perforating or crushing and burying in a safe place.

DIRECTIONS FOR USE: FOR USE ONLY BY COMMERCIAL PESTICIDE FORMULATORS.

ACCEPTED
2749-117
MAY 26 1971
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PEST CONTROL
REGISTERED UNDER NO. *2749-117*



Manufactured for:

ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICAL DIVISION

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368