

ACCEPTED
SEP - 6 1960

ACTIVE INGREDIENT	
Isopropyl m-chlorocarbamate	36.0%
INERT INGREDIENTS	64.0%
TOTAL	100.0%


CAUTION
KEEP OUT OF REACH OF CHILDREN

SHAKE WELL BEFORE USING

NET CONTENTS: GALLONS

1.0 GALLON

Manufactured for ACETO AGRICULTURAL CHEMICALS CORP., 126-02 Northern Boulevard, Flushing, New York 11368



CAUTION

Harmful if swallowed. Do not take internally. Avoid breathing of vapor or spray mist. Use with adequate ventilation. Avoid contact with eyes, skin and clothing. In case of contact, flush with plenty of water. For eyes, get medical attention. Keep away from domestic animals and foodstuffs. Avoid contamination of fertilizers, seed, plants and pesticides.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Do not store near heat or open flame. Keep container tightly closed when not in use. Avoid storage of SPUD NIC below freezing temperatures. Prolonged storage could result in crystal formation. If this happens, gradual warming to room temperature (75°F) and agitation will restore material to usable condition. Do not use direct heat or flame.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Shake well before using

SPUD NIC-3 is used for treating sprouting potatoes after storage and washing. SPUD NIC-3 will prevent wound periderm formation of potatoes, and therefore, it should only be used after bruises and cuts have healed. This normally takes a minimum of 2 weeks.

Do not apply in field. Do not use on seed potatoes.


Wash equipment thoroughly with soap and water before using the equipment for seed potatoes.

To inhibit sprouting of potatoes: The inhibition of sprouting at recommended rates is usually effective for up to 1 year. SPUD NIC-3 is applied as a 1% emulsion on the drying table after the potatoes have been washed. Dilute SPUD NIC-3 to a 1% emulsion by adding 1 gallon of SPUD NIC-3 to 35 gallons of water. Agitate to form a uniform emulsion. Apply at rate of 1 quart of 1% emulsion per 20 x 100 lbs. sacks (2,000 lbs.) of potatoes. Spray nozzle should be set to spray evenly across the washer rolls moving the potatoes. The dryer rolls will distribute the diluted SPUD NIC-3 to assure complete coverage of each potato.

DISCLAIMER

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

BEST AVAILABLE COPY



FLAMMABLE LIQUID, NOS UN1993

NOT REVIEWED 09-5-85
In Accordance with FR Notice 82-2
Based on Draft Labeling Dates

SPUD NIC-3

EMULSIFIABLE POTATO SPROUT INHIBITOR
One gallon of SPUD NIC-3 will treat 2,888 x 100 lbs. sacks (288,800 lbs.) of potatoes.

ACTIVE INGREDIENT:		
Isonopryl-m-chlorocarbamate*	36.0%
INERT INGREDIENTS:	64.0%
	TOTAL	100.0%

*Contains 3 lbs. Active Ingredient per gallon

CAUTION

KEEP OUT OF REACH OF CHILDREN


See Side Panel For Additional Precautionary Statements

SHAKE WELL BEFORE USING

NET CONTENTS GALLONS

EPA REG. NO. 2749-70

EPA EST. NO.



CAUTION

Harmful if swallowed. Do not take internally. Avoid breathing of vapor or spray mist. Use with adequate ventilation. Avoid contact with eyes, skin and clothing. In case of contact, flush with plenty of water. For eyes, get medical attention. Keep away from domestic animals and foodstuffs. Avoid contamination of fertilizers, seed plants and pesticides.

DISCLAIMER

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Do not store near heat or open flame. Keep container tightly closed when not in use.

Avoid storage of SPUD NIC-3 below freezing temperatures. Prolonged storage could result in crystal formation. If this happens, gradual warming to room temperature (75°F) and agitation will restore material to usable condition. Do not use direct heat or flame.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

of potatoes, and therefore, it should only be used after bruises and cuts have healed. This normally takes a minimum of 2 weeks.

Do not apply in field. Do not use on seed potatoes.

Wash equipment thoroughly with soap and water before using the equipment for seed potatoes.

To inhibit sprouting of potatoes: The inhibition of sprouting at recommended rates is usually effective for up to 1 year. SPUD NIC-3 is applied as a 1% emulsion on the drying table after the potatoes have been washed. Dilute SPUD NIC-3 to a 1% emulsion by adding 1 gallon of SPUD NIC-3 to 35 gallons of water. Agitate to form a uniform emulsion. Apply at rate of 1 quart of 1% emulsion per 20 x 100 lbs. sacks (2,000 lbs.) of potatoes. Spray nozzle should be set to spray evenly across the washer rolls moving the potatoes. The dryer rolls will distribute the diluted SPUD NIC-3 to assure complete coverage of each potato.

Shake well before using.

SPUD NIC-3 is used for treating sprouting potatoes after storage and washing. SPUD NIC-3 will prevent wound periderm formation.

Manufactured For:
**ACETO AGRICULTURAL
 CHEMICALS CORPORATION**
 126-02 Northern Blvd.
 Flushing, New York 11358