

ZINEB 75-W

NET WEIGHT 55 LBS

U.S.D.A. REG. NO. 2749-76

ACTIVE INGREDIENT	
ZINEB (ZINC ETHYLENE BISDITHIOCARBAMATE)	75%
INERT INGREDIENTS	25%
TOTAL	100%

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS

MAY BE HARMFUL IF INHALED OR SWALLOWED. DO NOT BREATHE DUST OR SPRAY MIST. AVOID CONTACT WITH EYES, SKIN OR CLOTHING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES GET MEDICAL ATTENTION.

KEEP CONTAINER CLOSED. KEEP AWAY FROM FIRE AND SPARKS. DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

DIRECTIONS FOR USE:

Pour recommended amount of this material through screen on surface of water in nearly filled spray tank or pre mix in a bucket half filled with water and pour mixture into nearly filled spray tank. Add balance of water to fill tank. Do not allow mixture to stand. Keep agitator running during filling, mixing and spraying operations. Do not use in low volume gear type spray equipment. This material is compatible with most of the commonly used insecticides, fungicides and nutritional sprays. Do not combine wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials being combined has proved them to be physically compatible.

The dosage rates on this label indicate pounds of ZINEB 75-W per 100 gallons of dilute spray. Use 100 to 125 gallon spray per acre for low crops such as beans, carrots and onions. Use 150 to 200 gallon spray per acre for large, heavier crops such as cucurbit, and tomato vines. Make thorough, full coverage applications.

Consult State Experimental Station or State Extension Service specialist for specific information as the timing, number and rate of applications needed will vary with local conditions.

FRUITS:

APPLE: Apple Blotch, Black Rot, Bitter Rot, Botryosphaeria Fruit Rot, Brooks Spot, Cedar Rust, Fly Speck, Frog-eye, Quince Rust, Scab, Sooty Blotch. Use 1 to 2 lbs. Begin applications in delayed dormant and continue through cover sprays. Use 1 to 1 1/2 lbs. when combined with Captan. Do not use within 7 days of harvest.

APPLE, PEACH: Fire Blight. Use 2 lbs. at 2 to 3 day intervals during periods of bloom opening and wet weather. Maintain recommended sanitary pruning practices. Do not apply within 7 days of harvest.

APRICOT, PEACH: Coryneum Blight (Shot Hole), Peach Leaf Curl. Use 2 lbs. in

postharvest or dormant sprays and in popcorn to jacket stages according to local spray program. CITRUS: Rust Mite and associated Russetting and Greasy Spot. Use 1 lb. Apply with thorough coverage. Make application from late June through early August, or earlier if rust mite begins to show.

SOUR CHERRY: Leaf Spot or Shot Hole. Use 1 lb. Apply at petal fall and 10 and 20 days after petal fall and immediately after harvest. Do not apply within 7 days of harvest.

CURRENT: Leaf Spot. Use 2 lbs. Apply 2 to 3 weeks after bloom. Repeat at 7 to 10 day intervals and immediately after harvest. Do not apply more than 200 gallons per acre. Do not apply within 7 days of harvest.

GRAPE: Black Rot, Downy Mildew, Ripe Rot. Use 1 1/2 lbs. just before bloom, just after bloom, 10 days after bloom, and 2 to 3 weeks later. Do not apply within 7 days of harvest.

STRAWBERRY: Leaf Spot. Use 1 1/2 to 2 lbs. Begin applications when new growth starts in spring and repeat at 7 to 10 day intervals until fruit begins to form. Leaf Scorch—Use 1 1/2 to 2 lbs. in new beds in late summer. Make 2 to 3 applications at 7 day intervals. Do not apply within 7 days of harvest.

VEGETABLES:

Use 1 1/2 to 2 lbs. per 100 gallons of dilute spray. Begin applications at first sign of disease and repeat at 7 to 10 day intervals or as needed to maintain control of the disease stated.

BEANS: Anthracnose—Begin when first blossoms appear. Downy Mildew, Rust. Begin when disease threatens. Do not apply within 7 days of picking.

BEETS, KALE, MUSTARD GREENS, SWISS CHARD, TURNIP: Cercospora, Leaf Spot, Downy Mildew. Do not apply within 7 days of harvest.

CARROT: Alternaria, Cercospora, Leaf Spot. Do not apply within 7 days of harvest if tops are to be used as food or feed.

CELERY: Early Blight, Late Blight. Begin applications when plants emerge in the plant beds. Repeat at 3 to 5 day intervals until plants are set in field. Continue applications at 7 day intervals. Remove residues remaining at harvest by washing and stripping.

CUCURBITS (Cantaloupe, Cucumber, Melons, Squash): Alternaria (Macrosporium) Blight, Anthracnose, Downy Mildew, Gummy Stem Blight. Direct spray to underside of leaves. Do not apply within 7 days of harvest on summer squash.

LETTUCE: Downy Mildew. Do not apply within 5 days of harvest on head lettuce or 7 days on leaf lettuce.

ONION: Botrytis Leaf Blight (Blast), Downy Mildew, Purple Blotch. Do not apply within 7 days of harvest on green or spring onions.

SPINACH: Downy Mildew (Blue Mold), White Rust. Do not apply within 7 days of harvest.

TOMATO: Anthracnose, Cladosporium Gray Mold, Early Blight, Septoria Leaf Spot, Stemphylium Leaf Spot.



Aceto Chemical Company
1000 North Main Street
P.O. Box 100
Cincinnati, Ohio 45201