

PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS
(& DOMESTIC ANIMALS)**

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause irritation of eyes, nose, throat and skin. May be harmful if inhaled or swallowed.

Do not breathe dust or spray mist. Use with adequate ventilation. Avoid contact with eyes, skin or clothing. Keep away from fire and sparks.

In case of contact, flush with plenty of water. For eyes, get medical attention.

If swallowed, induce vomiting by giving two glasses of water and sticking finger in throat.

Never give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

DO NOT APPLY DIRECTLY TO WATER

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with this labeling.


KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS

STORAGE AND DISPOSAL

Storage
Store in a cool dry area. Do not let AMAZIN PHASE 2 FUNGICIDE become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness and create vapors which may be flammable. Keep container closed when not in use.

Container Disposal
Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State or local authorities.

Pesticide Disposal
Pesticide spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures.



ACTIVE INGREDIENT:	
Maneb (Manganese Ethylenebisidithiocarbamate).....	80%
INERT INGREDIENTS:	20%
	TOTAL 100%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Statement of Practical Treatment

May Be Harmful If Swallowed

May Be Harmful if Inhaled.

May Cause Irritation Of Eyes, Nose, Throat And Skin.

In case of skin contact, flush with plenty of water.

In case of eye contact, flush with plenty of water – Get Medical Attention.

See Side Panel For Additional Precautionary Statements.

EST. NO. 3306-HL-1

EPA REG. NO. 2749-88-AA

NET CONTENTS: LBS.

**MFG. FOR: Aceto Agricultural Chemicals Corporation
126-02 Northern Boulevard
Flushing, New York 11368**

CONTROLS DISEASES ON THESE CROPS

FIELD CROPS

Sugar Beets
Corn
Tobacco

FRUITS AND NUTS

Almonds
Apples
Apricots
Cranberries
Grapes
Nectarines
Peaches

VEGETABLES

Broccoli
Cabbage
Cantaloupe
Carrots
Celery
Cucumbers
Endive
Lettuce
Lima Beans
Onions
Peppers
Potatoes
Pumpkins
Spinach
Squash
Tomatoes
Watermelon

APPLICATION FOR SPECIFIC CROPS:

ALMONDS (CALIFORNIA): Shot-hole (Fungus) and Brown Rot (Blossom and twig blight) — Use 1½ to 2 lbs. per 100 gals. (6 to 8 lbs. per acre). Begin application at the dormant stage. Apply in popcorn, full bloom and petal fall or every 7-10 days if bloom is staggered. Do not apply later than 5 weeks after petal fall. If applied after petal fall, do not feed hulls to dairy animals or animals being finished for slaughter.

APPLES: Scab, Cedar Apple Rust, Black Rot, Fly Speck, Sooty Blotch, Bitter Rot, White Rot—Use 1 to 2 lbs. per 100 gallons or up to 10½ lbs. per acre in concentrated sprays. Apply in pre-pink and repeat in pre-bloom, bloom and cover sprays. Follow state spray schedules.

Do not graze livestock in treated areas. Do not apply within 15 days of harvest in Arkansas, Delaware, Illinois, Indiana, Kansas, Kentucky, Maryland, Missouri, New Jersey, North Carolina, Ohio, Pennsylvania, South Carolina, Tennessee, Virginia, and West Virginia. Do not apply within 30 days of harvest in other states.

APRICOTS (CALIFORNIA): Brown Rot and Shot-hole (fungus) — Use 2 lbs. at red bud, early bloom and full bloom. On Green Rot or Jacket Rot, use 2 lbs. at full bloom and petal fall.

BROCCOLI: Downy Mildew—Apply 4 lbs. per acre. As a spray, use 100 to 150 gallons per acre. As an 8% dust, use 50 lbs. per acre. Apply when disease threatens and continue as needed at 7 to 14 day intervals. Shorten intervals when disease conditions are severe. Repeat after rains. If applied less than 3 days before harvest, remove residues by trimming and washing.

CABBAGE: Alternaria leaf spots and Downy Mildew—As a spray, use 1½ lbs. per 100 gallons and apply up to 150 gallons per acre. As a 8% dust, use 10½ to 26 lbs. per acre.

Apply 7 to 10 days after planting. If disease is present, apply earlier. Repeat every 3 days until plants are set in the field or until field-seeded plants are thinned. Do not apply within 7 days of picking.

CANTALOUPE AND SQUASH (TEXAS): Downy Mildew—Apply when disease threatens, repeat as needed every 7 to 10 days.

Do not apply within 5 days of harvest.

CARROTS: Alternaria and cercospora leaf spots — Apply when plants are 6 weeks old and repeat every 7 to 10 days.

CELERY: Early Blight (Cercospora) and Late Blight (septoria) — Apply when plants emerge in plant beds. Repeat at 3- to 5-day intervals in plant beds and 7- to 10-day intervals after plants are set in the field.

Do not apply within 14 days of harvest.

CORN: Helminthosporium Leaf Blight—Use 2 to 2½ lbs. per acre. Apply when disease first threatens. Repeat every 4 to 7 days. Do not feed treated forage to livestock.

CUCUMBERS: Downy Mildew, Anthracnose—Apply when disease threatens, repeat every 7 to 10 days. Do not exceed more than 3½ pounds AMAZIN PHASE 2 MANEB per acre per application. Do not apply within 5 days of harvest.

GRAPES (IN GREAT LAKES AREA): Black Rot—Apply up to 5 lbs. per acre. Do not apply later than 10 days after bloom. If you apply within 7 days of harvest. Do not apply more than 1½ lbs. per acre.

LETTUCE & ENDIVE: Downy Mildew—As a spray, use up to 150 gals. per acre. As a dust, use up to 50 lbs. per acre. Apply when disease is present. Repeat every 7 to 10 days. Do not apply within 10 days of picking. Remove residues from head of lettuce by stripping and trimming. Remove residues from leaf lettuce and endive by thorough washings.

LIMA BEANS: Downy Mildew and Anthracnose—As a spray, use up to 200 gallons per acre. As a dust, use 40 lbs. per acre. Apply when disease first appears. Repeat every 4 to 7 days as needed. Do not apply within 4 days before picking.

ONIONS: Downy Mildew, Purple Blotch (Blight) and Blaw (Botrytis Leaf Blight)—Apply when first spots appear. Repeat every 7 days as needed.

PEACHES: (Pacific States) and **NECTARINES** (California)—Shot-hole (Fungus) and Peach Leaf Curl (California only)—Apply dormant and popcorn sprays. For Brown Rot Blossom Blight, apply at pink bud and 25% to 75% full bloom (add a petal fall spray in Pacific Northwest). For fruit Brown Rot (California only), apply 2 lbs. per 100 gallons 4 weeks before harvest. If rot is found on green fruit, apply earlier and repeat at 7 to 14 day intervals as needed. On peaches, if applied within 14 days of harvest, remove residue by brushing. Do not apply later than 2 days before harvest. On nectarines, do not apply later than 14 days before harvest.

PEPPERS: Anthracnose, Ripe Rot, Cercospora Leaf Spot (Frog-eye Spot)—Use 3 lbs. per 100 gallons. For Cercospora Leaf Spot, apply when disease threatens. Repeat as needed at 7 to 10 day intervals. For Anthracnose and Ripe Rot, apply when the first fruits begin to turn red. Repeat as long as disease threatens at 7 to 10 day intervals. Do not apply more than 3½ lbs. AMAZIN PHASE 2 MANEB per acre per application.

POTATOES: Early and Late Blights—Use 1 to 2 lbs. per acre. Apply when plants are 2 to 6 inches high; repeat every 5 to 10 days for as long as blights threaten.

PUMPKINS: Downy Mildew, Angular Leaf Spot—Use 3 lbs. per acre. Use in a sufficient amount of water to wet plants thoroughly. Apply when disease threatens; repeat at 7 to 10 day intervals as needed.

ROSES: Cercospora Leaf Spot (Texas), Rust (California), Black Spot—Use 1½ lbs. Spray when first leaves unfold; repeat every 7 days as needed.

SPINACH: Downy Mildew (Blue Mold) and White Ruts—Apply during first sign of disease. Repeat every 7 to 10 days. Do not apply within 10 days of picking. Remove excess residues by thorough washing.

SUGAR BEETS: Cercospora Leaf Spot—As a spray, use 1½ to 2 lbs. per acre in 75 to 125 gallons of water. As a dust, use 30 to 40 lbs. per acre. Apply when disease threatens. 3 to 5 applications should be made. Repeat every 7 to 10 days as needed. Do not apply within 14 days of harvest at the spray dosage. At the dust dosages, do not apply within 10 days of harvest and do not feed treated tops to livestock.

TOBACCO: (North Carolina) Blue Mold—Use one-half pound per 100 gallons. Use 2½ to 5 gallons of diluted suspension per 100 sq. yds. Spray plant beds when plants are the size of a dime or when disease is reported in the area. Repeat every 3 days or more often if worked with rain. Continue until transplanting is complete.

TOMATOES: Early Blight, Late Blight Septoria Leaf Spot, Gray Leaf Spot (Stemphylium) and Anthracnose—Apply when fruit of first clusters is well formed. Repeat every 7 to 10 days. When weather conditions favor disease, more frequent applications may be necessary. When Anthracnose is a serious problem on canning tomatoes, use 3 lbs. per acre. Do not use on tender, young seedlings in green houses as injury may occur. Do not exceed more than 3½ lbs. AMAZIN PHASE 2 MANEB per acre per application. Do not apply within 5 days of harvest.

WATERMELONS: Downy Mildew, Anthracnose, Cercospora Leaf Spot, gummy stem blight—Use 1½ to 2 lbs. per 100 gallons of water. Apply as a full coverage spray up to 250 gallons per acre. Apply when runners form or when disease threatens. Repeat as needed at 7 to 10 day intervals. Under severe conditions, shorten intervals to 3 to 5 days. Do not apply within 5 days of harvest.

DIRECTION FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
DILUTION: Dilute this FLEXGARD with up to 10% by volume of water to make a Ready-To-Use solution containing not less than 2% Bis (tri-n-Butyltin) Oxide by weight.

APPLICATION

1. FLEXGARD waterbase preservative is intended for outdoor use only. Apply to wood products that will not be in contact with soil, such as boats, wood pilings, millwork, shingles and sidings.
2. Repeat application on textiles such as nylon web, rope and canvas as, if or when mildew appears.
3. The surface should be cleaned and dry before application.
4. Stir contents of container before each use.
5. Apply liberally by brush or dipping. Thoroughly cover cut ends and joints.
6. If applied by brush, use two coats.
7. If applied by dipping, allow article to remain in solution for 3 minutes.
8. Allow 6-8 hours drying time between coats. Longer drying time will not cause any adverse effect on performance.
9. Allow overnight drying before use.
10. Clean tools, protective gloves and clothing with water when finished.
11. Do not apply at or below freezing temperatures.

COVERAGE: Estimated coverage for diluted product: 275 square feet or less per gallon depending on application.

STORAGE AND DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Keep from freezing.

Household: Do not reuse empty container (bottle, can, bucket). Wrap (container) and put in trash collection.

Pesticide Disposal (except household): pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

Container Disposal (except household): metal containers-Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other approved state and local procedures.

General: Consult Federal, state and local disposal authorities for approved alternative procedures.

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals.

WARNING: May be fatal if swallowed, may be absorbed through skin. Causes skin irritation. Do not get in eyes, on skin or on clothing. Do not breathe vapor or spray mist. Wear protective clothing and rubber gloves when handling this product.

USE WITH ADEQUATE VENTILATION

Vapor: May cause injury if adequate ventilation is not insured. Do not use this product indoors, or any other confined areas, where vapors may concentrate and cause injury to plant or animal life. Keep containers closed when not in use.

PHYSICAL OR CHEMICAL HAZARDS: Not combustible (Waterbase)

ENVIRONMENTAL HAZARDS

This product is toxic to domestic animals. Treated sawdust and other wastes should not come in contact with domestic animals. Avoid contamination of feed or foodstuffs.

This product is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

Treated sawdust and other wood wastes should not be used as mulch where it will come in contact with useful living plants. Care should be taken to prevent drip or spray from contacting ornamental shrubs, trees, grass and other desirable vegetation.

NON WARRANTY: The information presented in this label is based upon the research and experience of Flexabar and is to the best of its knowledge accurate; however, no guarantee of its accuracy is made. Since Flexabar has no control over the conditions under which its products may be employed, and since information as to the physiological and other properties of its products is incomplete, Flexabar cannot guarantee the results obtained and assumes no liability whatsoever for possible damage or injury whether or not caused by following the suggestions in this label. Flexabar will not be liable for resulting or consequential damage. Flexabar warrants that none of its commercial products infringes the claims of any United States patent but no license is granted nor recommendation made hereby to practice any invention covered by any patent owned by or licensed to Flexabar or by others.