

PHOSALONE EC

FOR CONTROL OF INSECT PESTS AND EPIDERM MITES

ACTIVE INGREDIENT:

Phosalone O,O -Diethyl S - $\bar{Z}$ -(o -chloro-2-oxobenzoxazolin-
3:1)methylphosphorodithioate 34.8%

INERT INGREDIENTS:

TOTAL 65.2%
100.0%

BEST DOCUMENT AVAILABLE

*Equivalent to 3 pounds of phosalone per gallon.

NET CONTENTS: 5 GALLONS

EPA REG. NO: _____

WARNING: KEEP OUT OF REACH OF CHILDREN.

Phosalone is swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin. Do not get in eyes, on skin or on clothing. Wear protective clothing, natural rubber gloves and goggles. Do not breathe fumes or spray mist. Wear a mask or respirator effective for protection against PHOSALONE. Information as to suitable types of masks or respirators is available from the USDA. In case of contact, wash skin thoroughly with soap and water; for eyes, flush with plenty of water and get medical attention. Wash thoroughly after using and before eating or smoking. Wash contaminated clothing with soap and hot water before reuse.

NOTE TO PHYSICIAN: Causes cholinesterase inhibition. Atropine or 2-PAM (Pralidoxime methyl sulfate) are antidotal. The combined therapy of atropine and 2-PAM is more effective than either substance alone. Warning symptoms include weakness, headache, tightness in the chest, blurred vision, nonreactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

This product is toxic to fish and wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

Do not reuse empty container. Destroy empty container by perforating or crushing and burning in a safe place.

Do not store below 32°F. Keep away from heat and open flame.

Do not allow livestock to graze on crops in treated orchards.

DIRECTIONS FOR USE

BEST DOCUMENT AVAILABLE

GENERAL: PHOSALONE EC is an emulsifiable concentrate and is to be used with water. It is for use in all Power Operated Ground Sprayers and by hand. Keep spray mix agitated when spraying. Mix the quantities of PHOSALONE EC shown with water to give adequate coverage. For best results, mix with 100 to 150 gals. of water and apply as a complete coverage spray, normally, 1 to 200 sq. ft. per acre.

CROP	INSECT	PTS. 100 GALS.	SPECIAL INSTRUCTIONS
APPLES and PEARS	Aphidid (1st) Codling Moth	2.3 to 1-1/2	Make applications to local fruit pest control programs. Do not apply more than 17 1/2 lbs. (46-2 1/3 pts) per acre per season. Some russet-tinged Golden Delicious and other yellow varieties may follow use of PHOSALONE in some areas. Do not apply within 14 days of harvest. *except phosphate resistant strains.
	Red Banded Leaf Roller, Pear Leaf Rollers	1 to 2-2/3	
	Plum Curculio *Two-spotted Mite	1-1/2 to 2-2/3	
PEARS	Pear Psylla	2	
GRAPES (2)	Mites (including Pacific Mite)	2-1/2	Repeat application as required but not more than 2 lbs. (5 1/3 pts) per acre per season. Do not apply within 14 days of harvest. Applicable rates for leafhopper control on grapes are from 5-1/2 to 10-2/3 pts. per acre per application.
	Leaf Hopper	1	
	Leaf Hopper	5-1/2	

- (1) Registration for control is limited to the Northern United States.
(2) Registration is limited to California.

MINERAL OIL

AGRI-CHEMICAL COMPANY, INC.
Agricultural Chemical Division
11402 Northern Boulevard,
New York, N.Y. 11420

13