



RAMPART 15-G

15% GRANULES SYSTEMIC LOW-ODOR FORMULA INSECTICIDE FOR CONTROL OF CORN ROOTWORM AND WESTERN CORN ROOTWORM ON CORN.

WARNING

Apply this product only as specified on this label.

Not for use or storage in or around the house.

Keep away from domestic animals and foodstuffs.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

ENVIRONMENTAL CAUTIONS – This product is toxic to fish, shrimp, crabs and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab may be killed at the recommended dosage rates. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

In case of poisoning, contact physician immediately.

Note to Physician: Atropine is Antidote. 2-PAM is also Antidotal and may be administered in conjunction with Atropine. Warning symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

EPA REG. NO.
2749-105

ACTIVE INGREDIENT:

PHORATE (0,0-Diethyl S-(ethylthio) methyl-phosphorothioate).....	15%
INERT INGREDIENTS	85%
TOTAL	100%

DANGER:

KEEP OUT OF THE REACH OF CHILDREN



POISON

POISON

ANTIDOTE IS ATROPINE

Poisonous by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Do not get in eyes, on skin or clothing. Wear freshly laundered long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently. In case of contact, immediately remove contaminated clothing, wash thoroughly with soap and water before eating or smoking. Do not breathe dust. While emptying bags into equipment, pour downward and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach breathing zone. Fields may be reentered on day of treatment with soil applications of the product.

SEE WARNING STATEMENTS AND ANTIDOTE ON LEFT PANEL.

NET WEIGHT _____

GENERAL DIRECTIONS

WHEN USING RAMPART 15-G USE BY POURING DOWN WIND TO MINIMIZE EXPOSURE TO HANDLERS. SPILLAGE SHOULD BE REMOVED AND BURNED.

APPLY RAMPART 15-G WITH A GRANULAR PESTICIDE APPLICATION PROPERLY CALIBRATED TO ASSURE ACCURATE PLACEMENT AND PROPER DOSAGE.

DIRECTIONS FOR USE:

CORN (at planting time) – For control of corn rootworms: Use 4 to 6.5 lbs. per acre (40 inch row spacing) or 4 to 8 ozs. per 1000 feet of row (for any row spacing). Apply RAMPART 15-G in a 7 inch band over the row directly behind the planter shoe in front of the press wheel. Do not place RAMPART 15-G in direct contact with seed. Use the 6.5 lb. rate in areas having high rootworm populations.

CORN (post-emergence) – For control of corn rootworms: Use 4 to 6.5 lbs. per acre (40 inch row spacing) or 4 to 8 ozs. per 1000 feet of row (for any row spacing). Apply RAMPART 15-G at time of cultivation in a band at base of plants just ahead of cultivator shovels so as to cover granules with soil. Apply as soon as rootworm activity is observed. Do not make any later applications after emergence treatment.

NOTICE

FOLLOW DIRECTIONS CAREFULLY. TIMING AND METHOD OF APPLICATION, WEATHER AND CROP CONDITIONS, MIXTURES WITH OTHER CHEMICALS NOT SPECIFICALLY RECOMMENDED, AND OTHER INFLUENCING FACTORS IN THE USE OF THIS PRODUCT ARE BEYOND THE CONTROL OF THE SELLER. BUYER ASSUMES ALL RISKS OF USE, STORAGE AND HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

EPA EST. NO.
960-CO-02

ACCEPTED

treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

In case of poisoning, contact physician immediately.

Note to Physician: Atropine is Antidote. 2-PAM is also Antidotal and may be administered in conjunction with Atropine. Warning symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

EPA REG. NO.
2749-105

ANTIDOTE IS ATROPINE

Poisonous by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Do not get in eyes, on skin or clothing. Wear freshly laundered long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently. In case of contact, immediately remove contaminated clothing, wash thoroughly with soap and water before eating or smoking. Do not breathe dust. While emptying bags into equipment, pour downward and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach breathing zone. Fields may be reentered on day of treatment with soil applications of the product.

SEE WARNING STATEMENTS AND ANTIDOTE ON LEFT PANEL.

NET WEIGHT _____

per acre (40 inch row spacing) or 4 to 8 ozs. per 1000 feet of row (for any row spacing). Apply RAMPART 15-G in a 7 inch band over the row directly behind the planter shoe in front of the press wheel. Do not place RAMPART 15-G in direct contact with seed. Use the 6.5 lb. rate in areas having high rootworm populations.

CORN (post-emergence) - For control of corn rootworms: Use 4 to 6.5 lbs. per acre (40 inch row spacing) or 4 to 8 ozs. per 1000 feet of row (for any row spacing). Apply RAMPART 15-G at time of cultivation in a band at base of plants just ahead of cultivator shovels so as to cover granules with soil. Apply as soon as rootworm activity is observed. Do not make any later applications after emergence treatment.

NOTICE

FOLLOW DIRECTIONS CAREFULLY. TIMING AND METHOD OF APPLICATION, WEATHER AND CROP CONDITIONS, MIXTURES WITH OTHER CHEMICALS NOT SPECIFICALLY RECOMMENDED, AND OTHER INFLUENCING FACTORS IN THE USE OF THIS PRODUCT ARE BEYOND THE CONTROL OF THE SELLER. BUYER ASSUMES ALL RISKS OF USE, STORAGE AND HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

EPA EST. NO.
960-CO-02



Manufactured for:

ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICALS DIVISION

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368

