

**BOVINOX 80% SOLUBLE POWDER
ANIMAL INSECTICIDE**

ACTIVE INGREDIENT:


Dimethyl-(2,2,2-trichloro-1-hydroxymethyl)-phosphonate* 80%

INERT INGREDIENTS: 20%

TOTAL 100%

*Trichlorfon

EPA REG. NO. 2749-107
NET WEIGHT _____



**FOR CONTROL OF HORN FLIES, LICE
AND GRUBS ON BEEF
CATTLE.**

WARNING: KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM DOMESTIC ANIMALS AND
FOODSTUFFS.

THIS PRODUCT IS TOXIC TO FISH, WILDLIFE AND BIRDS. DO
NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR
DISPOSAL OF WASTES.

MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED
THROUGH THE SKIN. DO NOT GET IN EYES OR ON SKIN. DO
NOT BREATHE DUST OR SPRAY MIST. ALWAYS WASH THOR-
OUGHLY WITH SOAP AND WARM WATER AFTER HANDLING.
WASH CONTAMINATED CLOTHING WITH SOAP AND WARM WA-
TER BEFORE RE-USE. DO NOT CONTAMINATE FEED OR FOOD.

IF POISONING OCCURS, GET MEDICAL ATTENTION IMMEDIATE-
LY. PROLONGED EXPOSURE WILL RESULT IN CHOLINESTER-
ASE DEPRESSION.

NOTE TO PHYSICIAN: ATROPINE SULFATE IS ANTIDOTE. 2-PAM
IS ALSO ANTIDOTE AND MAY BE USED IN CONJUNCTION WITH
ANTROPINE.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CON-
TAINER BY PERFORATING OR CRUSHING AND BURYING IN A
SAFE PLACE.

BOVINOX 80% SOLUBLE POWDER IS A HIGHLY EFFECTIVE INSECTICIDE
THAT CAN BE USED IN ANY TYPE OF CONVENTIONAL SPRAY EQUIP-
MENT. BOVINOX 80% SOLUBLE POWDER COMPLETELY DISSOLVES IN
WATER FOR A MORE EFFECTIVE SPRAY SOLUTION. BOVINOX 80% SOLU-
BLE POWDER GIVES SYSTEMIC CONTROL OF GRUBS AND GIVES RESIDU-
AL CONTROL OF LICE AND HORN FLIES.

DIRECTIONS FOR USE:

FOR CATTLE GRUB CONTROL—TREATMENT OF CATTLE WITH BOVINOX 80% SOLU-
BLE POWDER CAN BE MADE ANY TIME DURING THE GRUB SEASON. HOWEVER,
FOR BEST RESULTS, TREAT EARLY IN THE GRUB CYCLE. TREAT AS SOON AS POS-
SIBLE AFTER ACTIVITY OF HORN FLY HAS STOPPED AND AT LEAST 6 WEEKS PRIOR
TO EXPECTED APPEARANCE OF GRUBS. YOU MAY APPLY BOVINOX 80% SOLUBLE
POWDER VERY NEAR OR DURING THE TIME GRUBS APPEAR BUT CONSIDERABLE
GRUB DAMAGE WILL HAVE ALREADY OCCURRED. USE AS SPRAY TREATMENT.
MIX 10 LBS. OF BOVINOX 80% SOLUBLE POWDER IN 100 GALLONS OF WATER. AP-
PLY THIS SOLUTION FOR COMPLETE WETTING TO RUN-OFF. APPLY IN A MANNER
THAT THE SKIN, NOT JUST THE HAIR, BECOMES THOROUGHLY WET. SPRAY PRES-
SURES OF 250 TO 350 LBS. ARE SUGGESTED. OPERATE BOX-TYPE SPRAY CHUTES
AT MAXIMUM PRESSURE FOR THOROUGH OVER-ALL WETTING AND PENETRATION.
A SINGLE APPLICATION IS ADEQUATE FOR GRUB CONTROL. DO NOT USE WITHIN
14 DAYS OF SLAUGHTER.

CATTLE HORN FLY CONTROL—USE WHEN HORN FLIES APPEAR AND CONSTITUTE A
PROBLEM. SPRAY TREATMENT: MIX 10 LBS. OF BOVINOX 80% SOLUBLE POWDER
IN 100 GALLONS OF WATER. APPLY THIS SOLUTION FOR COMPLETE WETTING AND
RUN-OFF. REPEAT AS NECESSARY BUT DO NOT USE WITHIN 14 DAYS OF SLAUGH-
TER.

CATTLE LICE CONTROL—USE WHEN LICE APPEAR AND CONSTITUTE A PROBLEM.
USE AS SPRAY TREATMENT: MIX 2½ LBS. OF BOVINOX 80% SOLUBLE POWDER IN
100 GALLONS OF WATER. APPLY THIS SOLUTION FOR COMPLETE WETTING TO
RUN-OFF. REPEAT AS NECESSARY BUT DO NOT USE WITHIN 14 DAYS OF SLAUGH-
TER.

DO NOT TREAT DAIRY ANIMALS. DO NOT TREAT ANIMALS LESS THAN 3 MONTHS
OLD. DO NOT TREAT ANIMALS THAT ARE SICK, CONVALESCENT, OR STRESSED
LIVESTOCK. DO NOT SPRAY IN A CONFINED, NON-VENTILATED AREA. DO NOT
USE IN CONJUNCTION WITH ORAL DRENCHES, OTHER MEDICATIONS, OR OTHER
ORGANIC PHOSPHATE-TYPE SPRAYS.



Manufactured For

ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICALS DIV

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368