

**PDQ  
4-(MCPB) SODIUM SALT  
WEED KILLER**

NET CONTENTS 5 GALLONS  
REG. NO. 2749-109

FOR CONTROL OF CANADA THISTLE AND CERTAIN OTHER WEEDS IN PEAS.

ACTIVE INGREDIENT  
SODIUM SALT OF 4-(2-METHYL-4-CHLOROPHENOXY)  
BUTYRIC ACID\* 23.5  
INERT INGREDIENTS 76.5  
100%

\*EQUIVALENT TO 23.5 LBS. OF 4-(2-METHYL-4-CHLOROPHENOXY) BUTYRIC ACID OR 27.5 LBS. OF 4-(2-METHYL-4-CHLOROPHENOXY) BUTYRIC ACID PER GALLON AT 8.3 F.

**CAUTION:** KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS

KEEP OUT OF LAKES, STREAMS, PONDS AND OTHER BODIES OF WATER.

MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. AVOID CONTACT WITH EYES, SKIN, NOSE, THROAT AND CLOTHING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

AVOID SPRAY DRIFT TO NEARBY CROPS. COARSE SPRAYS ARE LESS LIKELY TO DRIFT.

DO NOT STORE NEAR FERTILIZERS, SEEDS OR OTHER PESTICIDES. THOROUGHLY CLEAN SPRAY EQUIPMENT AFTER USING WITH PDQ. DO NOT USE EQUIPMENT TO SPRAY OTHER PESTICIDES UNTIL THE EQUIPMENT IS THOROUGHLY CLEAN. PROTECT FROM FREEZING. IF PDQ FREEZES, THAW AND REMIX BEFORE USING.

**GENERAL INFORMATION**

PDQ controls Canada Thistle better than any other herbicide used on peas. It is quite useful on early peas, but may be used for all varieties.

PDQ mixes readily with water and is suitable for use with all conventional spraying equipment.

PDQ controls Canada Thistle and will also control or suppress, depending on their stage of growth, other susceptible weeds such as annual morning glory, fanweed, lambsquarters, nightshade, pigweed, smartweed and sowthistle.

Sometimes temporary twisting of some pea varieties may occur after an application. Most canning varieties of peas may be effectively sprayed with PDQ but before using, consult with your State Extension Service or pea canning company field man for varietal differences to recommendations listed.

**DIRECTIONS FOR USE**

Apply PDQ when annual broadleaf weeds have emerged and before they are 3 inches tall. Canada Thistle should be sprayed after good shoot emergence has developed. Do not apply PDQ later than 3 nodes before first pea flowering as adverse effects on pea yields may result. Do not spray peas that lack moisture or when temperatures exceed 90 F. Applications made under adverse growing conditions (water logged soils or during periods of drought) will result in poor weed control and will injure peas.

**RECOMMENDED DOSAGE  
(DO NOT EXCEED RECOMMENDED DOSAGE)**

WEEDS AND STAGE OF GROWTH	AMOUNT OF PDQ TO USE	AMOUNT OF WATER TO USE
Canada Thistle 4 to 10 inches high	3 to 4 parts per acre	20 to 30 gals per acre
Other susceptible weeds less than 2 inches high (2 to 3 leaves)	2 to 3 parts per acre	10 to 15 gals per acre
Other susceptible weeds less than 2 inches high (2 to 3 leaves)	2 to 3 parts per acre	10 to 15 gals per acre
Other susceptible weeds less than 2 inches high (2 to 3 leaves)	2 to 3 parts per acre	10 to 15 gals per acre
Other susceptible weeds less than 2 inches high (2 to 3 leaves)	2 to 3 parts per acre	10 to 15 gals per acre



ACETO CHEMICALS