

DIREZ 50W TURF FUNGICIDE

ACTIVE INGREDIENT:

2,4-Dichloro-6-(0-Chloroanilino) -s-Triazine 50%

INERT INGREDIENTS: 50%

TOTAL 100%

NET WEIGHT _____

USDA REG. NO. 2749-120

WARNING: KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS.

THIS PRODUCT IS TOXIC TO FISH. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.

MAY BE HARMFUL IF SWALLOWED. MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. AVOID CONTACT WITH EYES, SKIN OR CLOTHING. AVOID INHALATION OF DUST. WASH THOROUGHLY AFTER HANDLING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

DO NOT GRAZE TREATED AREAS. DO NOT FEED CLIPPINGS TO LIVESTOCK.

DIREZ 50W TURF FUNGICIDE IS A BROAD SPECTRUM FUNGICIDE FOR MAJOR TURF AND LAWN DISEASES SUCH AS:

- HELMINTHOSPORIUM DISEASES
(MELTING-OUT, LEAF SPOT)
- COPPER SPOT
- SCLEROTINIA DOLLAR SPOT
- SNOW MOLD (TYPHULA)
- RUST
- BROWN PATCH

DIREZ 50W TURF FUNGICIDE IS TINTED GREEN SO THAT IT CAN BLEND WITH THE TURF AND ELIMINATE AN UNSIGHTLY APPEARANCE. DIREZ 50W TURF FUNGICIDE WILL NOT HARM SPRAY EQUIPMENT, CLOG NOZZLES OR CORRODE METAL PARTS OF THE SPRINGER. FOR A PREVENTATIVE PROGRAM, BEGIN APPLICATIONS BEFORE DISEASES APPEAR. THE DIREZ SPRAY PROGRAM SHOULD BE CONTINUED THROUGH THE SPRING AND SUMMER MONTHS.

DIRECTIONS FOR USE:
COMPLETE AND THOROUGH COVERAGE IS NECESSARY FOR BEST RESULTS.

DISEASE	POUNDS OF DIREZ 50W TURF FUNGICIDE	REMARKS
Brown patch, Copper Spot, Dollar Spot, Leaf Spot, Melting-out, Rust	¼ to ½	Apply specified dosage in 5 gallons of water per 1000 sq. ft. Repeat at 1 to 2 week intervals throughout the spring summer months.
Snow Mold (Typhula)	¼	Apply in 5 gallons of water per 1000 sq. ft. Apply after first killing frost in the fall. Make second application when the lawn is free of snow in mid-winter. Make third application after the final melting of snow in late winter or early spring.

ACCEPTED
OCT 5 1971
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 2749-120

DIREZ 50W TURF FUNGICIDE EFFECTIVELY PROTECTS GRASSES FROM VARIOUS DESTRUCTIVE DISEASES. COMPLETE AND THOROUGH COVERAGE OF AREA IS NECESSARY FOR SATISFACTORY CONTROL. FOR A PREVENTATIVE PROGRAM, BEGIN APPLICATIONS BEFORE DISEASES APPEAR AND REPEAT TREATMENTS AS NEEDED, USUALLY AT INTERVALS OF 10 TO 14 DAYS, EXCEPT DURING EXTENDED PERIODS OF COOL, MOIST CONDITIONS, WHEN APPLICATIONS EVERY 7 DAYS MAY BE ADVISABLE. THE DIREZ FUNGICIDE SPRAY PROGRAM SHOULD BE CONTINUED THROUGH THE SPRING AND SUMMER MONTHS.

PREVENTATIVE PROGRAM:
FOR CONTROL OF BROWN PATCH, COPPER SPOT, DOLLAR SPOT, LEAF SPOT, MELTING-OUT, RUST, SNOW MOLD (TYPHULA)—USE ¼ TO ½ POUND OF DIREZ 50W TURF FUNGICIDE IN 5 GALLONS OF WATER PER 1000 SQUARE FEET.



ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICALS DIV

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368