

# DIREZ 50 WP FOLIAGE FUNGI DE

**ACTIVE INGREDIENT:**

2,4-Dichloro-6 (0-Chloroanilino) -s-Triazine . . . 50%

**INERT INGREDIENTS:** . . . . . 50%

TOTAL 100%

**NET WEIGHT** \_\_\_\_\_

EPA REG. NO. 2749-122

**WARNING:** KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS.

THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS OR PONDS. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.

MAY BE HARMFUL IF SWALLOWED. MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. AVOID CONTACT WITH EYES, SKIN OR CLOTHING. AVOID INHALATION OF DUST. WASH THOROUGHLY AFTER HANDLING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

**DIRECTIONS FOR USE:**

Complete and thorough coverage of foliage is necessary for satisfactory control. Start applications before disease appears and repeat treatment as needed, usually at 7 to 10 day intervals. During wet weather, applications of 4 to 5 day intervals may be advisable. The DIREZ spray program should be continued throughout the growing season. Schedule applications with local conditions. Consult your State Agricultural Experimental Station for specific use information.

CROP	DISEASE	POUNDS PER ACRE OF DIREZ 50WP	REMARKS
<b>FIELD CROPS</b>			
Tobacco	Brown Spot (Alternaria)	2 lbs.	Apply specified dosage using approximately 100 gallons of water per acre for good coverage. Apply at weekly intervals during harvest, after each priming.
<b>FRUIT</b>			
Blackberries Dewberries Loganberries Raspberries	Cane and leaf rust (Kuehneola uredinis) Botrytis fruit rot Cane and leaf spot (Septoria rubi)	2-4 lbs.	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage. Do not apply within 5 days of harvest for strawberries nor within 14 days of harvest for the other berries.
Strawberries	Gray Mold (Botrytis) Leaf Spot Leaf Scorch	2-6 lbs.	
Blueberries Huckleberries	Double Spot Septoria Leaf Spot	3 lbs.	Apply specified dosage using 150 gallons of water per acre for good coverage. Do not apply within 14 days of harvest. Restricted for use only in North Carolina.
Cranberries	Berry Spot Black Rot End Rot Phomopsis Rot	6 lbs.	Apply specified dosage using 300 gallons of water per acre for good coverage. Do not apply within 14 days of harvest. Restricted for use only in Wisconsin.
<b>VEGETABLES</b>			
Celery	Early Blight Late Blight Rhizoctonia	2-6 lbs.	Apply specified dosage using 75 to 200 gallons of water per acre for good coverage. Treated celery should be washed and trimmed.
Cucumbers	Anthrachnose Downy Mildew Gummy stem blight Leaf Spots (Alternaria) (Cercospora)	4-6 lbs.	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage.

CROP	DISEASE	POUNDS PER ACRE OF DIREZ 50WP	REMARKS
Onions (bulb or dry) Garlic	Botrytis Purple Blotch (Alternaria)	2-6 lbs.	Apply specified dosage using 75 to 200 gallons of water per acre for good coverage. Not recommended for use in Florida.
Onions (Green) Shallots	Botrytis Purple Blotch (Alternaria)	2-3 lbs.	Apply specified dosage using 75 to 100 gallons of water per acre for good coverage. Not recommended for use in Florida.
Potatoes	Early Blight Late Blight	2 to 5 lbs.	Apply specified dosage using 75 to 175 gallons of water per acre for good coverage.
Pumpkins Squash	Anthrachnose Downy Mildew Gummy stem blight Leaf spots (Alternaria) (Cercospora)	4 to 6 lbs.	Apply specified dosage using 75 to 200 gallons of water per acre for good coverage.
Tomatoes	Anthrachnose Botrytis Early Blight Gray leaf spot Late Blight Septoria leaf spot	2 to 5 lbs.	Apply specified dosage using 75 to 175 gallons of water per acre for good coverage.
<b>MELONS</b>			
Cantaloupes Honeydew Muskmelons Watermelons Other Varieties	Anthrachnose Downy Mildew Gummy stem blight Leaf spots (Alternaria) (Cercospora)	4 to 6 lbs.	Apply specified dosage using 100 to 200 gallons of water per acre for good coverage.
<b>ORNAMENTALS</b>			
Gladioli	Leaf Spot (Stemphylium)	2 lbs.	Apply specified dosage in 100 gallons of water as a full coverage spray. Repeat twice weekly during periods of infection. Recommended for use in Florida only.

DO NOT EXCEED RECOMMENDED DOSAGE. DO NOT USE ON SEED CROPS INTENDED TO BE USED FOR FEED, FOOD OR FORAGE.

8 APR 1975



Manufactured for:

**ACETO CHEMICAL CO., INC.**  
AGRICULTURAL CHEMICALS DIV.

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368

ACETOPROTECTANT