




**PHORATE 15-G
15% GRANULES SYSTEMIC INSECTICIDE**



WARNING

Apply this product only as specified on this label.

Not for use or storage in or around the house.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOOD-STUFFS.

not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

ENVIRONMENTAL CAUTIONS — This product is toxic to fish, shrimp, crabs and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab may be killed at the recommended dosage rates. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

In case of poisoning, contact physician immediately.

Note to Physician: Atropine is Antidote. 2-PAM is also Antidotal and may be administered in conjunction with Atropine. Warning symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

EPA REG. NO.
2749-123

ACTIVE INGREDIENT:
PHORATE (0,0-Diethyl S-(ethylthio) methyl)-
phosphorothioate)..... 15%
INERT INGREDIENTS 85%
TOTAL 100%

DANGER:

KEEP OUT OF REACH OF CHILDREN.



POISON

POISON

ANTIDOTE IS ATROPINE

Poisonous by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Do not get in eyes, on skin or clothing. Wear freshly laundered long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently. In case of contact, immediately remove contaminated clothing, wash thoroughly with soap and water before eating or smoking. Do not breathe dust. While emptying bags into equipment, pour downward and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach breathing zone. Fields may be reentered on day of treatment with soil applications of the product.

SEE WARNING STATEMENTS AND ANTIDOTE ON LEFT PANEL.

NET WEIGHT _____

GENERAL DIRECTIONS:

WHEN USING PHORATE 15-G, USE BY POURING DOWN WIND TO MINIMIZE EXPOSURE TO HANDLERS. SPILLAGE SHOULD BE REMOVED AND BURNED.

APPLY PHORATE 15-G WITH A GRANULAR PESTICIDE APPLICATION PROPERLY CALIBRATED TO ASSURE ACCURATE PLACEMENT AND PROPER DOSAGE.

DIRECTIONS FOR USE:

POTATOES—FOR CONTROL OF APHIDS, LEAFHOPPERS, LEAF MINERS, PSYLLIDS, WIRE WORMS, FLEA BEETLE LARVAE AND REDUCTION OF FLEA BEETLE ADULTS. USE 13.2 LBS. PER ACRE ON SANDY OR LIGHT SOILS (38-INCH ROW SPACING). USE 20 LBS. PER ACRE ON HEAVY SILT OR CLAY LOAM SOILS (38-INCH ROW SPACING) OR 14½ OZS. FOR SANDY OR LIGHT SOILS PER 1000 FT. OF ROW (FOR 38 INCH ROW SPACING, 23½ OZS. FOR HEAVY SILT OR CLAY LOAM SOILS PER 1000 FT. OF ROW (FOR ANY ROW SPACING). DISTRIBUTE THE GRANULES EVENLY IN THE FURROW OR GRANULES MAY BE BANDED ON EACH SIDE OF THE ROW AT PLANTING TIME. WAIT 90 DAYS AFTER TREATMENT BEFORE HARVESTING POTATOES.

SUGAR BEETS—FOR CONTROL OF APHIDS, BEET ROOT MAGGOTS, LEAFHOPPERS, MITES AND LEAF MINERS. USE 6.6 LBS. PER ACRE (30-INCH ROW SPACING) OR 6 OZS. PER 1000 FT. (FOR ANY ROW SPACING). APPLY BY DRILLING OR BROADCAST AT SEEDING TIME. DO NOT PLACE PHORATE 15-G IN DIRECT CONTACT WITH THE SEED. DO NOT APPLY WITHIN 30 DAYS OF HARVEST. DO NOT FEED TREATED TOPS OR SILAGE TO DAIRY ANIMALS.

NOTICE

FOLLOW DIRECTIONS CAREFULLY. TIMING AND METHOD OF APPLICATION, WEATHER AND CROP CONDITIONS, MIXTURES WITH OTHER CHEMICALS NOT SPECIFICALLY RECOMMENDED, AND OTHER INFLUENCING FACTORS IN THE USE OF THIS PRODUCT ARE BEYOND THE CONTROL OF THE SELLER. BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

EPA EST. NO.
960-CO-02



**PHORATE 15-G
15% GRANULES SYSTEMIC INSECTICIDE**

ACCEPTED
NOV 26 1975
Under the Federal Insecticide,
Fungicide and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 2749-123

WARNING

Apply this product only as specified on this label.

Not for use or storage in or around the house.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOOD-STUFFS.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

ENVIRONMENTAL CAUTIONS - This product is toxic to fish, shrimp, crabs and other wildlife. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab may be killed at the recommended dosage rates. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

In case of poisoning, contact physician immediately.

Note to Physician: Atropine is Antidote. 2-PAM is also Antidotal and may be administered in conjunction with Atropine. Warning symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pinpoint pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

EPA REG. NO.
2749-123

ACTIVE INGREDIENT:	
PHORATE (0,0-D ethyl S-(ethylthio) methyl)-phosphorothioate).....	15%
INERT INGREDIENTS	85%
	TOTAL 100%

DANGER:

KEEP OUT OF REACH OF CHILDREN.



POISON

POISON

ANTIDOTE IS ATROPINE

Poisonous by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Do not get in eyes, on skin or clothing. Wear freshly laundered long-sleeved work clothing daily. While transferring from package to equipment, wear a clean cap and gloves (rubber or cotton). If cotton gloves are used, they must be laundered or discarded after each day's use. Rubber gloves should be washed with soap and water after each use. Do not wear the same gloves for other work. Destroy and replace gloves frequently. In case of contact, immediately remove contaminated clothing, wash thoroughly with soap and water before eating or smoking. Do not breathe dust. While emptying bags into equipment, pour downward and allow as little free fall as possible. Do not pour at face level and do not allow dust to reach breathing zone. Fields may be reentered on day of treatment with soil applications of the product.

SEE WARNING STATEMENTS AND ANTIDOTE ON LEFT PANEL.

NET WEIGHT

GENERAL DIRECTIONS:

WHEN USING PHORATE 15-G, USE BY POURING DOWN WIND TO MINIMIZE EXPOSURE TO HANDLERS. SPILLAGE SHOULD BE REMOVED AND BURNED.

APPLY PHORATE 15-G WITH A GRANULAR PESTICIDE APPLICATION PROPERLY CALIBRATED TO ASSURE ACCURATE PLACEMENT AND PROPER DOSAGE.

DIRECTIONS FOR USE:

POTATOES-FOR CONTROL OF APHIDS, LEAFHOPPERS, LEAF MINERS, PSYLLIDS, WIRE WORMS, FLEA BEETLE LARVAE AND REDUCTION OF FLEA BEETLE ADULTS. USE 13.2 LBS. PER ACRE ON SANDY OR LIGHT SOILS (38-INCH ROW SPACING). USE 20 LBS. PER ACRE ON HEAVY SILT OR CLAY LOAM SOILS (38-INCH ROW SPACING) OR 14 1/2 OZS. FOR SANDY OR LIGHT SOILS PER 1000 FT. OF ROW (FOR 38 INCH ROW SPACING, 23 1/2 OZS. FOR HEAVY SILT OR CLAY LOAM SOILS PER 1000 FT. OF ROW (FOR ANY ROW SPACING). DISTRIBUTE THE GRANULES EVENLY IN THE FURROW OR GRANULES MAY BE BANDED ON EACH SIDE OF THE ROW AT PLANTING TIME. WAIT 90 DAYS AFTER TREATMENT BEFORE HARVESTING POTATOES.

SUGAR BEETS FOR CONTROL OF APHIDS, BEET ROOT MAGGOTS, LEAFHOPPERS, MITES AND LEAF MINERS USE 6.6 LBS. PER ACRE (30-INCH ROW SPACING) OR 6 OZS. PER 1000 FT. (FOR ANY ROW SPACING). APPLY BY DRILLING OR BROADCAST AT SEEDING TIME. DO NOT PLACE PHORATE 15-G IN DIRECT CONTACT WITH THE SEED. DO NOT APPLY WITHIN 30 DAYS OF HARVEST. DO NOT FEED TREATED TOPS OR SILAGE TO DAIRY ANIMALS.

NOTICE

FOLLOW DIRECTIONS CAREFULLY. TIMING AND METHOD OF APPLICATION, WEATHER AND CROP CONDITIONS, MIXTURES WITH OTHER CHEMICALS NOT SPECIFICALLY RECOMMENDED AND OTHER INFLUENCING FACTORS IN THE USE OF THIS PRODUCT ARE BEYOND THE CONTROL OF THE SELLER. BUYER ASSUMES ALL RISKS OF USE STORAGE OR HANDLING OF THIS MATERIAL NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH.

EPA EST. NO.
960 CO-02



Manufactured for
ACETO CHEMICAL CO., INC.

AGRICULTURAL CHEMICALS DIVISION

12602 NORTHERN BOULEVARD, ELK GLEN, N.C. 27628