

2749-126

1/28/2011

1/8



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

JAN 28 2011

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Fred Trudwig, Jr
Aceto Agriculture Chemicals Corp
One Hollow Lane, Suite 201
Lake Success, NY 11042

Dear Mr Trudwig,

Subject Add Use on Alfalfa and Label Revisions
2 4-DB 175 Herbicide
- EPA Registration No 2749-126
- Your Submission Dated January 25 2011

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you submit one (1) copy of your final printed labeling before you release the product for shipment

A stamped copy of the labeling is enclosed for your records

If you have any questions concerning this letter please contact Mr James Stone at 703-305-7391

Sincerely yours,

A handwritten signature in black ink, appearing to read "K. V. Montague".

Kathryn V Montague
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

2,4-DB 175

**FOR THE CONTROL OF SEEDLING BROADLEAF WEEDS IN PEANUTS,
SOYBEANS, AND SEEDLING STANDS OF ALFALFA,
AND FOR USE ON ESTABLISHED STANDS OF ALFALFA**

ACTIVE INGREDIENT:

Dimethylamine Salt of 4-(2,4-dichlorophenoxy) butyric acid* 23.00%

OTHER INGREDIENTS: 77.00%

TOTAL: 100.00%

*Equivalent to 19.5% by weight or 1.75 lbs. per gallon of 4-(2,4-dichlorophenoxy) butyric acid

**KEEP OUT OF REACH OF CHILDREN
DANGER – PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS


For spill, leak, fire, exposure or accident call CHEMTREC® 1-800-424-9300. Outside the USA 1-703-527-3887

EPA REG. NO. 2749-126

EPA EST. NO. _____

NET CONTENTS: GALS.

Manufactured For:
Aceto Agricultural Chemicals Corporation
One Hollow Lane
Lake Success, NY 11042
Phone: 516-627-6000



**PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
DANGER PELIGRO**

Corrosive Causes irreversible eye damage May be fatal if swallowed Harmful if absorbed through skin Do not get in eyes or on clothing Avoid contact with skin

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are made of any waterproof material If you want more options follow the instructions for category A on the EPA chemical resistance category selection chart

All mixers loaders applicators and other handlers must wear

- long sleeved shirt and long pants
- shoes and socks
- chemical resistant gloves (except for pilots)
- chemical resistant apron when mixing loading cleaning up spills cleaning equipment or otherwise exposed to concentrate and
- Goggles or face shield

See Engineering Controls for additional requirements

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables exist use detergent and hot water Keep and wash PPE separately from other laundry

Engineering Control Statement

Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for Agricultural Pesticides [40 CFR 170 240(d)(6)] Pilots must wear the PPE required on this labeling for applicators

USER SAFETY RECOMMENDATIONS

Users should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash the outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

FIRST AID

IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 to 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice
IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 to 20 minutes • Call a poison control center or doctor for treatment advice

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment You may also contact CHEMTREC 1 800 424 9300 outside the USA 1 703 527 3887 for emergency medical treatment information

NOTE TO PHYSICIANS

For eye irritation examination by an ophthalmologist may be indicated If swallowed probable mucosal damage may contradict the use of gastric lavage Product contains a phenoxy herbicide chemical There is no specific antidote

ENVIRONMENTAL HAZARDS

This chemical is toxic to fish Do not apply directly to water to areas where surface water is present or to intertidal areas below the mean high water mark Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas Do not contaminate water when disposing of equipment washwaters Do not contaminate water intended for irrigation or domestic purposes Do not apply when weather conditions favor drift form target area

Groundwater Contamination Most cases of groundwater contamination involving phenoxy herbicides such as 2 4 DB have been associated with mixing/loading and disposal sites Caution should be exercised when handling 2 4 DB pesticides at such sites to prevent contamination of groundwater supplies Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills Placement of the mixing/loading equipment on an impervious pad to contain spill will help prevent groundwater contamination

This chemical has properties and characteristics associated with chemicals detected in groundwater The use of this chemical in areas where soils are permeable particularly where the water table is shallow may result in groundwater contamination Application around a cistern or well may result in groundwater contamination

4/8

DIRECTIONS FOR USE

It is a violation of federal law to use this product in any manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, shoes plus socks, chemical-resistant gloves made of any waterproof material and protective eyewear.

PRODUCT INFORMATION

Apply this product only as specified on this label.

NOTE: Avoid spray drift to susceptible plants and crops: cotton, tobacco, tomatoes, ornamentals, etc. Coarse sprays are less likely to drift.

Spray equipment previously used to apply another phenoxy must be thoroughly cleaned with alkali and water prior to use with 2,4-DB 175. Similarly, after using 2,4-DB 175, wash sprayer thoroughly before using again to spray susceptible crops. Observe all caution and limitations on labeling of all products used in mixtures.

Use agitation to keep solution well mixed, especially if it has been allowed to stand unagitated. Calibrate sprayer accurately.

Do not apply when crop is stressed from lack of moisture.
Do not apply this product through any type of irrigation system.
Do not use in or near greenhouses.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2 to 10 mph at the application site.

Additional requirements for aerial applications:

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for the displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications into temperatures inversions.

Additional requirements for ground boom application:

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

TANK MIXES

Unless otherwise prohibited on this label or the label of an intended tank mix product, this product may be applied in combination with any pesticide registered for the same crop, timing, and method of application. Observe the most restrictive label statements of various tank mix products used.

IMPORTANT PESTICIDE TANK MIXES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS ANY LIABILITY FOR LOSS INJURY OR DAMAGE RESULTING FROM A TANK MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER

COMPATIBILITY

Before full scale mixing of this product with other pesticides fertilizers secondary plant nutrients adjuvants surfactants or oils you must determine the compatibility of the proposed mixture Use proportionate quantities of each ingredient and mix in a small container Always mix one product thoroughly with the diluent before adding another product If no incompatibility is evident after 30 minutes the mixture is generally compatible for spraying To evaluate potential short term effects of applying the mixture test the tank mix combination on a few plants or a small area before larger scale treatments Wait at least 2 to 3 days for problems to become apparent

IMPORTANT MIXING WITH OTHER SUBSTANCES MAY INCREASE THE RISK OF MIXING INCOMPATIBILITIES REDUCED EFFECTIVENESS AND/OR CAUSE CROP INJURY OR LOSS ANY LIABILITY FOR LOSS INJURY OR DAMAGE RESULTING FROM A MIXTURE NOT SPECIFIED ON THIS LABEL OR IN MANUFACTURER'S SUPPLEMENTAL LABELING DISTRIBUTED FOR THIS PRODUCT IS SPECIFICALLY DISCLAIMED BY MANUFACTURER

ALFALFA

Application

For use in seedling alfalfa spray when the crop has reached the 1 to 2 trifoliate leaf stage and growing conditions are good In established alfalfa certain weeds will emerge in the Fall and overwinter in a rosette stage Best control of these weeds will result from application in late Fall or early Winter rather than in the Spring Do not apply after flowering

When To Apply

Weeds must be in the young seedling stage and actively growing to achieve satisfactory results For best results spray weeds in the 2 to 5 leaf stage of growth

Amount To Use

Apply this product as an overall spray by ground sprayer or airplane Apply at rates listed below according to weed problems Use the higher rate if weeds are past the seedling stage Aerial applications should be made in at least 5 gallons of water per acre while ground sprayer applications should be in a minimum of 10 gallons of water per acre Higher spray gallonage per acre will give better coverage and weed control Spray gallonage should give adequate coverage of the weeds without runoff The use of a non ionic or non ionic/anionic surfactant or crop oil concentrate approved for agricultural uses at their label rates will usually result in better weed control especially if weeds are beyond the seedling stage

Broadleaf Weeds	Rate of 2.4 DB 175 Per Acre
Annual Morningglory (<i>Ipomoea spp</i>) Cocklebur (<i>Xanthium spp</i>) Common Lambsquarters (<i>Chenopodium album</i>) Jimsonweed (<i>Datura stramonium</i>) Kochia or Mexican Fireweed (<i>Kochia scoparia</i>) Pigweed (<i>Amaranthus spp</i>) Velvetleaf (<i>Abutilon theophrasti</i>) Wild Turnip (<i>Brassica campestris</i>)	4 1/3 Pints
Black Mustard (<i>Brassica nigra</i>) Buckhorn Plantain (<i>Plantago lanceolate</i>) Common Ragweed (<i>Ambrosia artemisiifolia</i>) Curly Dock (<i>Rumex crispus</i>) Field Pennycress (Fanweed or Stinkweed) (<i>Thlaspi arvense</i>) Hedge Smartweed (<i>Polygonum scandens</i>) Ladysthumb (<i>Polygonum persicaria</i>) Prickly Lettuce (<i>Lactuca scariola</i>) Shepherdspurse (<i>Capsella bursa pastoris</i>) Sweetclover (Volunteer plants) (<i>Melilotus spp</i>) Wild Beet (<i>Beta maritima</i>) Wild Mustard (<i>Brassica Kaber</i>)	6 1/2 Pints

Do not graze established alfalfa nor feed straw or hay from established alfalfa to meat or dairy livestock within 30 days after application Do not graze or feed seedling alfalfa to meat or dairy livestock within 60 days after application Irrigation in particular overhead sprinkler irrigation should be delayed as long as possible (10 days or more) following application of this product to avoid washing the chemical into the root zone Do not apply when crop is stressed from lack of moisture

Apply as a postemergent spray When properly timed there is little or no effect on the crop In established alfalfa there may be some twisting of stems and malformation of leaves This condition is usually outgrown

Do not spray when daytime temperatures are expected to exceed 90 F within the next 2 or 3 days Do not apply if temperatures are likely to fall below 40 F during or shortly after treatment

Use of a surfactant or crop oil concentrate in the desert areas of California and Arizona may cause some crop injury under certain climatic and crop stress conditions Do not use a surfactant unless possible crop injury is acceptable

PEANUTS

Application

Apply this product postemergent as an overall spray by ground sprayer or airplane With ground applications use a boom sprayer with flat fan type nozzle Adjust the height of the boom above the crop so the growing terminals of weeds are sprayed Use low pressure (about 20 to 40 pounds per square inch) Do not use boom jet nozzles With airplane applications this product has been successfully applied using 5 gallons of total solution per acre The height of the application should insure thorough coverage of the tops of the weeds Best control has resulted from airplane application with coarse spray when cocklebur is at least crop high and actively growing Other weed species should be in the seedling stage and actively growing Airplane application is not recommended in the vicinity of susceptible crops

Do not apply if peanuts are suffering from lack of moisture

Amount To Use

Apply the rates as indicated in the tables below

Broadleaf Weeds	Rate of 2 4 DB 175 Per Acre
Cocklebur (<i>Xanthium spp</i>)	1 Pint (1gallon per 8 acres)
Annual Morningglory (<i>Ipomoea spp</i>) Common Lambsquarters (<i>Chenopodium album</i>) Common Ragweed (<i>Ambrosia artemisifolia</i>) Jimsonweed (<i>Datura stramonium</i>) Sicklepod (Coffeeweed) (<i>Cassia obtusifolia</i>) Velvetleaf (<i>Abutilon theophrasti</i>)	1 3/4 Pints (1 gallon per 4 1/2 acres)

Do not apply more than two (2) applications per season second application should be made no later than the late bloom stage of peanuts (about 90 to 100 days after planting) Do not feed treated peanut vines or peanut hay to meat or dairy livestock Do not apply to peanuts within 60 days of harvest

SOYBEANS

Application

Apply this product postemergent as an overall spray by ground sprayer or airplane With ground applications use a boom sprayer with flat fan type nozzles Adjust the height of the boom above the crop so the growing terminals of all weeds are sprayed Use low pressure (about 20 to 40 pounds per square inch) Do not use boom jet nozzles With airplane applications this product has been successfully applied using 5 gallons of total spray solution per acre The height of the application should insure thorough coverage of the tops of the weeds Best control has resulted when cocklebur is at least crop high and actively growing Airplane application is not recommended in the vicinity of susceptible crops Do not apply if soybeans are suffering from lack of moisture

Topical (Overhead or Over The Top) Applications

Apply from 7 to 10 days before soybeans bloom through mid bloom A good indication that the soybeans are about to bloom is when the soybean plants have turned a dark green color If application is made 7 to 10 days before bloom apply one (1) gallon of this product per 10 acres (12 8 fl oz per acre) if application is made from early bloom through mid bloom apply 1 gallon of this product per 8 acres (1 pint per acre) Use sufficient water to obtain adequate coverage The rates are for the control of cocklebur These rates will also stunt or partially control certain other weeds including annual morningglory velvetleaf and jimsonweed In the Midwest use only a directed application since topical application may reduce soybean yields

NOTICE Application should not be made to sparsely foliated stands stressed because of disease or lack of moisture It is essential that the canopy above the soybeans be complete To apply the product otherwise could result in crop damage such as stalk and stem splitting and reduction in yield

Directed Applications

For best results apply when weeds do not exceed 3 inches in height Good coverage of the growing terminals of weeds is essential for effective control Do not allow spray pattern to contact more than the lower one third of the soybean plants

Do not apply more than two (2) applications per season Precise application is essential to prevent damage to the crop In order to maintain the correct spraying height nozzles must be mounted on oiling shoes skid shoes or on cultivators with gauge wheels Do not use booms with drop nozzles

First Application Apply as directed spray when soybeans are 8 to 12 inches tall and weeds have emerged

Second Application Apply as directed spray no later than mid bloom stage

Amount To Use Apply the rates as indicated in the table below

Broadleaf Weeds	Rate of 2 4 DB 175 Per Acre
Cocklebur (<i>Xanthium spp</i>)	1 Pint (1 gallon per 8 acres)

Annual Morningglory- (<i>Ipomoea spp</i>) Common Lambsquarters (<i>Chenopodium album</i>) Common Ragweed (<i>Ambrosia artemisifolia</i>) Jimsonweed (<i>Datura stramonium</i>) Sicklepod (Coffeeweed) (<i>Cassia obtusifolia</i>) Velvetleaf (<i>Abutilon theophrasti</i>)	1 3/4 Pints (1 gallon per 4 1/2 acres)
---	---

While some temporary twisting of soybean plants may follow an application this will have little or no permanent effect on the crop

Applications made under adverse growing conditions such as periods of drought will result in injury to the beans under such conditions the crop should not be sprayed

- Do not spray this product on soybeans showing a general infestation of *Phytophthora*
- Do not apply more than the amount specified on this label
- Do not harvest soybeans within 60 days after application
- Do not feed/graze soybean forage or harvest hay for 60 days following any 2.4 DB application
- Do not use on soybeans grown West of the Rocky Mountains

Directed Band Applications of 2.4 DB 175

Application

Apply directed spray when beans are at least 8 inches tall and when weeds do not exceed 2 inches in height. Direct spray to cover weed foliage with minimum contact of the soybean plant. Do not spray higher than 3 inches on the soybean stem or crop injury may result. Use an 8002T Jet (or equivalent) or larger with minimum of 25 gallons per broadcast acre and spray pressure of 20 to 25 psi to avoid spray drift. For each 25 gallons of spray mixture a pint of surfactant such as Surfactant WK may be added. Continuous agitation in the spray tank is required to keep the material in suspension. Agitate by mechanical or hydraulic means if bypass or return line is used it should terminate at the bottom of the tank to minimize foaming. Openings in nozzle screen should be equal to or larger than 50 mesh. Apply with sprayer nozzles mounted on skid shoes, oiling shoes, or on cultivators with gauge wheels. Do not mount on booms with drop nozzles or on cultivators without gauge wheels.

Amount to Use

Apply the rates as indicated in the table below

Broadleaf Weeds	Rate of 2.4 DB 175 Per Broadcast Acre
Annual Morningglory (<i>Ipomoea spp</i>) Cocklebur (<i>Xanthium spp</i>) Common Lambsquarters (<i>Chenopodium album</i>) Common Ragweed (<i>Ambrosia artemisifolia</i>) Jimsonweed (<i>Datura stramonium</i>) Velvetleaf (<i>Abutilon theophrasti</i>)	1 Pint

Use a proportional amount of this product on the band (for example on a 12 inch band in 36 inch rows use 1/3 pint per acre). If a new flush of weeds occurs after initial treatment make a second application but do not apply within 60 days of harvest or make more than 2 applications per growing season. Do not allow spray or spray drift to contact growing terminals of beans as excessive crop injury will result. Do not feed/graze soybean forage or harvest hay for 60 days following any 2.4 DB application. Harvested soybeans may be used for feed or oil purposes.

NOTE Do not use on sands, loamy sands, gravelly soils, or on exposed sub soils. Do not use on soils containing less than 1/2% organic matter as crop injury may result. Observe all cautions and limitations of labeling of all products used in mixture.

STORAGE AND DISPOSAL

Do not contaminate water, foodstuffs, feed or seed by storage or disposal. Open dumping is prohibited.

PESTICIDE STORAGE Store at temperatures above 32 F. If product is allowed to freeze, warm to 50 F and agitate thoroughly before using. Store in a secure warehouse or storage building inaccessible to children and domestic animals. Avoid storing near open containers of fertilizer, seed or other pesticides. Keep container sealed when not in use. Store only in original container unless an emergency requires that a different container replace a damaged container. In such case, clearly label contents. Reduce stacking height where local conditions can affect package strength.

PESTICIDE DISPOSAL Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Nonrefillable Containers 5 Gallons or Less Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Nonrefillable containers larger than 5 gallons: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Refillable containers larger than 5 gallons: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Aceto Agricultural Chemicals Corporation. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ACETO AGRICULTURAL CHEMICALS CORPORATION MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Aceto Agricultural Chemicals Corporation is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, ACETO AGRICULTURAL CHEMICALS CORPORATION DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT ACETO AGRICULTURAL CHEMICALS CORPORATION'S ELECTION, THE REPLACEMENT OF PRODUCT.

(RV010611 RED-Comments)