

JUL 22 1993

S444300 $\frac{361}{17}$

Mr. Anthony Baldi
Aceto Agricultural Chemicals Corporation
One Hollow Lake
Lake Success, New York 11042-1215

Dear Mr. Baldi:

Subject: Dimethogon 25WP Systemic Insecticide
EPA Registration No.: 2749-134
Your Letter Dated July 1, 1993

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

Note that this acceptance of your label does not relieve you of your obligation to comply with the Worker Protection Standard (WPS). If any of your products are covered by the WPS, you are required to submit, and receive the Agency's approval by April 21, 1994, of a revised label reflecting the required label statements of 40 CFR 156, published in the FEDERAL REGISTER of August 21, 1992 (57 FR 38102). Under the WPS labeling regulations at 40 CFR 156, subpart K, § 156.200(c)(3), you are prohibited from distributing or selling any product within the scope of the WPS requirements after April 21, 1994, without amended labeling accepted by the Agency.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,



Robert A. Forrest
Product Manager (14)
Insecticide-Rodenticide Branch
Registration Division (H7504C)

enclosure

MADDEN Disk #3 7/21/93



DIMETHOGON 25WP

SYSTEMIC INSECTICIDE

FOR USE ON APPLES, PEARS AND GRAPES

ACTIVE INGREDIENT:	
DIMETHOATE [O, O-Dimethyl 8-[N-(methylcarbamoyl) Methyl] Phosphorodithioate]	25%
INERT INGREDIENTS:	
	75%
TOTAL	100%

ACCEPTED
with COMMENTS
in EPA Letter Dated

JUL 22 1993

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2749-134

**KEEP OUT OF REACH OF CHILDREN
WARNING — AVISO**

PRECAUCION AL USARIO. Si usted no lee ingles, no use este producto hasta que le etiquete le haya sido explicado en español.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not give anything by mouth to an unconscious person. Get medical attention.

If on Skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If in Eyes: Flush with plenty of water for 15 minutes. Get medical attention.

**FOR 24-HOUR EMERGENCY ASSISTANCE,
Call CHEMTREC (800) 424-9300**

NOTE TO PHYSICIAN: This product may cause Cholinesterase inhibition. Antidote is Atropine.

NET CONTENTS: 40 LBS.

**REGISTRATION FOR SPECIAL LOCAL NEED
FOR DISTRIBUTION AND USE ONLY WITHIN CALIFORNIA.
TRADE NAME: ACETO DIMETHOGON 25 WP**

Formulated For:
ACETO Agricultural Chemicals Corp.
One Hollow Lane, Lake Success, NY 11042.

EPA Reg. No. 2749-134-AA
EPA SLN No. CA-780044

EPA Est. No. 10651-CA-5

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

STORAGE: Store in a cool, dry area.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

All applicators, flaggers and all personnel involved with the mixing, loading, and transferring operations must wear the protective clothing and equipment enumerated below. Pilots are exempt from this requirement. The protective clothing and equipment to be worn is as follows:

- a. Impermeable gloves (for example, rubber or plastic covered reinforced gloves).
- b. Boots or boot covers.
- c. Long-sleeved shirt and long pants.
- d. Wide-brimmed hat.
- e. Respirators must be worn by flaggers and mixer/loaders.

AERIAL APPLICATION

Automatic flagging devices should be used whenever feasible.

If human flaggers are employed, they must wear the protective clothing and respirator specified on this label.

DIRECTIONS FOR USE: Dimethogon 25WP mixes readily in the spray tank and will not clog nozzles. For proper mixing, the spray tank should be 3/4 full of water before you add Dimethogon 25WP. Agitate to maintain a good suspension. Do not apply this product through any type of irrigation system.

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
WARNING-AVISO**

May be fatal if inhaled, swallowed or absorbed through the skin. May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes and skin. Avoid breathing vapor or spray mist. Wash thoroughly after handling. In case of contact, flush with plenty of water for at least 15 minutes. Get medical attention.

Keep away from domestic animals and foodstuffs. Do not contaminate or apply onto feed or foodstuffs.

ENVIRONMENTAL HAZARD

This pesticide is toxic to wildlife and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system.

RE-ENTRY

Do not enter treated area without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

NOTE: THE RE-ENTRY PERIOD FOR GRAPES IS 4 DAYS. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers.

Written or oral warnings must include the following information:

WARNING

(Insert area or field description) treated with Dimethogon on (insert date of application). Do not enter treated areas without protective clothing until sprays have dried. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Dimethogon and describe his condition. For further information, see the center panel of the pesticide label.

5 8 5

For best Aphid control, Dimethogen 25WP should be applied from green tip through early cover sprays with additional treatments as may be necessary. Sprays should be spaced no later than 2 weeks apart with a minimum of 2 sequential sprays.

APPLES AND PEARS: For control of Aphids, Leafhoppers, Pear Psylla, Mites (except Rust Mites): Use 1 to 2 lbs. in 100 gallons of water in full coverage spray. Do not apply during the bloom period. Do not apply within 28 days of harvest.

Foliar application to apple:
Do not apply when trees or substantial numbers of weeds in the orchard (grove) are in bloom.

Do not graze livestock in orchards that have been treated.

GRAPES (Grown in California) RAISIN AND WINE GRAPES, TABLE AND CANNING GRAPES: For control of Grape Leafhopper and Pacific Spider mites: Use 6 to 8 lbs. per acre. Apply through concentrate spray equipment in 20 to 40 gallons of water per acre or through dilute equipment in 200 to 400 gallons of water per acre depending upon vine growth density. Repeat applications as necessary. Do not apply within 28 days of harvest.

DO NOT ENTER TREATED AREA WITHIN 4 DAYS.

Dimethogen 25% Wettable Powder Systemic Insecticide may leave a visible deposit of powder on berries at maturity if treatment is made after berries reach 1/4 inch in size to avoid objectionable deposits of powder on berries grown for table and canning purposes. Do not apply treatments to these crops after berries reach 1/4 inch in diameter.

NOTICE OF WARRANTY

ACETO warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label, only when used in accordance with directions under normal use conditions. It is not possible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of ACETO. In no case shall ACETO be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. ACETO makes no warranties of merchantability or fitness for a particular purpose nor any other express or implied warranty except as stated above.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Label: California Statewide

Crop/Site/Commodity: Grapes (Calif. Raisin, Wine, Table and Canning Grapes).

Target Pest/Problem: Grape Leaf Hopper, Pacific Spider Mite.

Dosage: Do not use more than 16 lbs./per acre.

Dilution rate and method of application: Aerial application — Apply in 15 - 30 gallons of water per acre. Ground application — Apply through concentrate spray equipment in 20 - 40 gallons of water per acre, or through dilute equipment in 200 - 400 gallons of water per acre depending upon vine growth density. Repeat application as necessary.

Field re-entry safety interval: 4 DAYS

Preharvest interval: 28 DAYS

All applicable directions, restrictions, and precautions on the EPA-Registered label(s) are to be followed:

The labeling must be in the possession of the user at the time of pesticide application.

Registrant: ACETO Agricultural Chemicals Corp.
One Hollow Lane, Lake Success, NY 11042.

BEST AVAILABLE COPY