

ROUND UP POUR-ON CATTLE INSECTICIDE
For control of cattle grubs

ACTIVE INGREDIENT:

C. O-Diethyl O-[3-chloro-4-methyl-2-oxo-(2H)-1-benzopyran-7-yl]-phosphorodithioate 4%

INERT INGREDIENTS: 96%
TOTAL 100%

NET CONTENTS 1 GALLON
EPA REG. NO. 2749-135

WARNING: KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM DOMESTIC ANIMALS AND FOOD-STUFFS. THIS PRODUCT IS TOXIC TO FISH, WILDLIFE AND BIRDS. KEEP OUT OF ANY BODY OF WATER. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREAS TREATED. DO NOT APPLY WHERE RUN-OFF IS LIKELY TO OCCUR. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

MAY BE POISONOUS IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. DO NOT BREATHE DUST OR SPRAY MIST. USE WITH ADEQUATE VENTILATION AND PROTECTIVE CLOTHING. AVOID CONTACT WITH EYES, SKIN, NOSE AND THROAT. IN CASE OF CONTACT, WASH THOROUGHLY WITH SOAP AND WATER. WASH CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE REUSE.

IF POISONING OCCURS, GET PROMPT MEDICAL AID.

NOTE TO PHYSICIAN: PROLONGED EXPOSURE WILL RESULT IN CHOLINESTERASE DEPRESSION. ANTIDOTE IS ATROPINE SULFATE. 2-PAM IS ALSO ANTIDOTAL AND MAY BE ADMINISTERED IN CONJUNCTION WITH ATROPINE.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR FLAME.

POUR THE CORRECT AMOUNT OF ROUND UP POUR-ON SOLUTION UNIFORMLY ALONG THE ANIMAL'S BACK. ROUND UP POUR-ON SOLUTION MAY BE DIPPED FROM ANY CLEAN BUCKET OR PAIL.

TREATMENT OF CATTLE CAN BE MADE AT ANY TIME DURING THE GRUB SEASON. HOWEVER, FOR BEST RESULTS, TREAT EARLY IN THE GRUB CYCLE. TREAT AS SOON AS POSSIBLE AFTER ACTIVITY OF HEEL FLY HAS CEASED AND PREFERABLY AT LEAST 6 WEEKS BEFORE EXPECTED APPEARANCE OF GRUBS IN THE BACK. TREATMENTS MADE NEAR OR DURING THE TIME GRUBS APPEAR IN THE BACK WILL DESTROY THE GRUBS BUT CONSIDERABLE DAMAGE WILL HAVE ALREADY OCCURRED.

DO NOT TREAT LACTATING DAIRY ANIMALS. DO NOT TREAT ANIMALS THAT ARE LESS THAN 3 MONTHS OLD. DO NOT TREAT CONVALESCENT OR STRESSED LIVESTOCK.

DO NOT TREAT NON-LACTATING DAIRY ANIMALS WITHIN 14 DAYS OF FRESHENING. IF FRESHENING SHOULD OCCUR WITHIN 14 DAYS AFTER TREATMENT, DO NOT USE MILK AS HUMAN FOOD FOR THE BALANCE OF THE 14 DAY INTERVAL.

DO NOT TREAT OVERHEATED ANIMALS.

DO NOT TREAT ANIMALS FOR 10 DAYS BEFORE OR AFTER SHIPPING OR WEANING, OR AFTER EXPOSURE TO CONTAGIOUS AND INFECTIOUS DISEASES.

DO NOT TREAT IN A CONFINED, NON-VENTILATED AREA.

DO NOT USE IN CONJUNCTION WITH ORAL DRENCHES OR OTHER INTERNAL MEDICATIONS SUCH AS PHENOTHIAZINE OR WITH NATURAL OR SYNTHETIC PYRETHROIDS OR THEIR SYNERGISTS, OR WITH OTHER ORGANIC PHOSPHATES.

APPLY ROUND UP POUR-ON SOLUTION AT THE RATE OF 1/2 FLUID OUNCE PER 100 LBS. OF BODY WEIGHT. MEASURE THE CORRECT AMOUNT TO USE IN RELATION TO THE ANIMAL'S WEIGHT.

ONE GALLON OF ROUND UP POUR-ON SOLUTION WILL TREAT:

Weight of Animal (in lbs.)	200	300	400	500	600	700	800	900	1000
Number of Animals	128	85	64	51	42	36	32	28	25



Manufactured For:

ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICALS DIVISION

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368

ACCEPTED
9/13/72
EPA REG. NO. 2749-135
INERT INGREDIENTS
RODENTICIDE
CATTLE
COMMENTS