

**AMROL**  
**LIQUID WEED KILLER**  
**FOR WEED CONTROL ON NON-CROPLAND**

Contains 2 lbs. per gallon of AMITROLE  
ACTIVE INGREDIENT  
AMITROLE (3-Amino-1,2,4-Triazole) 21.1%  
INERT INGREDIENTS 78.9  
TOTAL 100.0

\*Equivalent to 2 lbs. per gallon

EPA REG. NO. 2749-13  
NET CONTENTS

IN INTERMOUNTAIN STATES AND  
AREAS USE AMROL LIQUID WEED KILLER  
INSTEAD OF 2,4-D AMITROLE  
IN INTERMOUNTAIN STATES AND  
AREAS USE AMROL LIQUID WEED KILLER  
INSTEAD OF 2,4-D AMITROLE

**WARNING** KEEP OUT OF THE REACH OF CHILDREN

KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS

MAY CAUSE IRRITATION OF NOSE AND THROAT. MAY BE HARMFUL IF INHALED OR SWALLOWED. DO NOT BREATHE SPRAY MIST. USE WITH ADEQUATE VENTILATION. AVOID CONTACT WITH EYES, SKIN OR CLOTHING. WASH THOROUGHLY AFTER HANDLING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION.

DO NOT STORE NEAR FERTILIZERS, SEEDS OR PESTICIDES OR USE IN A GREENHOUSE. DO NOT SPRAY OR ALLOW SPRAY DRIFT TO CONTAMINATE NEARBY CROPS.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

IMPORTANT: THIS PRODUCT MUST NOT BE USED ON CROPLAND.

DIRECTIONS FOR USE ON NON-CROPLAND SUCH AS ROADSIDES, DITCH BANKS AND FENCE ROWS

KEEP LIVESTOCK OFF TREATED AREAS. AVOID DRIFT OF SPRAY TO ADJACENT CROPLAND.

APPLY WHEN WEEDS ARE YOUNG AND GROWING ACTIVELY. SPRAY UNTIL WEEDS ARE THOROUGHLY WET.

RETREATMENT MAY BE NECESSARY TO CONTROL PLANTS NOT THOROUGHLY SPRAYED OR PLANTS ARISING FROM DORMANT SEEDS AND NUTS OR OLD ROOTS.

FOR SPOT TREATMENT OF SMALL AREAS USE A KNAPSACK SPRAYER WITH A MEDIUM SIZE NOZZLE. A VERY COARSE NOZZLE OR WATERING CAN IS NOT RECOMMENDED FOR MAKING SPOT TREATMENTS.

MIX 1/2 CUP (4 FLUID OUNCES) AMROL LIQUID WEED KILLER PER GALLON OF WATER. ONE GALLON OF SPRAY SOLUTION IS USUALLY SUFFICIENT TO COVER 272 SQUARE FEET OF SQUARE FOOT THOROUGHLY INFESTED WITH WEEDS.

| Weeds          | Gals. AMROL LIQUID WEED KILLER | Gals. Water | Directions  |
|----------------|--------------------------------|-------------|---|
| Canada Thistle | 1                              | 50          | Spray when plants are young and growing actively. Do not mow treated plants. Do not fall plow or disturb thistle to be treated in the spring. In the far West, treat during the bloom stage.  |
| Field Bindweed | 1                              | 50          | Spray when plants are young and growing actively. Do not mow treated plants. Do not fall plow or disturb bindweed to be treated in the spring. In the far West, treat during the bloom stage. |
| Field Bindweed | 1                              | 100         | Apply after catkins are fully formed and up to frost. Do not disturb sprayed plants. Spot treat any regrowth.   |
| Field Bindweed | 1                              | 50          | Spray when plants are 5 inches tall and growing actively.   |
| Field Bindweed | 2                              | 50          | Treat from mid-summer to early fall when shoots are growing vigorously. Spot treat any green regrowth with a solution of 1 gal. AMROL LIQUID WEED KILLER per 50 gals. of water.               |
| Nutsedge       | 1                              | 50          | Spray when shoots are 4-5 inches tall and growing actively. Where practical, cultivate treated areas when plants become white. Retreat when dormant nuts germinate.                           |
| Field Bindweed | 1                              | 50          | Spray in spring when weeds are in the bud to bloom stage. Do not mow treated plants. Spot treat any regrowth.   |
| Field Bindweed | 1                              | 50          | In the East and Mid West, spray when horse manure is in the bud to bloom stage. In the far West, treat from mid-summer to early frost. If practical, plow 2 weeks after spraying.             |
| Field Bindweed | 1                              | 50          | Spray in spring when weeds are in the bud to bloom stage. Do not mow treated plants. Spot treat any regrowth.   |
| Field Bindweed | 1                              | 50          | Spray when alfalfa is growing vigorously.   |
| Field Bindweed | 1                              | 50          | Spray in spring when weeds are in the bud to bloom stage. Do not mow treated plants. Spot treat any regrowth.   |
| Field Bindweed | 1                              | 50          | Apply anytime when dock is actively growing, preferably before development of bloom stalks.   |



| Weeds  | Gals. AMROL LIQUID WEED KILLER | Gals. Water | Directions  |
|--|--------------------------------|-------------|---|
| Wild Blackberry, Rubus                                     | 1                              | 50          | Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line. |
| Honeysuckle, Lonicera                                      | 2                              | 50          | Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line. |
| Salmal Berry, Western Dewberry, Blackberry, Big Leaf maple | 2                              | 50          | Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line. |
| Wild cherry, Ash, Locust and Sumac                         | 1                              | 50          | Treat from the time foliage is fully developed until plants begin to go dormant. For effective kill, all leaves, stems and suckers must be thoroughly wet to ground line. |

| ATTACHES                     | AMOUNTS PER ACRE                               |             |  |             |
|------------------------------|--|-------------|--|-------------|
|                              | Ground Spraying GALS. AMROL LIQUID WEED KILLER | GALS. WATER | Aerial Spraying GALS. AMROL LIQUID WEED KILLER | GALS. WATER |
| North-Central Eastern States | 1  | 50          | 1  | 50          |
| Southwestern States          | 1  | 50          | 1  | 50          |
| West - Apply                 | 1  | 50          | 1  | 50          |
| Apply after                  | 1  | 50          | 1  | 50          |
| Apply after                  | 1  | 50          | 1  | 50          |
| Apply after                  | 1  | 50          | 1  | 50          |
| Apply after                  | 1  | 50          | 1  | 50          |
| Apply after                  | 1  | 50          | 1  | 50          |

DO NOT APPLY WHERE WATER WILL BE USED FOR IRRIGATING DRINKING OR OTHER DOMESTIC PURPOSES.