

TREATED AREAS MAY BE KILLED. KEEP OUT OF LAKES, STREAMS AND PONDS. DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTE. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

MAY BE POISONOUS IF SWALLOWED. HARMFUL IF INHALED OR ABSORBED THROUGH THE SKIN. DO NOT GET IN EYES OR ON SKIN. DO NOT INHALE SPRAY MIST. USE WITH PROTECTIVE CLOTHING AND A FACE MASK. AFTER HANDLING, WASH WITH PLENTY OF SOAP AND WATER. WASH CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE REUSING.

IF POISONING OCCURS, GET MEDICAL ATTENTION IMMEDIATELY. PROLONGED EXPOSURE WILL RESULT IN CHOLINESTERASE DEPRESSION.

NOTE TO PHYSICIAN: ATROPINE SULFATE IS ANTIDOTAL. 2-PAM IS ALSO ANTIDOTAL AND MAY BE USED IN CONJUNCTION WITH ATROPINE. DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

ETIES OF SORGHUM WHICH ARE SENSITIVE TO PHOSPHATES. DO NOT USE TRINOX LIQUID INSECTICIDE ON CROPS NOT LISTED IN THIS LABEL. DO NOT USE TRINOX LIQUID INSECTICIDE ON FOOD CROPS GROWN IN GREENHOUSES.

RECOMMENDED APPLICATIONS

CROP	INSECT	PINTS TRINOX LIQUID INSECTICIDE	REMARKS
FIELD CROPS			
Alfalfa Clover	Alfalfa Caterpillar	¾ to 1	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Make only 1 application per cutting. Allow 4 days between treatment and cutting or grazing for rates up to 1 pint per acre and a 7 day interval for rates above 1 pint per acre.
	Alfalfa webworm	½ to 2	
	Western yellow-striped armyworm	1	
	Beet armyworm Variegated cutworm	1 to 2	
	Lygus bugs Stink bugs Tarnished plant bug	2	
Barley Flax Oats Wheat	Armyworms Beet webworm Variegated cutworm	1 to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary but not within 21 days of harvest. Allow 3 days between treatment and use of treated straw as feed for dairy or meat animals.
	Bertha armyworm Diamondback moth	2	

	Western yellow-striped armyworm	
	Beet armyworm Southern garden Leafhopper	2
	Black flea-hopper complex Cotton leaf perforator Leaf roller Lygus bugs Stink bugs	2 to 3
	Salt-marsh-caterpillar	3
Sugar Beets	Beet webworm Variegated cutworm	1 to 2
	Dipterous leaf miners	2
	Alfalfa webworm Beet armyworm	2 to 3
	Salt-marsh caterpillar	3
SEED FIELD CROPS		
Alfalfa Clover	Armyworms Lygus bugs Stink bugs Variegated cutworm	2 to 3

ACCEPTED
 1 MAR 19 1975
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 EPA Reg. No.
 registered under

2749-138



Manufactured For:

ACETO CHEMICAL CO.
 AGRICULTURAL CHEMICALS DIVISION

126-02 NORTHERN BOULEVARD, FLUSHING,

TRINOX LIQUID INSECTICIDE

IS NOT SIGNIFICANTLY
H AS PARASITES, PRED-
ALLY WHEN APPLIED AT
TES PER ACRE. THIS AD-
LIQUID INSECTICIDE IS
ON-SELECTIVE INSECTI-

IS READILY WITH WATER
FOR USE IN ALL POWER-
AND AIRCRAFT SPRAY-
ED AMOUNT OF TRINOX
REQUIRED AMOUNT OF
FIED DOSAGE OF TRINOX
DUNT OF WATER NECES-
RAGE OF FOLIAGE. THE
USED WILL DETERMINE

INSECTICIDE ON VARI-
IE SENSITIVE TO PHOS-
LIQUID INSECTICIDE ON
EL. DO NOT USE TRINOX
ROPS GROWN IN GREEN-

APPLICATIONS

TRINOX LIQUID INSECTICIDE	REMARKS
1	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Make only 1 application per cutting. Allow 4 days between treatment and cutting or grazing for rates up to 1 pint per acre and a 7 day interval for rates above 1 pint per acre.
2	
2	

CROP	INSECT	PINTS TRINOX LIQUID INSECTICIDE	REMARKS
Corn	Armyworms	1 to 2	Apply specified dosage per acre in sufficient water for complete coverage. Make one application per season when plants are 3 to 12 inches high and direct the spray to the lower portions of plants and to the soil around the base of the plants. Do not use corn for food, feed, or ensilage within 28 days of treatment.
	Cutworms		
Cotton	Cotton flea-hopper	½ to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Do not pasture treated fields within 14 days of application. Do not apply within 7 days of picking. TRINOX applications may occasionally cause marginal leaf burn. NOTE: For control of light to moderate infestation of cotton leafhoppers, use ½ pint of TRINOX per acre. Higher rates may be required if heavy infestations or migrations occur.
	Cotton leaf-worm	1 to 2	
	Darkling ground beetle		
	Western yellow-striped army-worm		
	Beet armyworm	2	
	Southern garden Leafhopper		
Black flea-hopper complex	2 to 3		
Cotton leaf perforator			
Leaf roller	3		
Lygus bugs			
Stink bugs			
Sugar Beets	Salt-marsh-caterpillar	3	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 14 days of harvest.
	Beet webworm	1 to 2	
	Variiegated cut-worm		
	Dipterous leaf miners	2	
	Alfalfa web-worm	2 to 3	
Beet army-worm			
Salt-marsh caterpillar	3		

CROP	INSECT	PINTS TRINOX LIQUID INSECTICIDE	REMARKS
			forage; however, do not cut green crop for these purposes.
VEGATABLES			
Beans (Dry & Snap)	Western bean cutworm	1 to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 14 days of harvest.
	Armyworms	2 to 3	
	Dipterous leaf miners		
	Lygus bugs		
	Mexican bean beetle		
	Stink bugs		
	Variiegated cut-worms		
Lima Beans	Western bean cut worm	1 to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Do not apply more than twice per season nor within 14 days of harvest.
	Armyworms	2 to 3	
	Dipterous leaf-miners		
	Lygus bugs		
	Mexican bean beetle		
	Stink bugs		
	Variiegated cutworm		
Cowpeas (Southern Peas	Armyworms	2 to 3	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Apply twice per season. Do not apply within 14 days of harvest.
	Dipterous leaf miners		
Black-eyed Peas	2 to 3		
Peas			
Crowder Peas)			
	Imported cabbage worm		
	Lygus bugs		
	Mexican bean beetle		
	Stink bugs		
	Variiegated cutworm		
ORNAMENTALS			
Ornamental Flowers, Shrubs & Trees	Armyworms	2 to 3	Mix specified dosage in 100 gallons of water and spray plants until they are thoroughly wet. For use in hand type sprayers, mix 2 to 3 tablespoonsfull per gallon of water. Repeat applications as required. CAUTION: Injury to
	Bagworms		
	Climbing cut-worms		
	Dipterous leaf miners		
	Lygus bugs		
	Tarnished plant bug		
	Tobacco bud-		

IS SENSITIVE TO FROST
LIQUID INSECTICIDE ON
EL. DO NOT USE TRINOX
CROPS GROWN IN GREEN-

APPLICATIONS

RS Dx DID NCIDE	REMARKS
1	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Make only 1 application per cutting. Allow 4 days between treatment and cutting or grazing for rates up to 1 pint per acre and a 7 day interval for rates above 1 pint per acre.
2	
2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary but not within 21 days of harvest. Allow 3 days between treatment and use of treated straw as feed for dairy or meat animals.

	striped army-worm		application. Do not apply within 7 days of picking. TRINOX applications may occasionally cause marginal leaf burn. NOTE: For control of light to moderate infestation of cotton leafhoppers, use 1/2 pint of TRINOX per acre. Higher rates may be required if heavy infestations or migrations occur.
	Beet armyworm Southern garden Leafhopper	2	
	Black flea-hopper complex Cotton leaf perforator Leaf roller Lygus bugs Stink bugs	2 to 3	
	Salt-marsh-caterpillar	3	
Sugar Beets	Beet webworm Variegated cutworm	1 to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 14 days of harvest.
	Dipterous leaf miners	2	
	Alfalfa webworm Beet army-worm	2 to 3	
	Salt-marsh caterpillar	3	
SEED FIELD CROPS			
Alfalfa Clover	Armyworms Lygus bugs Stink bugs Variegated cutworm	2 to 3	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary up to 7 days before harvest. Chaff from alfalfa and clover seed crop may be used for feed or

Lima Beans	Mexican bean beetle Stink bugs Variegated cutworm	2 to 3	Apply 1 gallon per acre. Do not apply more than twice per season nor within 14 days of harvest.
Cowpeas (Southern Peas Black-eyed Peas Crowder Peas)	Armyworms Dipterous leaf miners Imported cabbage worm Lygus bugs Mexican bean beetle Stink bugs Variegated cutworm	2 to 3	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Apply twice per season. Do not apply within 14 days of harvest.
ORNAMENTALS			
Ornamental Flowers, Shrubs & Trees	Armyworms Bagworms Climbing cutworms Dipterous leaf miners Lygus bugs Tarnished plant bug Tobacco budworm Webworms	2 to 3	Mix specified dosage in 100 gallons of water and spray plants until they are thoroughly wet. For use in hand type sprayers, mix 2 to 3 table-spoons full per gallon of water. Repeat applications as required. CAUTION: Injury to foliage has occurred on certain varieties of carnations, hydrangias and zinnias. When large plantings of these flowers are to be sprayed for the first time, treat only a few plants and observe for 4 to 5 days before-spraying the entire plantings.
	Nantucket pine tip moth Zimmerman pine moth	2	

Manufactured For:

ACETO CHEMICAL CO., INC.

AGRICULTURAL CHEMICALS DIVISION

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368

ONOFF IS LIKELY TO
 TAMINATE WATER BY
 IENT OR DISPOSAL OF
 PRODUCT ONLY AS
 BEL.

SWALLOWED. HARM-
 ABSORBED THROUGH
 T IN EYES OR ON SKIN.
 RAY MIST. USE WITH
 G AND A FACE MASK.
 WASH WITH PLENTY OF
 WASH CONTAMINATED
 AND HOT WATER BE

GET MEDICAL AT-
 LY. PROLONGED EX-
 IN CHOLINESTERASE

ATROPINE SULFATE IS
 ALSO ANTIDOTAL AND
 UNCTION WITH ATRO-
 E EMPTY CONTAINER.
 TAINER BY PERFORA-
 ID BURYING IN A SAFE

LIQUID INSECTICIDE ON FOOD CROPS GROWN IN GREEN-
 HOUSES.

RECOMMENDED APPLICATIONS

CROP	INSECT	PINTS TRINOX LIQUID INSECTICIDE	REMARKS
FIELD CROPS			
Alfalfa Clover	Alfalfa Caterpillar	¾ to 1	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Make only 1 application per cutting. Allow 4 days between treatment and cutting or grazing for rates up to 1 pint per acre and a 7 day interval for rates above 1 pint per acre.
	Alfalfa webworm	½ to 2	
	Western yellow-striped armyworm	1	
	Beet armyworm Variegated cutworm	1 to 2	
	Lygus bugs Stink bugs Tarnished plant bug	2	
Barley Flax Oats Wheat	Armyworms Beet webworm Variegated cutworm	1 to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary but not within 21 days of harvest. Allow 3 days between treatment and use of treated straw as feed for dairy or meat animals.
	Bertha armyworm Diamondback moth	2	

Southern garden Le
 Black flea
 hopper co
 Cotton lea
 forator
 Leaf rolle
 Lygus bug
 Stink bug

Salt-marsh
 erpillar

Sugar Beets

Beet web
 Variegated
 worm

Dipterous
 miners

Alfalfa we
 worm
 Beet army
 worm

Salt-marsh
 erpillar

SEED
 FIELD CROPS

Alfalfa
 Clover

Armywor
 Lygus bug
 Stink bug
 Variegated
 cutworm

ACCEPTED

MAR 19 1975

Manufactured For:

PINTS TRINOX LIQUID INSECTICIDE	REMARKS
1 to 2	Apply specified dosage per acre in sufficient water for complete coverage. Make one application per season when plants are 3 to 12 inches high and direct the spray to the lower portions of plants and to the soil around the base of the plants. Do not use corn for food, feed, or ensilage within 28 days of treatment.
½ to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Do not pasture treated fields within 14 days of application. Do not apply within 7 days of picking. TRINOX applications may occasionally cause marginal leaf burn. NOTE: For control of light to moderate infestation of cotton leafhoppers, use ½ pint
1 to 2	
2 er	
2 to 3	

CROP	INSECT	PINTS TRINOX LIQUID INSECTICIDE	REMARKS
			forage; however, do not cut green crop for these purposes.
VEGATABLES			
Beans (Dry & Snap)	Western bean cutworm	1 to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Repeat as necessary. Do not apply within 14 days of harvest.
	Armyworms Dipterous leaf miners Lygus bugs Mexican bean beetle Stink bugs Variegated cutworms	2 to 3	
Lima Beans	Western bean cut worm	1 to 2	Apply specified dosage per acre using sufficient water for complete coverage but not less than 1 gallon per acre. Do not apply more than twice per season nor within 14 days of harvest.
	Armyworms Dipterous leaf-miners Lygus bugs Mexican bean beetle Stink bugs Variegated cutworm	2 to 3	
Cowpeas (Southern Peas Black-eyed	Armyworms Dipterous leaf miners Imported cab-	2 to 3	Apply specified dosage per acre using sufficient water for complete