

**DIPHACINONE  
RODENTICIDE  
ANTI-COAGULANT FOR RAT AND MICE CONTROL**

**ACTIVE INGREDIENT.**

DIPHACINONE (2-Diphenylacetyl-1, 3-indandione)	0.1%
INERT INGREDIENTS:	99.9%
TOTAL:	100.0%

EPA REG. NO 2749-139

NET WEIGHT 5 LBS

**CAUTION:** KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM HUMANS, WILDLIFE, DOMESTIC ANIMALS AND PETS. IF SWALLOWED BY HUMANS, DOMESTIC ANIMALS OR PETS, THIS MATERIAL MAY REDUCE THE CLOTTING ABILITY OF THE BLOOD AND CAUSE BLEEDING. IN SUCH CASE, INTRAVENOUS AND ORAL ADMINISTRATION OF VITAMIN K COMBINED WITH BLOOD TRANSFUSIONS ARE INDICATED AS IN THE CASE OF HEMORRHAGE CAUSED BY OVERDOSES OF BISHYDROXYCOUMARIN.

ANTIDOTE: GIVE A TABLESPOONFUL OF SALT IN A GLASS OF WARM WATER AND REPEAT UNTIL VOMIT FLUID IS CLEAR. CALL A PHYSICIAN IMMEDIATELY.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

**DIRECTIONS FOR USE:**

- (A) THOROUGHLY MIX ONE PART (BY WEIGHT) OF RODENTICIDE WITH NINETEEN PARTS OF DRY CEREAL BAIT.
- (B) PLACE BAIT IN LOCATIONS WHERE RATS AND MICE FEED, WATER OR TRAVEL.
- (C) BAIT PLACEMENTS ALONG WALLS, IN CORNERS, AND IN CONCEALED LOCATIONS WHERE RATS AND MICE FEED ARE DESIRABLE.
- (D) BAITS SHOULD BE PLACED IN AREAS INACCESSIBLE TO CHILDREN, PETS, AND DOMESTIC ANIMALS, OR IN TAMPER-PROOF BAIT BOXES.

**FOR RAT CONTROL:**

- (A) PLACE AT LEAST FOUR OUNCES OF BAIT AT EACH BAIT LOCATION
- (B) MAINTAIN AN UNINTERRUPTED SUPPLY OF BAIT IN THE STATIONS FOR APPROXIMATELY 10 DAYS.

**FOR MOUSE CONTROL:**

- (A) PLACE TABLESPOON AMOUNTS OF BAIT AT 8-10 FEET INTERVALS
- (B) MAINTAIN AN UNINTERRUPTED SUPPLY OF BAIT FOR AT LEAST 15 DAYS.

NOTE: IF BAIT MOLDS, DECOMPOSES OR BECOMES SOUR, REPLACE WITH FRESH MATERIALS.



Manufactured by

**ACETO CHEMICAL CO., INC.**  
AGRICULTURAL CHEMICALS DIV.

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368