

ACCEPTED

Jul 12, 1972

2749-142

FUNDEX
FUNGICIDE

FOR CONTROL OF SCAB AND RUST ON APPLES

ACTIVE INGREDIENTS:

Manganous Dimethyldithiocarbamate	48.0%
Manganous Benzothiazylmercaptide	2.4%
Thiram (Tetramethylthiuram disulfide)	12.4%
2,2' - Dithiobisbenzothiazole	2.2%

INERT INGREDIENTS:

	35.0%
TOTAL	100.0%

NET WEIGHT

EPA REG. NO. 2749-142

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

KEEP AWAY FROM FIRES AND SPARKS

THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS AND PONDS. DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTE.

MAY BE HARMFUL IF SWALLOWED OR INHALED. MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. AVOID BREATHING DUST OR SPRAY MIST. USE WITH ADEQUATE VENTILATION. AVOID CONTACT WITH EYES, SKIN OR CLOTHING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.


FUNDEX IS ESPECIALLY RECOMMENDED FOR USE IN AN ALL-SEASON PROGRAM ON RUSSET - SUSCEPTIBLE VARIETIES SUCH AS GOLDEN DELICIOUS, GRIMES GOLDEN AND JONATHON.

DIRECTIONS FOR USE

Apples: For early season control of scab and rust. Use FUNDEX in the pre-bloom, bloom period calyx and first cover sprays at the rate of 2 lbs. per 100 gallons of water.

Apples: For late season control of scab. Use in the second and third cover applications at the rate of 1½ lbs. per 100 gallons of water and in any subsequent applications at the rate of 1 lb. per 100 gallons of water. If disease conditions are severe, higher dosages up to 2 lbs. per 100 gallons of water may be required in later application.

If FUNDEX is applied as a concentrated spray, use ¾ of the above dosages per 100 gallons of water multiplied by the number of times the spray is concentrated.



Manufactured for

ACETO CHEMICAL CO., INC.

AGRICULTURAL CHEMICALS DIV.

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368