



ACTIVE INGREDIENT:  
DIMETHYL (2, 2, 2-TRICHLORO-1-HYDROXYETHYL)  
PHOSPHONATE (TRICHLORFON) 8%  
INERT INGREDIENTS: 92%  
TOTAL 100%

EPA REG. NO. 2749-147

NET CONTENTS \_\_\_\_\_

## WARNING:

KEEP OUT OF REACH OF CHILDREN.  
KEEP AWAY FROM DOMESTIC ANIMALS.  
DO NOT CONTAMINATE FEED OR FOOD.

THIS PRODUCT IS TOXIC TO FISH, WILDLIFE AND BIRDS. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.

MAY BE POISONOUS IF SWALLOWED. HARMFUL IF ABSORBED THROUGH THE SKIN. DO NOT GET IN EYES OR ON SKIN. WASH THOROUGHLY WITH SOAP AND WARM WATER AFTER HANDLING. WASH CONTAMINATED CLOTHING WITH SOAP AND HOT WATER BEFORE REUSE.

IF POISONING OCCURS, OBTAIN PROMPT MEDICAL AID. EXPOSURE MAY RESULT IN CHOLINESTERASE DEPRESSION.

## BOVINOX POUR ON CATTLE INSECTICIDE FOR EFFECTIVE CONTROL OF CATTLE GRUBS

NOTE TO PHYSICIAN: ATROPINE SULFATE IS ANTIDOTAL. 2-PAM IS ALSO ANTIDOTAL AND MAY BE ADMINISTERED IN CONJUNCTION WITH ATROPINE.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

### DIRECTIONS FOR USE:

BOVINOX POUR-ON CATTLE INSECTICIDE is a ready-to-use preparation that requires no further dilution before use on beef cattle and non-lactating dairy cattle. A single application placed on the backline of an animal usually gives effective control of cattle grubs and reduction of lice populations.

Apply BOVINOX POUR-ON solution at the rate of 1/2 fluid ounce per 100 pounds of body weight. Measure the correct amount to use in relation to the animal's weight.

### 1 GALLON OF BOVINOX POUR-ON CATTLE INSECTICIDE WILL TREAT

Weight of Animals (lbs.)	200	300	400	500	600	700	800	900	1000
Number of Animals	128	85	64	51	42	36	32	28	25

Pour the correct amount of solution uniformly along the animal's back. The BOVINOX POUR-ON solution may be dipped from any clean bucket or pail.

Treat as soon as possible after activity of heel fly has ceased and at least 6 weeks prior to expected appearance of grubs in the back. This is to avoid treatment at time

grubs are in the esophagus (gullet) or spinal canal. Some mild to moderate side reactions may occur if application is made while grubs are in the area of the esophagus or spinal canal. These reactions, which may be evidenced by salivation, bloat, urticaria and ataxia, are usually reversible, however, should reactions of this type occur, consult a veterinarian. Applications either before or after this stage or grub development will not result in these side reactions which are caused by substances released from the killed grubs.

### PROTECT FROM EXTREME HEAT AND PROTECT FROM SUNLIGHT

DO NOT TREAT OVER-HEATED ANIMALS.  
DO NOT TREAT LACTATING DAIRY CATTLE.  
DO NOT TREAT ANIMALS LESS THAN 3 MONTHS OLD.  
DO NOT TREAT ANIMALS THAT ARE SICK, CONVALESCENT OR STRESSED LIVESTOCK.

Do not treat dry dairy animals within 21 days of freshening. If freshening occurs within 21 days of treatment, do not use milk for food for the balance of the 21 day period.

Do not treat cattle within 21 days of slaughter.

Do not treat animals for 10 days before or after shipping or weaning, or after exposure to contagious or infectious diseases.

WARNING: Trichlorfon is a cholinesterase inhibitor. Do not use this product on animals simultaneously or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals. Consult veterinarian at first sign of adverse reaction. Atropine is antidote.



Manufactured for:

ACETO CHEMICAL CO., INC.  
AGRICULTURAL CHEMICALS DIV.

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368