

CHLOROBENZILATE 4E MITICIDE

ACTIVE INGREDIENT:	
ETHYL 4,4'-DICHLOROBENZILATE*	45.5%
XYLENE	43.5%
INERT INGREDIENTS	11.0%
TOTAL	100.0%

*EQUIVALENT TO 4 LBS. PER GALLON

NET CONTENTS 5 GALLONS

EPA REG. NO. 2749-155

WARNING: KEEP OUT OF REACH OF CHILDREN.

POISONOUS BY SKIN CONTACT, INHALATION OR SWALLOWING. RAPIDLY ABSORBED THROUGH THE SKIN. DO NOT GET IN EYES, SKIN OR ON CLOTHING. DO NOT BREATHE SPRAY MIST. WASH THOROUGHLY AFTER HANDLING AND BEFORE EATING OR SMOKING.

FISH CAUTION. THIS PRODUCT IS TOXIC TO FISH. DO NOT CONTAMINATE ANY BODY OF WATER BY DIRECT APPLICATION, CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES AND CONTAINERS.

DO NOT REUSE EMPTY CONTAINER. DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

FLAMMABLE MIXTURE. KEEP AWAY FROM HEAT OR OPEN FLAME. STORE AT TEMPERATURES ABOVE 32° F.

GENERAL INFORMATION

CHLOROBENZILATE 4E MITICIDE IS AN EMULSIFIABLE SOLUTION INTENDED FOR APPLICATION AFTER DILUTION WITH WATER. THE RATES BELOW ARE FOR DILUTE SPRAYS (RATE PER 100 GALS. OF WATER) UNLESS OTHERWISE STATED. IN GENERAL APPLICATION SHOULD BE MADE WHEN MITES FIRST APPEAR. FOR SEVERE INFESTATIONS, OR IF CONDITIONS ARE FAVORABLE FOR MITE BUILDUP, REPEATED APPLICATIONS WILL BE NECESSARY. DO NOT USE WITH HIGHLY ALKALINE MATERIALS SUCH AS LIME.

DIRECTIONS FOR USE

Agricultural Crops:

Almonds - European Red Mites, Pacific Spider Mites, Brown Mites (Clover Mites), Two-spotted Spider Mites. Use 1/2 - 3/4 pt. per 100 gals. of water and apply as full coverage sprays when mites first appear. Repeat applications as necessary. Do not apply within 21 days of harvest. Treated almond hulls may be used for feed for livestock.

NOTE: DO NOT APPLY TO KAPAREIL VARIETY OF ALMONDS SINCE INJURY MAY OCCUR. CHLOROBENZILATE 4E IS COMPATIBLE WITH MOST COMMONLY USED LIGHTWEIGHT SUMMER OILS.

Apples and Pears - Two-spotted Spider Mites, European Red Mites, Brown Mites (Clover Mites), McDaniel Mites. Apply 1/2 - 3/4 pt. per 100 gals. of water in full coverage sprays, or repeat applications after 7 - 10 days, particularly if the lower dosage is used. In concentrate applications apply 5-6 pts. per acre in sufficient water to assure thorough coverage. Applications should not be made before second cover spray.

NOTE: DO NOT APPLY TO DELICIOUS, JONATHAN, OR MCINTOSH APPLES SINCE FRUIT INJURY MAY OCCUR ON THESE VARIETIES.

DO NOT APPLY TO APPLES WITHIN 14 DAYS, OR TO PEARS WITHIN 7 DAYS OF HARVEST.

Cherries - Cherry Rust Mites: Apply 3/4 pt. per 100 gals. of water, after fruit harvest.

NOTE: DO NOT USE ON PLUMS, PRUNES OR PEACHES SINCE INJURY MAY OCCUR.

Citrus - Grapefruit, Oranges, Lemons, Limes, Tangelos, Tangerines, Kumquats:

Citrus Rust Mites: Apply 1/4 - 1/2 pt. per 100 gals. of water or 5 pts. per acre in sufficient water to assure thorough coverage when mites first appear. For severe infestations, or if conditions are favorable for mite buildup, repeated applications will be necessary.

Aerial Applications: Apply 1-1/2 pts. in 10 gallons of water per acre as a supplemental treatment.

Flat Mites, Six-Spotted Mites: Texas Citrus Fruit Mites: Apply 1/2 pt. per 100 gals. of water, or apply 5 pts. per acre in sufficient water to assure thorough coverage.

Citrus Red Mites: Purple Mites: Apply 1/2 pt. per 100 gals. of water, or apply 5 pts. per acre in sufficient water to assure thorough coverage for quick cleanup of this pest. For seasonal control either repeated applications or use in combinations with other miticides will be required.

Citrus Bud Mites, Pacific Spider Mites, Yuma Spider Mites: Apply 1/2 - 3/4 pt. per 100 gals. of water, or apply 5 - 7-1/2 pts. per acre in sufficient water to assure thorough coverage.

Cotton - Spider Mites (Atlantic, Pacific, Schoene, Two-spotted): Apply 1 quart per acre in sufficient water to assure thorough coverage, when leaves start to show red and yellow speckling. Direct spray so under sides of leaves receive thorough coverage.

Aerial Application: Apply 1 qt. in a minimum of 3 gals. of water per acre.

NOTE: DO NOT APPLY AFTER BOLLS BEGIN TO OPEN. DO NOT GRAZE OR FEED TREATED FORAGE OR GIN WASTES TO LIVESTOCK.

Melons - Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persian Melons:

Two-spotted Spider Mites: Apply 1 qt. per acre broadcast in sufficient water to assure thorough coverage. When treating young plants that do not require broadcast treatment, reduce the amount of CHLOROBENZILATE 4E in proportion to the area treated. For example, when melons are planted in 60 inch rows and vines 20 inches wide are treated, apply 1/3 qt. per acre.

Walnuts - European Red Mites: Apply 1/2 - 3/4 pt. per 100 gals. of water as a full coverage spray, or apply 6 - 8 pts. per acre in sufficient water to assure thorough coverage.

NOTE: DO NOT APPLY AFTER HUSKS BEGIN TO SPLIT OPEN.

Ornamentals and Lawns:

Holly - Southern Red Mites: Mix 1/2 - 3/4 pt. per 100 gals. of water and apply when mites are first noted. Repeat as necessary.

Spruce, Arborvitae, and Hemlock - Spruce Spider Mites: Mix 1/2 - 3/4 pt. per 100 gals. of water and apply when mites are first noted. Repeat as necessary.

Taxus and Azalea - Two-spotted Spider Mites: Mix 1/2 to 3/4 pt. per 100 gals. of water and apply when mites are first noted. Repeat as necessary.

Lawns and Shrubbery - Clover Mites: To control clover mites and prevent their migration to adjoining dwellings, mix 1/2 - 3/4 pt. per 100 gals. of water and apply as a drenching spray to infested lawn and shrubbery within 10 feet of dwellings. The foundation and exterior walls of dwellings should be similarly treated to the height of the first floor windows. Repeat application in 7-10 days if necessary.



Manufactured by

ACETO CHEMICAL CO., INC.