



cobex

SUPPLEMENTAL LABELING

Sunflower:

Apply preplant and incorporate COBEX at broadcast rates of 1-1/3 pints (1/3 lb/Acre) on coarse soils; 2 pints (1/2 lb/Acre) on medium soils; and 2-2/3 pints (2/3 lb/Acre) on fine soils.

1624-104

IMPORTANT

BEFORE USING COBEX, CAREFULLY READ THE COBEX LABEL (EPA Reg. No. 1624-104) AND FOLLOW THE PRECAUTIONS AND DIRECTIONS.

THIS BULLETIN CONTAINS NEW OR SUPPLEMENTAL DIRECTIONS FOR THE USE OF COBEX. USE THIS BULLETIN IN CONJUNCTION WITH THE REGULAR PACKAGE LABEL (EPA Reg. No. 1624-104-AA).



cobex

SUPPLEMENTAL LABELING

Soybeans - Midwest United States Only

COBEX at 3/8 lb (1-1/2 pints— per acre applied to medium-fine soils with less than 2.5% organic matter will control foxtail, crabgrass, pigweed and lambs-quarters.

To obtain control of smartweed and to substantially reduce populations of morningglory, velvetleaf, ragweed and teaweed, use 2 pints COBEX per acre.

ACCEPTED
JUN 23 1976
1624-104

IMPORTANT

BEFORE USING COBEX, CAREFULLY READ THE COBEX LABEL (EPA Reg. No. 1624-104) AND FOLLOW THE PRECAUTIONS AND DIRECTIONS.

THIS BULLETIN CONTAINS NEW OR SUPPLEMENTAL DIRECTIONS FOR THE USE OF COBEX. USE THIS BULLETIN IN CONJUNCTION WITH THE REGULAR PACKAGE LABEL (EPA Reg. No. 1624-104-AA).