

SOLEX 4G

WEED-KILLING FERTILIZER

FOR NON-CROPLAND USE

ACTIVE INGREDIENT:

m- (3,3-Dimethylureido phenyl) tert-
butylcarbamate

4.0%

INERT INGREDIENTS:

TOTAL 96.0%
100.0%

EPA REG. NO. _____

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

May be harmful if swallowed. May cause irritation of eyes, nose,
throat and skin. Avoid inhalation of dust. Avoid contact with
skin, eyes or clothing.

Keep away from domestic animals and foodstuffs.

Do not reuse empty container. Destroy empty container by perforating
or crushing and burying in a safe place.

Thoroughly clean spreading equipment before using for other purposes
(or do not use same spreading equipment for other purposes).

DIRECTIONS FOR USE

Apply SOLEX 4G just before or during the period of active growth of
the weeds to be controlled. Sufficient moisture from rainfall or
artificial means is necessary after application to carry the chemical
into the root zones. The time of treatment should precede the rainy
season. Where limited rainfall occurs during the active growth period,
this material will not provide satisfactory control of hard-to-kill
deep-rooted perennials.

BEST DOCUMENT AVAILABLE

100

60 to 100

140 to 200

300 to 400

USE THESE TABLES TO DETERMINE THE CORRECT PORT FOR YOUR PORTAL. THE PORTAL IS THE PORTAL THAT IS USED TO ACCESS THE PORTAL. THE PORTAL IS THE PORTAL THAT IS USED TO ACCESS THE PORTAL.

Manufactured Port:

Manufactured Port: 1000
Manufactured Port: 1000
Manufactured Port: 1000
Manufactured Port: 1000

BEST DOCUMENT AVAILABLE