

BROMOXYNIL WEED KILLER

FOR CONTROL OF BUCKWHEAT, PURPLE MUSTARD,
TARWEED, FIDDLENECK AND GROMWELL WEEDS IN WHEAT AND BARLEY

ACTIVE INGREDIENT:

Octanoic acid ester of Bromoxynil (3,5-dibromo-4-hydroxybenzonitrile)*

33.4%

INERT INGREDIENTS:

66.6%

TOTAL

100.0%

*Contains 2 lbs. of Active per gallon.

NET CONTENTS 1 GALLON
EPA REG. NO. 2749-179

CAUTION: KEEP OUT OF THE REACH OF CHILDREN. KEEP AWAY FROM DOMESTIC ANIMALS AND FOODSTUFFS.

THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS OR PONDS.

MAY BE HARMFUL IF SWALLOWED. MAY CAUSE IRRITATION OF EYES, NOSE, THROAT AND SKIN. AVOID INHALATION OF SPRAY MIST. AVOID CONTACT WITH SKIN, EYES OR CLOTHING.

IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES, GET MEDICAL ATTENTION.

DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

THOROUGHLY CLEAN SPRAY EQUIPMENT WITH A SUITABLE CHEMICAL BEFORE USING FOR OTHER PURPOSES.

AVOID SPRAY DRIFT INTO NEIGHBORING CROPS AND PASTURES.

DO NOT STORE NEAR FERTILIZERS, SEEDS, INSECTICIDES, OR FUNGICIDES.

DO NOT CONTAMINATE WATER USED FOR IRRIGATION OR DOMESTIC PURPOSES.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

Apply 1½ pints of BROMOXYNIL per acre after wheat or barley emerge (2-3 leaf) and before weeds are past 3-4 leaf stage.

Use 2 pints of BROMOXYNIL per acre when weeds are between 3-4 leaf stage and 6-8 inches in height, but before flower formation.

Where fiddleneck of tarweed is a problem or where weeds are growing during a drought, the higher rate is suggested. If lower dosages are to be used, consult local agricultural authorities for the rate required to control weeds in your particular area.

Do not apply to wheat or barley during or after boot stage.

Spray in enough water for good distribution. Use 10 or more gallons per acre with ground sprayers or 5 or more gallons per acre with aerial sprayers.

Among the weeds controlled, when label directions are followed, are the following:

Fiddleneck	(Amsinckia intermedia)
Tarweed	(Hemizonia spp.)
Wild Buckwheat	(Polygonum convolvulus)
Tartary Buckwheat	(Fagopyrum tataricum)
Blue (purple)	(Chotidpots tenella)
Tumble (Jim Hill) mustard	(Sisymbrium altissimum)
Wild mustard	(Brassica kaber)
Field pennycress, fanweed	(Thlaspi arvense)
Corn gromwell	(Lithospermum arvense)
Cow Cockle	(Saponaria vaccaria)
Lambsquarters	(Chenopodium album)
London Rocket	(Sisymbrium irio)
Shepherdspurse	(Capsella bursa-pastoris)
Silver-leaf nightshade	(Solanum elaeagnifolium)
Groundsel	(Senecio vulgaris)

This treatment will not control established perennial broad-leaf weeds such as Canada thistle, bindweed or horsetail rush.



Manufactured for

ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICALS DIVISION

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368