


# ETHION 25WP INSECTICIDE



**ACTIVE INGREDIENT:**

Ethion (0,0,0',0'-Tetraethyl S,  
S'-methylene bisphosphorodithioate

25%

**INERT INGREDIENTS:**

75%

TOTAL 100%

EPA REG. NO: 2749-213  
NET WEIGHT \_\_\_\_\_

**WARNING: KEEP OUT OF THE REACH OF CHILDREN.**

MAY BE FATAL IF SWALLOWED. ABSORBED THROUGH SKIN. DO NOT BREATHE VAPOR, DUST OR SPRAY MIST. AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WASH THOROUGHLY AFTER HANDLING. AVOID STORAGE NEAR FOOD AND FEED PRODUCTS.

THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL. THIS PRODUCT IS TOXIC TO BEES AND SHOULD NOT BE APPLIED WHEN BEES ARE ACTIVELY VISITING THE AREA.

DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREAS TREATED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.

Rinse equipment and DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

APPLE (40)	Apple Rust Mite Clover (Brown Almond) Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 8-10 lbs. per acre	DO NOT EXCEED 4 applications during fruiting period.
APPLE (30)	Apple Rust Mite Clover (Brown Almond) Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 8-10 lbs. per acre	DO NOT EXCEED 2 applications during fruiting period.
APPLE (20)	Apple Rust Mite Clover (Brown Almond) Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 8-10 lbs. per acre	DO NOT EXCEED 4 applications during fruiting period.

DO NOT USE on Astrachen, Davey, Duchess, Margaret Pratt, Melba, Wealthy Williams Red (Red Williams), Yellow Transparent or any varieties maturing at or before early McIntosh apples as injury is likely to occur.

CHERRY	Clover (Brown Almond) Mite European Red Mite McDaniel Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 6-10 lbs. per acre	Use as a post-harvest treatment only.
GRAPE (30) (Eastern Use Only)	Grape Berry Moth Grape Leafhopper Two-spotted Spider Mite	3 lbs. per 100 gals. of water finished spray per acre 9 lbs. per acre	Apply a single application during the first first week in August.
PEACH (30) NECTARINE (30)	Clover (Brown Almond) Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 5-7 lbs. per acre	DO NOT EXCEED 2 applications during fruiting period.
PEAR (60)	Codling Moth  Pear Psylla	2 lbs. per 100 gals. of water 10-20 lbs. per acre 1½ to 2 lbs. per 100 gals. of water 8-20 lbs. per acre	DO NOT EXCEED 3 applications after the first cover spray.
PEAR (30)	Clover (Brown Almond) Mite	1 lb. per 100 gals. of water	DO NOT EXCEED 3 applications after the

LIMES (21)	Six-
ORANGES (0)	Tea
TANGELOS (0)	Black
TANGERINES (0)	Brown
	California
	Chandler
	Florida
	Gloves
	Purple
	Sweet
	Yellow
	Clara
	FOR

BEANS (4)	LY
Succulent and Dry	
BEANS (2)	Pot
	ho
	Be
	Be
	Ma
	Be
	ri
	Sp
BEAN SEED	Sh

REC  
 Sep-28-1972  
 2749-213

# ETHION 25% INSECTICIDE

ACTIVE INGREDIENT:  
 Ethion (0,0,0',0'-Tetraethyl S,  
 S'-methylene bisphosphorodithioate) 25%

INERT INGREDIENTS: 75%

TOTAL 100%

EPA REG. NO: 2749-213  
 NET WEIGHT

## WARNING: KEEP OUT OF THE REACH OF CHILDREN.

MAY BE FATAL IF SWALLOWED. ABSORBED THROUGH SKIN. DO NOT BREATHE VAPOR, DUST OR SPRAY MIST. AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WASH THOROUGHLY AFTER HANDLING. AVOID STORAGE NEAR FOOD AND FEED PRODUCTS.

THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL. THIS PRODUCT IS TOXIC TO BEES AND SHOULD NOT BE APPLIED WHEN BEES ARE ACTIVELY VISITING THE AREA.

DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREAS TREATED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.

Rinse equipment and DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

### DIRECTIONS FOR USE:

Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in ( ) following crop. Do not use in undiluted form.

APPLE (60)	Codling Moth	2 lbs. per 100 gals. of water 16-20 lbs. per acre	DO NOT EXCEED 4 applications during fruiting period.
APPLE (40)	Codling Moth	2 lbs. per 100 gals. of water 16-20 lbs. per acre	DO NOT EXCEED 2 applications during fruiting period.

APPLE (40)	Apple Rust Mite Clover (Brown) Almond Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 8-10 lbs. per acre	DO NOT EXCEED 4 applications during fruiting period.	LIME ORANGE TANG
APPLE (30)	Apple Rust Mite Clover (Brown) Almond Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 8-10 lbs. per acre	DO NOT EXCEED 2 applications during fruiting period.	
APPLE (20)	Apple Rust Mite Clover (Brown) Almond Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 8-10 lbs. per acre	DO NOT EXCEED 4 applications during fruiting period.	
DO NOT USE on Astrachen, Davey, Duchess, Margaret Pratt, Melba, Wealthy Williams Red (Red Williams), Yellow Transparent or any varieties maturing at or before early McIntosh apples as injury is likely to occur.				
CHERRY	Clover (Brown) Almond Mite European Red Mite McDaniel Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 6-10 lbs. per acre	Use as a post-harvest treatment only.	
GRAPE (30) (Eastern Use Only)	Grape Berry Moth Grape Leafhopper Two-spotted Spider Mite	3 lbs. per 100 gals. of water of finished spray per acre 9 lbs. per acre	Apply a single application during the first week in August.	
PEACH (30) NECTARINE (30)	Clover (Brown) Almond Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 5-7 lbs. per acre	DO NOT EXCEED 2 applications during fruiting period.	BEAN SUGAR and BEAN
PEAR (60)	Codling Moth  Pear Psylla	2 lbs. per 100 gals. of water 10-20 lbs. per acre 1 1/2 to 2 lbs. per 100 gals. of water 8-20 lbs. per acre	DO NOT EXCEED 3 applications after the first cover spray.	
PEAR (30)	Clover (Brown) Almond Mite European Red Mite Pear Rust Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 5-10 lbs. per acre 8-10 lbs. for Rust Mite	DO NOT EXCEED 3 applications after the first cover spray.	BEAN TREA
PLUMS (21) PRUNES (21)	Clover (Brown) Almond Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 5-7 lbs. per acre	DO NOT EXCEED 3 applications during the fruiting period.	CUCU SUMM SQUA
PLUMS (45) PRUNES (45)	European Fruit Lecanium	1 1/2 lbs. per 100 gals. of water 7-10 lbs. per acre	DO NOT EXCEED 2 applications during fruiting period.	EGGP
STRAWBERRY (2)	Strawberry Leaf Roller Two-spotted Spider Mite	4 lbs. per acre 2 lbs. per acre		MELC
GRAPEFRUIT (0) LEMONS (21)	Citrus Red Mite Citrus Rust Mite			



Manufactured for  
**ACETO CHE**  
 AGRICULT  
 176 02 NORTHERN BOUL

# THION 25WP INSECTICIDE

SEED 4 ap-  
ing fruiting

SEED 2 ap-  
ing fruiting

SEED 4 ap-  
ing fruiting

ba, Wealthy  
turing at or

harvest  
ly.

ap-  
ing the first  
August.

SEED 2 ap-  
ing  
id.

SEED 3 ap-  
for the

SEED 3 ap-  
for the  
ay.

SEED 3  
during the  
ed

SEED 2  
during  
ed

ured for

LIMES (21)	Six-spotted Mite		
ORANGES (10)	Texas Citrus Mite		
TANGELOS (10)	Black Scale		
<b>TANGERINES (0)</b>			
	Brown Soft Scale		
	Calif. Red Scale		
	Chaff Scale		
	Florida Red Scale		
	Glover Scale		
	Purple Scale		
	Snow Scale		
	Yellow Scale		
	Citrus Whitefly		
	<b>FOR MITES:</b>	1-1.5 lbs. per 100 gals. of water combined with 0.5-0.7% summer oil (Florida & Texas areas) or 1 to 1.5 lbs. per 100 gals. of water combined with 1-2 gals. summer oil (West Coast)	<b>DO NOT APPLY</b> more than 2000 gals. of spray per acre. On lemons and limes, do not apply more than 1X per season. On tangerines do not apply more than 2X per season. On grapefruit, oranges, tangelos and tangerines do not repeat applications within 90 days of the previous treatment. <b>DO NOT APPLY</b> oil within 3 weeks of a sulfur application. <b>DO NOT USE</b> oil on navel oranges
	<b>FOR WHITEFLY:</b>	WHERE SNOW SCALE is a problem, spray trunk, limbs and twigs thoroughly. Apply shortly after Whiteflies have laid their eggs.	

## VEGETABLES

BEANS (4) Succulent and Dry	Lygus Bugs	4 lbs. per acre.	<b>DO NOT FEED</b> treated bean foliage to livestock.
BEANS (2)	Potatoe Leaf hopper	2 lbs. per acre.	
	Bean Leaf Beetle		<b>DO NOT FEED</b> treated bean foliage to livestock.
	Mexican Bean Beetle	1-2 lbs. per acre	
	Two spotted Spider Mite		
BEAN SEED TREATMENT	Seed Corn Maggot	4 ozs. per 100 lbs. of seed	Mix and apply to seed as a slurry treatment. Combine with a suitable fungicide for seed treatment if desired. Do not use treated seed for food, feed or oil purposes.
CUCUMBER (0) SUMMER SQUASH (0)	Serpentine Leaf Miner	2 lbs. per acre	
	Two spotted Mite		
EGGPLANT (0)	Two spotted Mite	1-2 lbs. per acre	
MELONS (0)	Melon leaf Miner	2 lbs. per acre	
	Serpentine Leaf Miner		
	Two spotted Spider Mite		

ONION SEED TREATMENT	Onion Maggot		Treat seed according to the number of feet of row planted per acre. On upland soil mix 4 lbs. with the amount of seed to be planted in 1 acre of rows 12" apart. Mix 3 lbs. with the amount of seed to be planted in 1 acre of rows 15" apart and 2 lbs. on seeds to be planted in 1 acre rows 24" apart. On muck or heavy type soil increase dosage by 50-100 percent.
PEPPER (0) PIMENTO (0)	Serpentine Leaf Miner	2 lbs. per acre	
	Two-spotted Mite	1-2 lbs. per acre	
TOMATOES (2)	Tropical Mite	2 lbs. per acre	
	Two spotted Spider mite		
	Leaf Miner		

## OTHER CROPS

ALMONDS (30)	Clover (Brown Almond) Mite	1 lbs. per 100 gals. of water	<b>DO NOT EXCEED</b> 1 application during fruiting period.
	European Red Mite	5-8 lbs. per acre	
	Two-spotted Spider Mite		
CHESTNUTS FILBERTS PECANS	Aphids	10 lbs. per acre	<b>DO NOT EXCEED</b> two applications during fruiting period. On Chestnuts and pecans <b>DO NOT APPLY</b> after husk split. On Filberts, <b>DO NOT APPLY</b> when nut is visible. <b>DO NOT GRAZE</b> livestock in treated groves.
	Mites		
WALNUTS	European Red Mite	6-10 lbs. per acre	<b>DO NOT EXCEED</b> two applications during the fruiting period. <b>DO NOT APPLY</b> after husk split.
	Two spotted Spider Mite		
	Walnut Aphid		
	Walnut Husk Fly		
SORGHUM (grown for seed & grain)	Sorghum Midge	1-2 lbs. per acre	<b>DO NOT EXCEED</b> 3 applications. <b>DO NOT</b> harvest or feed forage within 30 days after treatment. Some leaf burning may occur on phosphate sensitive varieties.
	Sorghum Mite		
COTTON	Cotton leafworm	Use 2 lbs. in sufficient water to treat 1 acre	<b>DO NOT APPLY</b> after bolls open. <b>DO NOT</b> graze dairy or meat animals in treated fields. Workers entering treated fields within 24 hours of application should wear protective clothing.
ORNAMENTALS (Out of doors)	Two spotted Spider Mite	1 lb. per 100 gals	Application to Anderson Yew may cause some needle drop.

TO CHEMICAL CO. INC.  
AGRICULTURAL CHEMICALS  
NORTHERN BOULEVARD

**SPECIFIED ON THIS LABEL. THIS PRODUCT IS TOXIC TO BEES AND SHOULD NOT BE APPLIED WHEN BEES ARE ACTIVELY VISITING THE AREA.**

**DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT APPLY WHEN WEATHER CONDITIONS FAVOR DRIFT FROM AREAS TREATED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR DISPOSAL OF WASTES.**

**Rinse equipment and DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.**

**DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.**

**DIRECTIONS FOR USE:**

**Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe use limitations. If insect control is required beyond the following use patterns, supplement the control program with other suitable pesticides. Observe days interval between last application and harvest indicated by number in ( ) following crop. Do not use in undiluted form.**

<b>APPLE (60)</b>	<b>Codling Moth</b>	<b>2 lbs. per 100 gals. of water 16-20 lbs. per acre</b>	<b>DO NOT EXCEED 4 applications during fruiting period.</b>
<b>APPLE (40)</b>	<b>Codling Moth</b>	<b>2 lbs. per 100 gals. of water 16-20 lbs. per acre</b>	<b>DO NOT EXCEED 2 applications during fruiting period.</b>

GRA  
(East  
Use C  
  
PEAC  
NECT  
(3  
  
PEAF  
  
PEAF  
  
PLUM  
PRU  
  
PLUM  
PRU  
  
STRA  
  
GRA  
LEM

# ETHION 25WP INSECTICIDE

DO NOT EXCEED 4 applications during fruiting period.

DO NOT EXCEED 2 applications during fruiting period.

DO NOT EXCEED 4 applications during fruiting period.

Att, Melba, Wealthy varieties maturing at or

as a post-harvest treatment only.

Apply a single application during the first week in August.

DO NOT EXCEED 2 applications during fruiting period.

DO NOT EXCEED 3 applications after the first cover spray.

DO NOT EXCEED 3 applications after the first cover spray.

LIMES (21)  
ORANGES (0)  
TANGELOS (0)

Six-spotted Mite  
Texas Citrus Mite  
Black Scale

TANGERINES (0)  
Brown Soft Scale  
Calif. Red Scale  
Chaff Scale  
Florida Red Scale  
Glover Scale  
Purple Scale  
Snow Scale  
Yellow Scale  
Citrus Whitefly  
FOR MITES:

1-1.5 lbs. per 100 gals. of water combined with 0.5-0.7% summer oil (Florida & Texas areas) or 1 to 1.5 lbs. per 100 gals. of water combined with 1-2 gals. summer oil (West Coast)  
WHERE SNOW SCALE is a problem, spray trunk, limbs and twigs thoroughly  
Apply shortly after Whiteflies have laid their eggs.

DO NOT APPLY more than 2000 gals. of spray per acre. On lemons and limes, do not apply more than 1X per season. On tangerines do not apply more than 2X per season. On grapefruit, oranges, tangelos and tangerines do not repeat applications within 90 days of the previous treatment. DO NOT APPLY oil within 3 weeks of a sulfur application. DO NOT USE oil on navel oranges.

FOR WHITEFLY:

## VEGETABLES

BEANS (1) Succulent and Dry	Lygus Bugs	4 lbs. per acre.	DO NOT FEED treated bean foliage to livestock.
BEANS (2)	Potatoe Leaf-hopper Bean Leaf Beetle Mexican Bean Beetle Two-spotted Spider Mite	2 lbs. per acre. 1-2 lbs. per acre	DO NOT FEED treated bean foliage to livestock.
BEAN SEED TREATMENT:	Seed Corn Maggot	4 ozs. per 100 lbs. of seed	Mix and apply to seed as a slurry treatment. Combine with a suitable fungicide.

ONION SEED TREATMENT

Onion Maggot

Treat seed according to the number of feet of row planted per acre. On upland soil mix 4 lbs. with the amount of seed to be planted in 1 acre of rows 12" apart. Mix 3 lbs. with the amount of seed to be planted in 1 acre of rows 18" apart and 2 lbs. on seeds to be planted in 1 acre rows 24" apart. On muck or heavy-type soil increase dosage by 50-100 percent.

PEPPER (0) PIMENTO (0)	Serpentine Leaf Miner Two-spotted Mite	2 lbs. per acre 1-2 lbs. per acre
TOMATOES (2)	Tropical Mite Two-spotted Spider mite Leaf Miner	2 lbs. per acre

## OTHER CROPS

ALMONDS (30)	Clover (Brown Almond) Mite European Red Mite Two-spotted Spider Mite	1 lbs. per 100 gals. of water 5-8 lbs. per acre	DO NOT EXCEED 1 application during fruiting period.
CHESTNUTS FILBERTS PECANS	Aphids Mites	10 lbs. per acre	DO NOT EXCEED two applications during fruiting period. On Chestnuts and pecans DO NOT APPLY after husk split. On Filberts, DO NOT APPLY when nut is visible. DO NOT GRAZE livestock in treated groves.
WALNUTS	European Red Mite Two-spotted Spider Mite Walnut Aphid Walnut Husk Fly	6-10 lbs. per acre	DO NOT EXCEED two applications during the fruiting period. DO NOT APPLY after husk split.
SORGHUM (grown for	Sorghum Midges Sorghum Mite	1-2 lbs. per acre	DO NOT EXCEED 3 applications. DO NOT harvest or feed forage

as a post-harvest  
treatment only.

Apply a single ap-  
plication during the first  
week in August.

DO NOT EXCEED 2 ap-  
plications during  
fruiting period.

DO NOT EXCEED 3 ap-  
plications after the  
cover  
spray.

DO NOT EXCEED 3 ap-  
plications after the  
cover spray.

DO NOT EXCEED 3 ap-  
plications during the  
fruiting period.

DO NOT EXCEED 2 ap-  
plications during  
fruiting period.

**FOR WHITEFLY:**

**SCALE** is a problem,  
spray trunk, limbs  
and twigs thoroughly  
Apply shortly after  
Whiteflies have laid  
their eggs.

**VEGETABLES**

<b>BEANS (4)</b> Succulent and Dry	Lygus Bugs	4 lbs. per acre.	<b>DO NOT FEED</b> treated bean foliage to livestock.
<b>BEANS (2)</b>	Potatoe Leaf- hopper Bean Leaf Beetle Mexican Bean Beetle Two-spotted Spider Mite	2 lbs. per acre.  1-2 lbs. per acre	<b>DO NOT FEED</b> treated bean foliage to livestock.
<b>BEAN SEED TREATMENT:</b>	Seed Corn Maggot	4 ozs. per 100 lbs. of seed	Mix and apply to seed as a slurry treatment. Combine with a suitable fungicide for seed treatment if de- sired. Do not use treated seed for food, feed or oil purposes.
<b>CUCUMBER (0) SUMMER SQUASH (0)</b>	Serpentine Leaf Miner Two-spotted Mite	2 lbs. per acre	
<b>EGGPLANT (0)</b>	Two-spotted Mite	1-2 lbs. per acre	
<b>MELONS (0)</b>	Melon leaf Miner Serpentine Leaf Miner Two-spotted Spider Mite	2 lbs. per acre	

**OTHER CROPS**

<b>ALMONDS (30)</b>	Clover (Brown Almond) Mite European Red Mite Two-spotted Spider Mite	1 lb. per 100 gals. of water 5-8 lbs. per acre	<b>DO NOT EXCEED 1 ap- plication</b> during fruiting period.
<b>CHESTNUTS FILBERTS PECANS</b>	Aphids Mites	10 lbs. per acre	<b>DO NOT EXCEED two ap- plications</b> during fruiting period. On Chestnuts and pecans <b>DO NOT APPLY</b> after husk split. On Filberts, <b>DO NOT APPLY</b> when nut is visible. <b>DO NOT GRAZE</b> livestock in treated groves.
<b>WALNUTS</b>	European Red Mite Two-spotted Spider Mite Walnut Aphid Walnut Husk Fly	6-10 lbs. per acre	<b>DO NOT EXCEED two ap- plications</b> during the fruiting period. <b>DO NOT APPLY</b> after husk split.
<b>SORGHUM</b> (grown for seed & grain)	Sorghum Midge Sorghum Mite	1-2 lbs. per acre	<b>DO NOT EXCEED 3 ap- plications. DO NOT</b> harvest or feed forage within 30 days after treatment. Some leaf burning may occur on phosphate sensitive varieties.
<b>COTTON</b>	Cotton leafworm	Use 2 lbs. in sufficient water to treat 1 acre.	<b>DO NOT APPLY</b> after bolls open. <b>DO NOT</b> graze dairy or meat animals in treated fields. Workers entering treated fields within 24 hours of application should wear protective clothing.
<b>ORNAMENTALS</b> (Out-of-doors)	Two-spotted Spider Mite	1 lb. per 100 gals.	Application to An- derson Yew may cause some needle drop.

Manufactured for:

**DETO CHEMICAL CO., INC.**  
AGRICULTURAL CHEMICALS DIV.

202 NORTHERN BOULEVARD, FLUSHING, N.Y. 11367