

MANEB 8 DUST
POTATO SEED TREATER
FUNGICIDE

ACTIVE INGREDIENT:

Maneb (Manganese ethylenebisdithiocarbamate) 8.0%

INERT INGREDIENTS:

92.0%
TOTAL 100.0%

EPA REG. NO. 2749-222

NET WEIGHT 50 LBS.

IMPORTANT

STORE IN A COOL, DRY PLACE. DO NOT ALLOW TO BECOME WET OR OVER-HEATED IN STORAGE. KEEP CONTAINER CLOSED WHEN NOT IN USE. DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

CONTROLS FUSARIUM SEED-PIECE DECAY, THE MOST COMMON SEED- AND SOIL-BORNE FUNGUS DISEASE OF POTATOES.

HELPS PREVENT SEED-PIECE ROT UNDER ADVERSE SPROUTING CONDITIONS (COLD AND WET, OR DRY SOIL CONDITIONS).

PROTECTS SEED WHICH IS HELD FOR PLANTING.

DIRECTIONS FOR USE:

POTATO SEED-PIECE: Cutting and Treating -- Before planting, cutting machines or knives, trays, tables, barrels or sacks, equipment, trucks and planters should be thoroughly cleaned and sanitized. If not sprouted, subject seed potatoes to temperature of 50° to 60° F. for about 10 to 15 days before handling them.

Dosage and Application: Use 1 lb. of MANEB 8 DUST per 100 lbs. of cut seed-pieces. Apply it so that the entire surface of the seed-piece is thoroughly covered by the fungicide dust. Place treated seed in clean, disease-free containers or store in bulk at proper temperature and humidity. Prevent inhalation of dust by use of exhaust fans during treatment.

The cut and treated seed may be planted immediately, but preferably should be held for a week or more at cool temperature and high relative humidity. This holding period promotes suberization and healing of the cut surfaces. Good suberization reduces the susceptibility of the seed-piece to infection by other disease organisms, especially in cold, wet soil.

Do not use excess treated seed-pieces for food or feed.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

AVOID INHALING DUST -- AVOID CONTACT WITH SKIN.
AVOID CONTAMINATION OF FEED AND FOOD.
STORE IN A COOL, DRY PLACE.
PROTECT FROM EXCESSIVE HEAT.

MAY CAUSE IRRITATION TO NOSE, THROAT, EYES AND SKIN.
DO NOT BREATHE DUST. IN CASE OF SKIN CONTACT, WASH WITH WATER; FOR EYES, WASH COPIOUSLY WITH WATER AND GET MEDICAL ATTENTION.

THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS OR PONDS. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.



Manufactured For

ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICALS DIV.

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368

POTATO SEED TREATER

FUNGICIDE

IMPORTANT

ACTIVE INGREDIENT:

Maneb (Manganese ethylenebisdithiocarbamate)

8.0%

INERT INGREDIENTS:


92.0%

TOTAL

100.0%

EPA REG. NO. 2749-222

NET WEIGHT 50 LBS.



STORE IN A COOL, DRY PLACE. DO NOT ALLOW TO BECOME WET OR OVER-HEATED IN STORAGE. KEEP CONTAINER CLOSED WHEN NOT IN USE. DO NOT REUSE EMPTY CONTAINER. DESTROY EMPTY CONTAINER BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE.

CONTROLS FUSARIUM SEED-PIECE DECAY, THE MOST COMMON SEED- AND SOIL-BORNE FUNGUS DISEASE OF POTATOES.

HELPS PREVENT SEED-PIECE ROT UNDER ADVERSE SPROUTING CONDITIONS (COLD AND WET, OR DRY SOIL CONDITIONS).

PROTECTS SEED WHICH IS HELD FOR PLANTING.

DIRECTIONS FOR USE:

POTATO SEED-PIECE: Cutting and Treating — Before planting, cutting machines or knives, trays, tables, barrels or sacks, equipment, trucks and planters should be thoroughly cleaned and sanitized. If not sprouted, subject seed potatoes to temperature of 50° to 60° F. for about 10 to 15 days before handling them.

Dosage and Application: Use 1 lb. of MANEB 8 DUST per 100 lbs. of cut seed-pieces. Apply it so that the entire surface of the seed-piece is thoroughly covered by the fungicide dust. Place treated seed in clean, disease-free containers or store in bulk at proper temperature and humidity. Prevent inhalation of dust by use of exhaust fans during treatment.

The cut and treated seed may be planted immediately, but preferably should be held for a week or more at cool temperature and high relative humidity. This holding period promotes suberization and healing of the cut surfaces. Good suberization reduces the susceptibility of the seed-piece to infection by other disease organisms, especially in cold, wet soil.

Do not use excess treated seed-pieces for food or feed.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.

AVOID INHALING DUST — AVOID CONTACT WITH SKIN.
AVOID CONTAMINATION OF FEED AND FOOD.
STORE IN A COOL, DRY PLACE.
PROTECT FROM EXCESSIVE HEAT.

MAY CAUSE IRRITATION TO NOSE, THROAT, EYES AND SKIN.
DO NOT BREATHE DUST. IN CASE OF SKIN CONTACT, WASH WITH WATER; FOR EYES, WASH COPIOUSLY WITH WATER AND GET MEDICAL ATTENTION.

THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS OR PONDS. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.



Manufactured For:

ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICALS DIV.

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368