


DICHLOBENIL 10-G AQUATIC WEED KILLER

ACTIVE INGREDIENT:
Dichlobenil (2,6-Dichlorobenzonitrile)
INERT INGREDIENTS:



EPA REG. NO: 2749-255
NET WEIGHT: 50 lbs.

10.0%
90.0%
TOTAL 100.0%

CAUTION: KEEP OUT OF REACH OF CHILDREN

HARMFUL IF SWALLOWED. DO NOT BREATHE DUST. DO NOT ALLOW CONTACT WITH EYES OR ON SKIN. USE WITH CAUTION. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. KEEP OUT OF REACH OF CHILDREN. DO NOT STORE WITH SEED, FEED OR FOOD PRODUCTS

DO NOT REUSE EMPTY CONTAINER. DESTROY BY BURYING IN A SAFE PLACE.

DO NOT APPLY TO WATER WHICH WILL BE USED FOR CROP IRRIGATION, FOR LIVESTOCK WATER OR FOR HUMAN CONSUMPTION. DO NOT USE FISH FROM TREATED WATER FOR FOOD OR FEED FOR A PERIOD OF 90 DAYS AFTER APPLICATION. DO NOT USE IN COMMERCIAL FISH OR SHELLFISH WATER. CONSULT YOUR STATE FISH GAME DEPARTMENT BEFORE APPLYING DICHLLOBENIL 10-G TO PUBLIC WATERS.

DICHLLOBENIL 10-G IS RECOMMENDED FOR THE CONTROL OF SUBMERGED AQUATIC WEEDS IN NON-FLOWING WATER, SUCH AS PONDS, RESERVOIRS AND LAKES. DICHLLOBENIL 10-G APPLIED BEFORE WEED GROWTH BEGINS WILL GIVE SEASON LONG CONTROL OF SEVERAL SPECIES OF WEEDS INCLUDING THE FOLLOWING:

ELODEA
NORTHERN WATERMILFOIL
NAIADS
COONTAIL
CHARA
PONDWEEDS (Potamogeton spp)

DICHLLOBENIL 10-G APPLIED BEFORE WEED EMERGENCE PREVENTS WEED GROWTH. THIS AVOIDS THE NEED TO CONTROL EXISTING STANDS OF VEGETATION AND THE ACCOMPANYING RISK OF FISH LOSS. DICHLLOBENIL 10-G DOES NOT HARM FISH WHEN USED AS DIRECTED.

DIRECTIONS

Apply DICHLLOBENIL 10-G during early spring either to exposed pond bottoms or through the water. DICHLLOBENIL 10-G is active through the soil rather than through the water, therefore, uniform distribution of the granules over the entire area to be treated is essential. It is particularly suited for use as a partial pond treatment, treatment around boat docks, and swimming areas.

TREATMENT OF EXPOSED BOTTOMS AND SHORELINES: Apply DICHLLOBENIL 10-G at a rate of 70-100 pounds per acre. Complete and uniform coverage of the area, whether it be applied to the bottom or into the water is necessary for control. Use the lower rate if the soil is wet at the time of application, or if the water is less than one foot deep. In the case of treatment after drawdown, the pond should be refilled promptly.

TREATMENT THROUGH WATER: Distribute DICHLLOBENIL 10-G uniformly over the surface of the water using a rate of 100-150 lbs. per acre of surface area. The lower rate should be used in shallow water less than 2 feet deep. The higher rate in deep water over 5-6 feet. Application should be made during periods of calm when wave action is at a minimum. Use of a non-motorized boat is recommended to avoid excessive water turbulence.

DICHLLOBENIL 10-G should be spread using an applicator. If areas to be treated are small, satisfactory application can be made by hand, throwing the granules with a small scoop. DICHLLOBENIL 10-G is designed to sink rapidly in water.



Manufactured for:

ACETO CHEMICAL CO., INC.
AGRICULTURAL CHEMICALS DIV.

126-02 NORTHERN BOULEVARD, FLUSHING, N.Y. 11368